


► Performance on the pneumonia mortality measure by hospital characteristics: **safety-net status, teaching status, and urban or rural location.**

The Centers for Medicare & Medicaid Services (CMS) periodically investigates select hospital characteristics that may impact a hospital's performance on the following mortality measure: hospital-level 30-day risk-standardized mortality rate (RSMR) following pneumonia [1]. The pneumonia mortality measure includes Medicare fee-for-service (FFS) and Veterans Health Administration (VA) beneficiaries aged 65 or older [2]. The pneumonia mortality measure assesses death from any cause within 30 days of a hospital admission for pneumonia, regardless of whether the patient dies while still in the hospital or after discharge from the hospital [2]. The pneumonia mortality measure has been publicly reported on [Hospital Compare](#) since 2008 and has been included in the Hospital Value-Based Purchasing (HVBP) Program since 2013 [3].

**FIGURE I** Distributions of hospital RSMRs (%) for pneumonia by safety-net status, July 2011-June 2014.



Variation in RSMRs reflects differences in performance among hospitals; lower RSMRs suggest better quality, and higher RSMRs suggest worse quality. To understand the impact of hospital safety-net status, teaching status, and urban or rural location, we examined RSMRs among hospitals with these characteristics with 25 or more qualifying admissions. Therefore, we evaluated the pneumonia RSMRs for a total of 4,165 hospitals by comparing 1,206 safety-net hospitals against 2,959 non-safety-net hospitals, 1,126 teaching hospitals against 3,039 non-teaching hospitals, and 3,155 urban hospitals against 1,010 rural hospitals for the July 2011 – June 2014 reporting period.


Safety-net hospitals are defined as those committed to caring for populations without stable access to care, specifically public hospitals or private hospitals with a Medicaid caseload greater than one standard deviation above their respective state's mean private hospital Medicaid caseload [4]. Teaching Hospitals provide post-graduate education for physicians completing residency and fellowship [4]. Urban and rural hospitals are defined by hospital self-identification [4].

Prepared for CMS by Yale New Haven Health Services Corporation (YNHHSC) Center for Outcomes Research and Evaluation (CORE) September 2015

HOSPITAL CHARACTERISTICS

To ensure accurate assessment of each hospital, the pneumonia mortality measure uses a statistical model to adjust for key differences in patient risk factors that are clinically relevant and that have a strong relationship with the mortality outcome [2].

**FIGURE 2** Distributions of hospital RSMRs (%) for pneumonia by teaching status, July 2011-June 2014.




**TABLE I** Distributions of hospital RSMRs (%) for pneumonia overall, by safety-net status, teaching status, and urban or rural location, July 2011-June 2014.

	Pneumonia RSMR (%)						
	Overall; n=4165	Safety-net hospitals; n=1206	Non-safety-net hospitals; n=2959	Teaching hospitals; n=1126	Non-teaching hospitals; n=3039	Urban hospitals; n=3155	Rural hospitals; n=1010
Maximum	20.3	20.3	19.8	19.9	20.3	20.3	19.3
90%	13.9	14.0	13.8	13.5	14.0	13.8	14.0
75%	12.6	12.9	12.5	12.2	12.7	12.5	12.9
Median (50%)	11.5	11.8	11.3	11.1	11.6	11.3	11.8
25%	10.5	10.9	10.3	10.0	10.7	10.3	10.9
10%	9.6	10.1	9.5	9.2	9.8	9.5	10.1
Minimum	6.9	7.5	6.9	7.1	6.9	6.9	7.7

Prepared for CMS by Yale New Haven Health Services Corporation (YNHHSC) Center for Outcomes Research and Evaluation (CORE) September 2015

The median pneumonia RSMR for all hospitals was 11.5% (interquartile range [IQR]: 10.5%-12.6%; Table 1). The median pneumonia RSMR for safety-net hospitals was 11.8% (IQR: 10.9%-12.9%) and for non-safety-net hospitals was 11.3% (IQR: 10.3%-12.5%; Figure 1 and Table 1). The median pneumonia RSMR for teaching hospitals was 11.1% (IQR: 10.0%-12.2%) and for non-teaching hospitals was 11.6% (IQR: 10.7%-12.7%; Figure 2 and Table 1). The median pneumonia RSMR for urban hospitals was 11.3% (IQR: 10.3%-12.5%) and for rural hospitals was 11.8% (IQR: 10.9%-12.9%; Figure 3 and Table 1).

**FIGURE 3** Distributions of hospital RSMRs (%) for pneumonia by urban or rural location, July 2011-June 2014.



Safety-net hospitals had a median pneumonia RSMR that was 0.5 percentage points higher than non-safety-net hospitals, teaching hospitals had a median pneumonia RSMR that was 0.5 percentage points lower than non-teaching hospitals, and urban hospitals had a median pneumonia RSMR that was 0.5 percentage points lower than rural hospitals.

1. Medicare Hospital Quality Chartbook 2014: Performance Report on Outcome Measures. Prepared by Yale New Haven Health Services Corporation Center for Outcomes Research and Evaluation for the Centers for Medicare and Medicaid Services 2014; <http://www.cms.gov/Medicare/Quality-Initiatives-Patient-Assessment-Instruments/HospitalQualityInits/Downloads/Medicare-Hospital-Quality-Chartbook-2014.pdf>. Accessed 16 June 2015.
2. Dorsey K, Grady J, Desai N, et al. 2015 Condition-Specific Measures Updates and Specifications Report Hospital-Level 30-Day Risk-Standardized Mortality Measures: Acute Myocardial Infarction – Version 9.0, Heart Failure – Version 9.0, Pneumonia – Version 9.0, Chronic Obstructive Pulmonary Disease – Version 4.0, Stroke – Version 4.0; <https://www.qualitynet.org/dcs/ContentServer?c=Page&pagename=QnetPublic%2FPage%2FQnetTier4&cid=1163010421830>. Accessed 26 June 2015.
3. “Hospital Inpatient Prospective Payment Systems for Acute Care Hospitals, Final Rule.” Federal Register / 22 August 2014; <http://federalregister.gov/a/2014-18545>. Accessed 16 June 2015.
4. AHA Annual Survey Database Fiscal Year 2013; <http://www.ahadataviewer.com/book-cd-products/aha-survey/>. Accessed 26 June 2015.

Prepared for CMS by Yale New Haven Health Services Corporation (YNHHSC) Center for Outcomes Research and Evaluation (CORE) September 2015