


SOCIODEMOGRAPHIC STATUS

► Performance on the pneumonia readmission measure: **Hospitals that serve high and low proportions of African-American patients.**

The Centers for Medicare & Medicaid Services (CMS) periodically investigates select hospital practices that may impact a hospital's performance on the following readmission measure: hospital-level 30-day risk-standardized readmission rate (RSRR) following pneumonia [1]. The pneumonia readmission measure includes Medicare fee-for-service (FFS) and Veterans Health Administration (VA) beneficiaries aged 65 or older [2]. The pneumonia readmission measure assesses the occurrence of unplanned readmission for any cause within 30 days after discharge from hospitalization for pneumonia [2]. The pneumonia readmission measure has been publicly reported on [Hospital Compare](#) since 2009 and has been included in the Hospital Readmissions Reduction Program (HRRP) since 2012 [3].

FIGURE I Distributions of pneumonia RSRRs (%) for hospitals with the lowest and highest proportions of African-American patients, July 2011-June 2014.



Prepared for CMS by Yale New Haven Health Services Corporation (YNHHSC) Center for Outcomes Research and Evaluation (CORE) September 2015

SOCIODEMOGRAPHIC STATUS

Variation in RSRRs reflects differences in performance among hospitals; lower RSRRs suggest better quality, and higher RSRRs suggest worse quality. To understand the impact of caring for African-American patients, we examined RSRRs among hospitals with high and low proportions of African-American patients. Therefore, we compared the pneumonia RSRRs for the 831 hospitals with the lowest overall proportion of African-American Medicare FFS patients (0% of a hospital's Medicare FFS patients) to the 423 hospitals with the highest proportion of African-American Medicare FFS patients ($\geq 22.1\%$ of a hospital's Medicare FFS patients) for the July 2011 – June 2014 reporting period. Hospitals with the lowest and highest proportions of African-American patients are designated as those that fall within the lowest and highest deciles of all hospitals with 25 or more qualifying discharges, respectively. The proportion of African-American Medicare FFS patients for each hospital was determined using the Medicare Part A Inpatient Claims from 2013. All hospitals with 0% African-American patients were included in the lowest decile. To ensure accurate assessment of each hospital, the pneumonia readmission measure uses a statistical model to adjust for key differences in patient risk factors that are clinically relevant and that have a strong relationship with the readmission outcome [2]. Please note that VA hospitals are not included in this analysis.

TABLE I Distributions of pneumonia RSRRs (%) for hospitals with the lowest and highest proportions of African-American patients, July 2011-June 2014.

	Pneumonia RSRR (%)	
	Lowest proportion (0%) African-American patients; n=831	Highest proportion ($\geq 22.1\%$) African-American patients; n=423
Maximum	20.8	21.5
90%	17.7	18.9
75%	17.1	18.1
Median (50%)	16.6	17.4
25%	16.2	16.7
10%	15.8	16.2
Minimum	13.4	14.5

The median pneumonia RSRR for hospitals with the highest proportion of African-American patients was 17.4% (interquartile range [IQR]: 16.7%-18.1%). The median pneumonia RSRR for hospitals with the lowest proportion of African-American patients was 16.6% (IQR: 16.2%-17.1%; Figure 1 and Table 1).

Hospitals with the lowest proportion of African-American patients had a median pneumonia RSRR that was 0.8 percentage points lower than hospitals with the highest proportion.

1. Medicare Hospital Quality Chartbook 2014: Performance Report on Outcome Measures. Prepared by Yale New Haven Health Services Corporation Center for Outcomes Research and Evaluation for the Centers for Medicare and Medicaid Services 2014; <http://www.cms.gov/Medicare/Quality-Initiatives-Patient-Assessment-Instruments/HospitalQualityInits/Downloads/Medicare-Hospital-Quality-Chartbook-2014.pdf>. Accessed 16 June 2015.

2. Dorsey K, Grady J, Desai N, et al. 2015 Condition-Specific Measures Updates and Specifications Report Hospital-Level 30-Day Risk-Standardized Readmission Measures: Acute Myocardial Infarction – Version 8.0, Heart Failure – Version 8.0, Pneumonia – Version 8.0, Chronic Obstructive Pulmonary Disease – Version 4.0, Stroke – Version 4.0; <https://www.qualitynet.org/dcs/ContentServer?c=Page&pagename=QnetPublic%2FPage%2FQnetTier4&cid=1219069855841>. Accessed 26 June 2015.

3. “Hospital Inpatient Prospective Payment Systems for Acute Care Hospitals, Final Rule.” Federal Register / 22 August 2014; <http://federalregister.gov/a/2014-18545>. Accessed 16 June 2015.

Prepared for CMS by Yale New Haven Health Services Corporation (YNHHC) Center for Outcomes Research and Evaluation (CORE) September 2015