


SOCIODEMOGRAPHIC STATUS

► Performance on the acute myocardial infarction mortality measure: **Hospitals that serve high and low proportions of African-American patients.**

The Centers for Medicare & Medicaid Services (CMS) periodically investigates select hospital practices that may impact a hospital's performance on the following mortality measure: hospital-level 30-day risk-standardized mortality rate (RSMR) following acute myocardial infarction (AMI) [1]. The AMI mortality measure includes Medicare fee-for-service (FFS) and Veterans Health Administration (VA) beneficiaries aged 65 or older [2]. The AMI mortality measure assesses the occurrence of death for any cause within 30 days after hospital admission for AMI [2]. The AMI mortality measure has been publicly reported on [Hospital Compare](#) since 2007 and has been included in the Hospital Value-Based Purchasing (HVBP) Program since 2013 [3].

FIGURE I Distributions of AMI RSMRs (%) for hospitals with the lowest and highest proportions of African-American patients, July 2011-June 2014.



Prepared for CMS by Yale New Haven Health Services Corporation (YNHHSC) Center for Outcomes Research and Evaluation (CORE) September 2015

SOCIODEMOGRAPHIC STATUS

Variation in RSMRs reflects differences in performance among hospitals; lower RSMRs suggest better quality, and higher RSMRs suggest worse quality. To understand the impact of caring for African-American patients, we examined RSMRs among hospitals with high and low proportions of African-American patients. Therefore, we compared the AMI RSMRs for the 244 hospitals with the lowest overall proportion of African-American Medicare FFS patients (0% of a hospital's Medicare FFS patients) to the 245 hospitals with the highest overall proportion of African-American Medicare FFS patients ($\geq 23.7\%$ of a hospital's Medicare FFS patients) for the July 2011 – June 2014 reporting period. Hospitals with the lowest and highest proportions of African-American patients are designated as those that fall within the lowest and highest deciles of all hospitals with 25 or more qualifying admissions, respectively. The proportion of African-American Medicare FFS patients for each hospital was determined using the Medicare Part A Inpatient Claims from 2013. All hospitals with 0% African-American patients were included in the lowest decile. To ensure accurate assessment of each hospital, the AMI mortality measure uses a statistical model to adjust for key differences in patient risk factors that are clinically relevant and that have a strong relationship with the mortality outcome [2]. Please note that VA hospitals are not included in this analysis.

TABLE I Distributions of AMI RSMRs (%) for hospitals with the lowest and highest proportions of African-American patients, July 2011-June 2014.

	AMI RSMR (%)	
	Lowest proportion (0%) African-American patients; n=244	Highest proportion ($\geq 23.7\%$) African-American patients; n=245
Maximum	17.5	17.9
90%	16.0	15.8
75%	15.2	15.1
Median (50%)	14.3	14.2
25%	13.5	13.4
10%	12.7	12.7
Minimum	11.5	10.4

The median AMI RSMR for hospitals with the highest proportion of African-American patients was 14.2% (interquartile range [IQR]: 13.4%-15.1%). The median AMI RSMR for hospitals with the lowest proportion of African-American patients was 14.3% (IQR: 13.5%-15.2%; Figure 1 and Table 1).

Hospitals with the lowest proportion of African-American patients had a median AMI RSMR that was 0.1 percentage points higher than that of hospitals with the highest proportion.

1. Medicare Hospital Quality Chartbook 2014: Performance Report on Outcome Measures. Prepared by Yale New Haven Health Services Corporation Center for Outcomes Research and Evaluation for the Centers for Medicare and Medicaid Services 2014; <http://www.cms.gov/Medicare/Quality-Initiatives-Patient-Assessment-Instruments/HospitalQualityInits/Downloads/Medicare-Hospital-Quality-Chartbook-2014.pdf>. Accessed 16 June 2015.

2. Dorsey K, Grady J, Desai N, et al. 2015 Condition-Specific Measures Updates and Specifications Report Hospital-Level 30-Day Risk-Standardized Mortality Measures: Acute Myocardial Infarction – Version 9.0, Heart Failure – Version 9.0, Pneumonia – Version 9.0, Chronic Obstructive Pulmonary Disease – Version 4.0, Stroke – Version 4.0; <https://www.qualitynet.org/dcs/ContentServer?c=Page&pagename=QnetPublic%2FPage%2FQnetTier4&cid=1163010421830>. Accessed 26 June 2015.

3. “Hospital Inpatient Prospective Payment Systems for Acute Care Hospitals, Final Rule.” Federal Register / 22 August 2014; <http://federalregister.gov/a/2014-18545>. Accessed 16 June 2015.

Prepared for CMS by Yale New Haven Health Services Corporation (YNHHC) Center for Outcomes Research and Evaluation (CORE) September 2015