

|  |
|--|
| <b>OASIS ITEM</b>  |
| <p><b>(M1600)</b> Has this patient been treated for a <b>Urinary Tract Infection</b> in the past 14 days?</p> <p> <input type="checkbox"/> 0 - No<br/> <input type="checkbox"/> 1 - Yes<br/> <input type="checkbox"/> NA - Patient on prophylactic treatment<br/> <input type="checkbox"/> UK - Unknown         </p>   |
| <b>ITEM INTENT</b>   |
| Identifies treatment of urinary tract infection during the past 14 days.   |
| <b>TIME POINTS ITEM(S) COMPLETED</b>   |
| Start of care<br>Resumption of care<br>Discharge from agency – not to inpatient facility   |
| <b>RESPONSE—SPECIFIC INSTRUCTIONS</b>  |
| <ul style="list-style-type: none"> <li>The term “past fourteen days” is the two-week period immediately preceding the start/resumption of care or discharge. This means that for purposes of counting the 14-day period, the date of admission is day 0 and the day immediately prior to the date of admission is day 1. For example, if the patient’s SOC date is August 20, any treatment for a UTI occurring on or after August 6 would be considered.</li> <li>Unknown is not an option at discharge from agency.</li> <li>Select Response 0 – No, if patient has not been treated for a UTI within the past two weeks, including if the patient had symptoms of a UTI or a positive culture for which the physician did not prescribe treatment, or the treatment ended more than 14 days ago.</li> <li>Select Response 1 – Yes, when the patient has been prescribed an antibiotic within the past 14 days specifically for a confirmed or suspected UTI.</li> <li>Select Response 1 – Yes, if the patient is on prophylactic treatment <u>and</u> develops a UTI.</li> <li>Select Response NA – if the patient is on prophylactic treatment to prevent UTIs.</li> </ul> |
| <b>DATA SOURCES / RESOURCES</b>  |
| <ul style="list-style-type: none"> <li>Patient/caregiver interview</li> <li>Physician orders</li> <li>Review of health history</li> <li>Referral information</li> <li>Physician</li> <li>Medication list</li> </ul>  |

|  |
|--|
| <b>OASIS ITEM</b>  |
| <p><b>(M1610) Urinary Incontinence or Urinary Catheter Presence:</b></p> <p><input type="checkbox"/> 0 - No incontinence or catheter (includes anuria or ostomy for urinary drainage) [ <i>Go to M1620</i> ]</p> <p><input type="checkbox"/> 1 - Patient is incontinent</p> <p><input type="checkbox"/> 2 - Patient requires a urinary catheter (i.e., external, indwelling, intermittent, suprapubic) [ <i>Go to M1620</i> ]</p>  |
| <b>ITEM INTENT</b>   |
| Identifies presence of urinary incontinence or condition that requires urinary catheterization of any type, including intermittent or indwelling. The etiology (cause) of incontinence is not addressed in this item.  |
| <b>TIME POINTS ITEM(S) COMPLETED</b>   |
| <p>Start of care</p> <p>Resumption of care</p> <p>Follow-up</p> <p>Discharge from agency - not to inpatient facility</p>   |
| <b>RESPONSE—SPECIFIC INSTRUCTIONS</b>  |
| <ul style="list-style-type: none"> <li>• Select Response 0 if the patient has anuria or an ostomy for urinary drainage (e.g., an ileal conduit), or if the patient has a urinary diversion that is pouched (ileal conduit, urostomy, ureterostomy, nephrostomy), with or without a stoma</li> <li>• Select Response 1 if the patient is incontinent AT ALL (i.e., “occasionally,” “only when I sneeze,” “sometimes I leak a little bit,” etc.).</li> <li>• Select Response 1 if the patient is incontinent or is dependent on a timed-voiding program. Timed voiding is defined as scheduled toileting assistance or prompted voiding to manage incontinence based on identified patterns. Time voiding is a compensatory strategy; it does not cure incontinence.</li> <li>• Select Response 2 if a catheter or tube is utilized for drainage (even if catheterizations are intermittent).</li> <li>• Select Response 2 if the patient requires the use of a urinary catheter for any reason (e.g., retention, post-surgery, incontinence). Select Response 2 and follow the skip pattern if the patient is <u>both</u> incontinent and requires a urinary catheter.</li> <li>• A leaking urinary drainage appliance is not incontinence.</li> <li>• Select Response 2 if a catheter was inserted during the comprehensive assessment.</li> <li>• If a catheter was discontinued during the comprehensive assessment or if a catheter is both inserted and discontinued during the comprehensive assessment, Response 0 or 1 would be appropriate, depending on whether or not the patient is continent.</li> <li>• Assessment strategies: Review the urinary elimination pattern as you take the health history. Does the patient admit having difficulty controlling the urine, or is he/she embarrassed about needing to wear a pad so as not to wet on clothing? Do you have orders to change a catheter? Is your stroke patient using an external catheter? Be alert for an odor of urine, which might indicate there is a problem with bladder sphincter control. If the patient receives aide services for bathing and/or dressing, ask for input from the aide (at follow-up assessment). This information can then be discussed with the patient. Urinary incontinence may result from multiple causes, including physiologic reasons, cognitive impairments, or mobility problems.</li> </ul> |

**DATA SOURCES / RESOURCES (cont'd for OASIS Item M1610)**

- Patient/caregiver interview
- Observation
- Physical assessment
- Physician orders
- Review of health history
- Referral information

|  |
|--|
| <b>OASIS ITEM</b>  |
| <p><b>(M1615) When does Urinary Incontinence occur?</b></p> <p> <input type="checkbox"/> 0 - Timed-voiding defers incontinence<br/> <input type="checkbox"/> 1 - Occasional stress incontinence<br/> <input type="checkbox"/> 2 - During the night only<br/> <input type="checkbox"/> 3 - During the day only<br/> <input type="checkbox"/> 4 - During the day and night         </p>  |
| <b>ITEM INTENT</b>   |
| Identifies when the urinary incontinence occurs.   |
| <b>TIME POINTS ITEM(S) COMPLETED</b>   |
| Start of care<br>Resumption of care<br>Discharge from agency - not to inpatient facility   |
| <b>RESPONSE—SPECIFIC INSTRUCTIONS</b>  |
| <ul style="list-style-type: none"> <li>• Select Response 0 if timed-voiding defers incontinence. Timed voiding determines the patient's pattern for voiding and schedules toileting to prevent episodes of leaking. The patient can self-schedule toileting or the caregiver can prompt or bring the patient to the toilet. Time voiding is a compensatory strategy; it does not cure incontinence. If timed voiding does not defer incontinence, do not select Response 0.</li> <li>• Select Response 1 – Occasional stress incontinence - when the patient is unable to prevent escape of relatively small amounts of urine when coughing, sneezing, laughing, lifting, moving from sitting to standing position, or other activities (stress), which increase abdominal pressure.</li> <li>• If urinary incontinence happens with regularity or in other circumstances than those described in the definition of stress incontinence, determine when the incontinence usually occurs and select Response 2, 3, or 4 as appropriate.</li> <li>• Select Response 2 – During the night only – when the patient's incontinence occurs while the patient is sleeping at night.</li> <li>• Select Response 3 – During the day only – when the patient's incontinence occurs while the patient is up/awake during the day. Includes incontinence during daytime naps.</li> <li>• Select Response 4 – During the day and night – when the patient is incontinent when sleeping at night and up/awake during the day.</li> </ul> |
| <b>DATA SOURCES / RESOURCES</b>  |
| <ul style="list-style-type: none"> <li>• Patient/caregiver interview</li> <li>• Observation</li> <li>• Physical assessment</li> <li>• Review of health history</li> <li>• Referral information</li> </ul>  |

|   |
|---|
| <b>OASIS ITEM</b>   |
| <p><b>(M1620) Bowel Incontinence Frequency:</b></p> <ul style="list-style-type: none"> <li><input type="checkbox"/> 0 - Very rarely or never has bowel incontinence</li> <li><input type="checkbox"/> 1 - Less than once weekly</li> <li><input type="checkbox"/> 2 - One to three times weekly</li> <li><input type="checkbox"/> 3 - Four to six times weekly</li> <li><input type="checkbox"/> 4 - On a daily basis</li> <li><input type="checkbox"/> 5 - More often than once daily</li> <li><input type="checkbox"/> NA - Patient has ostomy for bowel elimination</li> <li><input type="checkbox"/> UK - Unknown</li> </ul>  |
| <b>ITEM INTENT</b>  |
| Identifies how often the patient experiences bowel incontinence. Refers to the frequency of a symptom (bowel incontinence), not to the etiology (cause) of that symptom. This item does <u>not</u> address treatment of incontinence or constipation (e.g., a bowel program).   |
| <b>TIME POINTS ITEM(S) COMPLETED</b>  |
| <p>Start of care</p> <p>Resumption of care</p> <p>Follow-up</p> <p>Discharge from agency - not to inpatient facility</p>  |
| <b>RESPONSE—SPECIFIC INSTRUCTIONS</b>   |
| <ul style="list-style-type: none"> <li>• Responses are arranged in order of least to most frequency of bowel incontinence.</li> <li>• Response 4 – On a daily basis – indicates that the patient experiences bowel incontinence once per day.</li> <li>• Response NA is used if patient has an ostomy for bowel elimination.</li> <li>• Unknown is not an option at follow-up or discharge.</li> <li>• Assessment strategies: Review the bowel elimination pattern as you take the health history. Observe the cleanliness around the toilet when you are in the bathroom. Note any visible evidence of soiled clothing. Ask the patient if she/he has difficulty controlling stools, has problems with soiling clothing, uncontrollable diarrhea, etc. The patient's responses to these items may make you aware of an as yet unidentified problem that needs further investigation. If the patient is receiving aide services, question the aide about evidence of bowel incontinence at follow-up time points. This information can then be discussed with the patient. Incontinence may result from multiple causes, including physiologic reasons, mobility problems, or cognitive impairments.</li> </ul> |
| <b>DATA SOURCES / RESOURCES</b>   |
| <ul style="list-style-type: none"> <li>• Patient/caregiver interview</li> <li>• Observation</li> <li>• Physical assessment</li> <li>• Review of health history</li> <li>• Referral information</li> </ul>   |

| OASIS ITEM   |
|--|
| <p><b>(M1630) Ostomy for Bowel Elimination:</b> Does this patient have an ostomy for bowel elimination that (within the last 14 days): a) was related to an inpatient facility stay, <u>or</u> b) necessitated a change in medical or treatment regimen?</p> <p><input type="checkbox"/> 0 - Patient does <u>not</u> have an ostomy for bowel elimination.</p> <p><input type="checkbox"/> 1 - Patient's ostomy was <u>not</u> related to an inpatient stay and did <u>not</u> necessitate change in medical or treatment regimen.</p> <p><input type="checkbox"/> 2 - The ostomy <u>was</u> related to an inpatient stay or <u>did</u> necessitate change in medical or treatment regimen.</p>  |
| ITEM INTENT  |
| Identifies whether the patient has an ostomy for bowel elimination and, if so, whether the ostomy was related to a recent inpatient stay or caused a change in medical treatment plan.   |
| TIME POINTS ITEM(S) COMPLETED  |
| Start of care<br>Resumption of care<br>Follow-up   |
| RESPONSE—SPECIFIC INSTRUCTIONS   |
| <ul style="list-style-type: none"> <li>Applies to any type of ostomy for bowel elimination (e.g., colostomy, ileostomy). This item only addresses bowel ostomies, not other types of ostomies (e.g., urinary ostomies, tracheostomies).</li> <li>If an ostomy has been reversed, then the patient does <u>not</u> have an ostomy at the time of assessment.</li> <li>If patient does not have an ostomy for bowel elimination, select Response 0 – Patient does <u>not</u> have an ostomy for bowel elimination.</li> <li>If the patient does have an ostomy for bowel elimination, determine whether the ostomy was related to an inpatient stay or necessitated a change in the medical or treatment regimen within the last 14 days.</li> <li>The term “past fourteen days” is the two-week period immediately preceding the start/resumption of care or follow-up assessment. This means that for purposes of counting the 14-day period, the date of admission/assessment is day 0 and the day immediately prior to the date of admission/assessment is day 1. For example, if the patient’s SOC date is August 20, any ostomy related to an inpatient stay or requiring medical or treatment regimen change that occurred on or after August 6 would be considered.</li> </ul> |
| DATA SOURCES / RESOURCES   |
| <ul style="list-style-type: none"> <li>Patient/caregiver interview</li> <li>Physician orders</li> <li>Review of health history</li> <li>Referral information</li> <li>Physician</li> <li>Supplies list</li> </ul>  |