

ICD-10-CM Official Guidelines for Coding and Reporting 2010

Narrative changes appear in bold text

Items underlined have been moved within the guidelines since the 2009 version

Italics are used to indicate revisions to heading changes

The Centers for Medicare and Medicaid Services (CMS) and the National Center for Health Statistics (NCHS), two departments within the U.S. Federal Government's Department of Health and Human Services (DHHS) provide the following guidelines for coding and reporting using the International Classification of Diseases, 10th Revision, Clinical Modification (ICD-10-CM). These guidelines should be used as a companion document to the official version of the ICD-10-CM as published on the NCHS website. The ICD-10-CM is a morbidity classification published by the United States for classifying diagnoses and reason for visits in all health care settings. The ICD-10-CM is based on the ICD-10, the statistical classification of disease published by the World Health Organization (WHO).

These guidelines have been approved by the four organizations that make up the Cooperating Parties for the ICD-10-CM: the American Hospital Association (AHA), the American Health Information Management Association (AHIMA), CMS, and NCHS.

These guidelines are a set of rules that have been developed to accompany and complement the official conventions and instructions provided within the ICD-10-CM itself. **The instructions and conventions of the classification take precedence over guidelines.** These guidelines are based on the coding and sequencing instructions in Volumes I, and II of ICD-10-CM, but provide additional instruction. Adherence to these guidelines when assigning ICD-10-CM diagnosis codes is required under the Health Insurance Portability and Accountability Act (HIPAA). The diagnosis codes (Volumes 1-2) have been adopted under HIPAA for all healthcare settings. A joint effort between the healthcare provider and the coder is essential to achieve complete and accurate documentation, code assignment, and reporting of diagnoses and procedures. These guidelines have been developed to assist both the healthcare provider and the coder in identifying those diagnoses and procedures that are to be reported. The importance of consistent, complete documentation in the medical record cannot be overemphasized. Without such documentation accurate coding cannot be achieved. The entire record should be reviewed to determine the specific reason for the encounter and the conditions treated.

The term encounter is used for all settings, including hospital admissions. In the context of these guidelines, the term provider is used throughout the guidelines to mean physician or any qualified health care practitioner who is legally accountable for establishing the patient's diagnosis. Only this set of guidelines, approved by the Cooperating Parties, is official.

The guidelines are organized into sections. Section I includes the structure and conventions of the classification and general guidelines that apply to the entire classification, and chapter-specific guidelines that correspond to the chapters as they are arranged in the classification. Section II includes guidelines for selection of principal diagnosis for non-outpatient settings. Section III includes guidelines for reporting additional diagnoses in non-outpatient settings. Section IV is for outpatient coding and reporting. It is necessary to review all sections of the guidelines to fully understand all of the rules and instructions needed to code properly.

ICD-10-CM Official Guidelines for Coding and Reporting	1
Section I. Conventions, general coding guidelines and chapter specific guidelines.....	7
A. Conventions for the ICD-10-CM	7
1. The Alphabetic Index and Tabular List	7
2. Format and Structure:	7
3. Use of codes for reporting purposes	7
4. Placeholder character	7
5. 7 th Characters	8
6. Abbreviations.....	8
a. Index abbreviations.....	8
b. Tabular abbreviations.....	8
7. Punctuation	8
8. Use of “and”.....	8
9. Other and Unspecified codes	9
a. “Other” codes.....	9
b. “Unspecified” codes.....	9
10. Includes Notes.....	9
11. Inclusion terms.....	9
12. Excludes Notes.....	9
a. Excludes1	9
b. Excludes2.....	9
13. Etiology/manifestation convention (“code first”, “use additional code” and “in diseases classified elsewhere” notes).....	10
14. “And”	11
15. “With”	11
16. “See” and “See Also”.....	11
17. “Code also note”	11
18. Default codes	11
19. Syndromes	11
B. General Coding Guidelines.....	11
1. Locating a code in the ICD-10-CM	11
2. Level of Detail in Coding	12
3. Code or codes from A00.0 through T88.9, Z00-Z99.8.....	12
4. Signs and symptoms	12
5. Conditions that are an integral part of a disease process	12
6. Conditions that are not an integral part of a disease process.....	12
7. Multiple coding for a single condition.....	12
8. Acute and Chronic Conditions.....	13
9. Combination Code	13
10. Late Effects (Sequela).....	14
11. Impending or Threatened Condition.....	14
12. Reporting Same Diagnosis Code More than Once	14
13. Laterality	14
14. Documentation for BMI and Pressure Ulcer Stages.....	15
C. Chapter-Specific Coding Guidelines	15
1. Chapter 1: Certain Infectious and Parasitic Diseases (A00-B99).....	15

a.	Human Immunodeficiency Virus (HIV) Infections	15
b.	Infectious agents as the cause of diseases classified to other chapters	17
c.	Infections resistant to antibiotics	18
d.	Sepsis, Severe Sepsis, and Septic Shock	18
2.	Chapter 2: Neoplasms (C00-D49)	22
a.	Treatment directed at the malignancy	22
b.	Treatment of secondary site	22
c.	Coding and sequencing of complications	22
d.	Primary malignancy previously excised	23
e.	Admissions/Encounters involving chemotherapy, immunotherapy and radiation therapy ..	24
f.	Admission/encounter to determine extent of malignancy	24
g.	Symptoms, signs, and abnormal findings listed in Chapter 18 associated with neoplasms .	25
h.	Admission/encounter for pain control/management	25
i.	Malignancy in two or more noncontiguous sites	25
j.	Disseminated malignant neoplasm, unspecified	25
k.	Malignant neoplasm without specification of site	25
l.	Sequencing of neoplasm codes	25
m.	Current malignancy versus personal history of malignancy	27
n.	Leukemia in remission versus personal history of leukemia	27
o.	Aftercare following surgery for neoplasm	27
p.	Follow-up care for completed treatment of a malignancy	27
q.	Prophylactic organ removal for prevention of malignancy	27
r.	Malignant neoplasm associated with transplanted organ	28
3.	Chapter 3: Disease of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	28
4.	Chapter 4: Endocrine, Nutritional, and Metabolic Diseases (E00-E89)	28
a.	Diabetes mellitus	28
5.	Chapter 5: Mental and behavioral disorders (F01 – F99)	30
a.	Pain disorders related to psychological factors	30
6.	Chapter 6: Diseases of Nervous System and Sense Organs (G00-G99)	31
a.	Dominant/nondominant side	31
b.	Pain - Category G89	31
7.	Chapter 7: Diseases of Eye and Adnexa (H00-H59)	34
8.	Chapter 8: Diseases of Ear and Mastoid Process (H60-H95)	34
9.	Chapter 9: Diseases of Circulatory System (I00-I99)	35
a.	Hypertension	35
b.	Atherosclerotic coronary artery disease and angina	37
c.	Intraoperative and Postprocedural cerebrovascular accident	37
d.	Sequelae of Cerebrovascular Disease	37
e.	Acute myocardial infarction (AMI)	38
10.	Chapter 10: Diseases of Respiratory System (J00-J99)	39
a.	Chronic Obstructive Pulmonary Disease [COPD] and Asthma	39
b.	Acute Respiratory Failure	40
c.	Influenza due to certain identified influenza influenza viruses	40
11.	Chapter 11: Diseases of Digestive System (K00-K94)	41
12.	Chapter 12: Diseases of Skin and Subcutaneous Tissue (L00-L99)	41
a.	Pressure ulcer stage codes	41

13.	Chapter 13: Diseases of the Musculoskeletal System and Connective Tissue (M00-M99) ...	42
a.	Site and laterality	42
b.	Acute traumatic versus chronic or recurrent musculoskeletal conditions	43
c.	Coding of Pathologic Fractures	43
d.	Osteoporosis.....	43
14.	Chapter 14: Diseases of Genitourinary System (N00-N99)	44
a.	Chronic kidney disease	44
15.	Chapter 15: Pregnancy, Childbirth, and the Puerperium (O00-O9A).....	45
a.	General Rules for Obstetric Cases	45
b.	Selection of OB Principal or First-listed Diagnosis.....	46
c.	Pre-existing conditions versus conditions due to the pregnancy	47
d.	Pre-existing hypertension in pregnancy.....	48
e.	Fetal Conditions Affecting the Management of the Mother.....	48
f.	HIV Infection in Pregnancy, Childbirth and the Puerperium	48
g.	Diabetes mellitus in pregnancy.....	49
h.	Long term use of insulin	49
i.	Gestational (pregnancy induced) diabetes	49
j.	Sepsis and septic shock complicating abortion, pregnancy, childbirth and the puerperium	49
k.	Puerperal sepsis.....	50
l.	Alcohol and tobacco use during pregnancy, childbirth and the puerperium	50
m.	Poisoning, toxic effects, adverse effects and underdosing in a pregnant patient.....	50
n.	Normal Delivery, Code O80	51
o.	The Peripartum and Postpartum Periods.....	51
p.	Code O94, Sequelae of complication of pregnancy, childbirth, and the puerperium.....	52
q.	Abortions.....	52
16.	Chapter 16: Newborn (Perinatal) Guidelines (P00-P96)	53
a.	General Perinatal Rules.....	53
b.	Observation and Evaluation of Newborns for Suspected Conditions not Found	55
c.	Coding Additional Perinatal Diagnoses.....	55
d.	Prematurity and Fetal Growth Retardation.....	55
e.	Low birth weight and immaturity status	55
f.	Bacterial Sepsis of Newborn.....	56
g.	Stillbirth.....	56
17.	Chapter 17: Congenital malformations, deformations, and chromosomal abnormalities (Q00-Q99).....	56
18.	Chapter 18: Symptoms, signs, and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99).....	57
a.	Use of symptom codes	57
b.	Use of a symptom code with a definitive diagnosis code	57
c.	Combination codes that include symptoms	57
d.	Repeated falls.....	57
e.	Glasgow coma scale.....	58
f.	Functional quadriplegia	58
g.	SIRS due to Non-Infectious Process.....	58
g.	Death NOS	59

19.	Chapter 19: Injury, poisoning, and certain other consequences of external causes (S00-T88).....	59
a.	Code Extensions.....	59
b.	Coding of Injuries	60
c.	Coding of Traumatic Fractures	60
d.	Coding of Burns and Corrosions.....	61
e.	Adverse Effects, Poisoning , Underdosing and Toxic Effects.....	64
f.	Adult and child abuse, neglect and other maltreatment.....	67
g.	Complications of care	67
20.	Chapter 20: External Causes of Morbidity (V01- Y99).....	70
a.	General External Cause Coding Guidelines	70
b.	Place of Occurrence Guideline	71
c.	Activity Code	72
d.	Place of Occurrence, Activity, <i>and Status</i> Codes Used with other External Cause Code....	72
e.	If the Reporting Format Limits the Number of External Cause Codes	72
f.	Multiple External Cause Coding Guidelines	73
g.	Child and Adult Abuse Guideline.....	73
h.	Unknown or Undetermined Intent Guideline	73
i.	Late Effects of External Cause Guidelines	74
j.	Terrorism Guidelines	74
k.	External cause status	75
21.	Chapter 21: Factors influencing health status and contact with health services (Z00-Z99)...	75
a.	Use of Z codes in any healthcare setting	75
b.	Z Codes indicate a reason for an encounter.....	76
c.	Categories of Z Codes.....	76
Section II. Selection of Principal Diagnosis.....		90
A.	Codes for symptoms, signs, and ill-defined conditions	91
B.	Two or more interrelated conditions, each potentially meeting the definition for principal diagnosis.	91
C.	Two or more diagnoses that equally meet the definition for principal diagnosis.....	91
D.	Two or more comparative or contrasting conditions.	91
E.	A symptom(s) followed by contrasting/comparative diagnoses.....	91
F.	Original treatment plan not carried out.....	91
G.	Complications of surgery and other medical care.....	92
H.	Uncertain Diagnosis.....	92
I.	Admission from Observation Unit.....	92
1.	Admission Following Medical Observation.....	92
2.	Admission Following Post-Operative Observation	92
J.	Admission from Outpatient Surgery	92
Section III. Reporting Additional Diagnoses.....		93
A.	Previous conditions.....	93
B.	Abnormal findings	94
C.	Uncertain Diagnosis.....	94

Section IV. Diagnostic Coding and Reporting Guidelines for Outpatient Services	94
A. Selection of first-listed condition.....	95
1. Outpatient Surgery	95
2. Observation Stay	95
B. Codes from A00.0 through T88.9, Z00-Z99.....	95
C. Accurate reporting of ICD-10-CM diagnosis codes	95
D. Codes that describe symptoms and signs.....	96
E. Encounters for circumstances other than a disease or injury	96
F. Level of Detail in Coding	96
1. ICD-10-CM codes with 3, 4, or 5 digits	96
2. Use of full number of digits required for a code.....	96
G. ICD-10-CM code for the diagnosis, condition, problem, or other reason for encounter/visit.....	96
H. Uncertain diagnosis.....	96
I. Chronic diseases.....	97
J. Code all documented conditions that coexist.....	97
K. Patients receiving diagnostic services only.....	97
L. Patients receiving therapeutic services only	97
M. Patients receiving preoperative evaluations only.....	98
N. Ambulatory surgery	98
O. Routine outpatient prenatal visits.....	98
P. Encounters for general medical examinations with abnormal findings.....	98
Q. Encounters for routine health screenings	98

Section I. Conventions, general coding guidelines and chapter specific guidelines

The conventions, general guidelines and chapter-specific guidelines are applicable to all health care settings unless otherwise indicated. **The conventions and instructions of the classification take precedence over guidelines.**

A. Conventions for the ICD-10-CM

The conventions for the ICD-10-CM are the general rules for use of the classification independent of the guidelines. These conventions are incorporated within the Index and Tabular of the ICD-10-CM as instructional notes.

1. The Alphabetic Index and Tabular List

The ICD-10-CM is divided into the Index, an alphabetical list of terms and their corresponding code, and the Tabular List, a chronological list of codes divided into chapters based on body system or condition. The Index is divided into two parts, the Index to Diseases and Injury, and the Index to External Causes of Injury. Within the Index of Diseases and Injury there is a Neoplasm Table and a Table of Drugs and Chemicals.

See Section I.C2. General guidelines

See Section I.C.19. Adverse effects, poisoning, underdosing and toxic effects

2. Format and Structure:

The ICD-10-CM Tabular List contains categories, subcategories and codes. Characters for categories, subcategories and codes may be either a letter or a number. All categories are 3 characters. A three-character category that has no further subdivision is equivalent to a code. Subcategories are either 4 or 5 characters. Codes may be 4, 5, 6 or 7 characters. That is, each level of subdivision after a category is a subcategory. The final level of subdivision is a code. All codes in the Tabular List of the official version of the ICD-10-CM are in bold. Codes that have applicable 7th characters are still referred to as codes, not subcategories. A code that has an applicable 7th character is considered invalid without the 7th character.

The ICD-10-CM uses an indented format for ease in reference

3. Use of codes for reporting purposes

For reporting purposes only codes are permissible, not categories or subcategories, and any applicable 7th character is required.

4. Placeholder character

The ICD-10-CM utilizes a placeholder character “X”. The “X” is used as a 5th character placeholder at certain 6 character codes to allow for future

expansion. An example of this is at the poisoning, adverse effect and underdosing codes, categories T36-T50. Where a placeholder exists, the X must be used in order for the code to be considered a valid code.

5. 7th Characters

Certain ICD-10-CM categories have applicable 7th characters. The applicable 7th character is required for all codes within the category, or as the notes in the Tabular List instruct. The 7th character must always be the 7th character in the data field. If a code that requires a 7th character is not 6 characters, a placeholder X must be used to fill in the empty characters.

6. Abbreviations

a. Index abbreviations

NEC “Not elsewhere classifiable”

This abbreviation in the Index represents “other specified”.

When a specific code is not available for a condition, the Index directs the coder to the “other specified” code in the Tabular.

b. Tabular abbreviations

NEC “Not elsewhere classifiable”

This abbreviation in the Tabular represents “other specified”.

When a specific code is not available for a condition the Tabular includes an NEC entry under a code to identify the code as the “other specified” code.

NOS “Not otherwise specified”

This abbreviation is the equivalent of unspecified.

7. Punctuation

[] Brackets are used in the tabular list to enclose synonyms, alternative wording or explanatory phrases. Brackets are used in the Index to identify manifestation codes.

() Parentheses are used in both the Index and Tabular to enclose supplementary words that may be present or absent in the statement of a disease or procedure without affecting the code number to which it is assigned. The terms within the parentheses are referred to as nonessential modifiers.

:

Colons are used in the Tabular List after an incomplete term which needs one or more of the modifiers following the colon to make it assignable to a given category.

8. Use of “and”

When the term “and” is used in a narrative statement it represents and/or.

9. Other and Unspecified codes

a. “Other” codes

Codes titled “other” or “other specified” are for use when the information in the medical record provides detail for which a specific code does not exist. Index entries with NEC in the line designate “other” codes in the Tabular. These Index entries represent specific disease entities for which no specific code exists so the term is included within an “other” code.

b. “Unspecified” codes

Codes (usually a code with a 4th digit 9 or 5th digit 0 for diagnosis codes) titled “unspecified” are for use when the information in the medical record is insufficient to assign a more specific code. For those categories for which an unspecified code is not provided, the “other specified” code may represent both other and unspecified.

10. Includes Notes

This note appears immediately under a three-digit code title to further define, or give examples of, the content of the category.

11. Inclusion terms

List of terms is included under some codes. These terms are the conditions for which that code is to be used. The terms may be synonyms of the code title, or, in the case of “other specified” codes, the terms are a list of the various conditions assigned to that code. The inclusion terms are not necessarily exhaustive. Additional terms found only in the Index may also be assigned to a code.

12. Excludes Notes

The ICD-10-CM has two types of excludes notes. Each type of note has a different definition for use but they are all similar in that they indicate that codes excluded from each other are independent of each other.

a. Excludes1

A type 1 Excludes note is a pure excludes note. It means “NOT CODED HERE!” An Excludes1 note indicates that the code excluded should never be used at the same time as the code above the Excludes1 note. An Excludes1 is used when two conditions cannot occur together, such as a congenital form versus an acquired form of the same condition.

b. Excludes2

A type 2 excludes note represents “Not included here”. An excludes2 note indicates that the condition excluded is not part of the condition

represented by the code, but a patient may have both conditions at the same time. When an Excludes2 note appears under a code, it is acceptable to use both the code and the excluded code together, when appropriate.

13. Etiology/manifestation convention (“code first”, “use additional code” and “in diseases classified elsewhere” notes)

Certain conditions have both an underlying etiology and multiple body system manifestations due to the underlying etiology. For such conditions, the ICD-10-CM has a coding convention that requires the underlying condition be sequenced first followed by the manifestation. Wherever such a combination exists, there is a “use additional code” note at the etiology code, and a “code first” note at the manifestation code. These instructional notes indicate the proper sequencing order of the codes, etiology followed by manifestation.

In most cases the manifestation codes will have in the code title, “in diseases classified elsewhere.” Codes with this title are a component of the etiology/manifestation convention. The code title indicates that it is a manifestation code. “In diseases classified elsewhere” codes are never permitted to be used as first listed or principal diagnosis codes. They must be used in conjunction with an underlying condition code and they must be listed following the underlying condition. See category F02, Dementia in other diseases classified elsewhere, for an example of this convention.

There are manifestation codes that do not have “in diseases classified elsewhere” in the title. For such codes a “use additional code” note will still be present and the rules for sequencing apply.

In addition to the notes in the Tabular, these conditions also have a specific Index entry structure. In the Index both conditions are listed together with the etiology code first followed by the manifestation codes in brackets. The code in brackets is always to be sequenced second.

An example of the etiology/manifestation convention is dementia in Parkinson’s disease. In the index, code G20 is listed first, followed by code F02.80 or F02.81 in brackets. Code G20 represents the underlying etiology, Parkinson’s disease, and must be sequenced first, whereas codes F02.80 and F02.81 represent the manifestation of dementia in diseases classified elsewhere, with or without behavioral disturbance.

“Code first” and “Use additional code” notes are also used as sequencing rules in the classification for certain codes that are not part of an etiology/manifestation combination.

See Section I.B.7. Multiple coding for a single condition.

14. “And”

The word “and” should be interpreted to mean either “and” or “or” when it appears in a title.

15. “With”

The word “with” in the Alphabetic Index is sequenced immediately following the main term, not in alphabetical order.

16. “See” and “See Also”

The “see” instruction following a main term in the Index indicates that another term should be referenced. It is necessary to go to the main term referenced with the “see” note to locate the correct code.

A “see also” instruction following a main term in the index instructs that there is another main term that may also be referenced that may provide additional index entries that may be useful. It is not necessary to follow the “see also” note when the original main term provides the necessary code.

17. “Code also note”

A “code also” note instructs that two codes may be required to fully describe a condition, but this note does not provide sequencing direction.

18. Default codes

A code listed next to a main term in the ICD-10-CM Index is referred to as a default code. The default code represents that condition that is most commonly associated with the main term, or is the unspecified code for the condition. If a condition is documented in a medical record (for example, appendicitis) without any additional information, such as acute or chronic, the default code should be assigned.

19. Syndromes

Follow the Alphabetic Index guidance when coding syndromes. In the absence of index guidance, assign codes for the documented manifestations of the syndrome.

B. General Coding Guidelines

1. Locating a code in the ICD-10-CM

To select a code in the classification that corresponds to a diagnosis or reason for visit documented in a medical record, first locate the term in the Index, and then verify the code in the Tabular List. Read and be guided by instructional notations that appear in both the Index and the Tabular List.

It is essential to use both the Index and Tabular List when locating and assigning a code. The Index does not always provide the full code. Selection of the full code, including laterality and any applicable 7th character can only be done in the Tabular list. A dash (-) at the end of an Index entry indicates that additional characters are required. Even if a dash is not included at the Index entry, it is necessary to refer to the Tabular list to verify that no 7th character is required.

2. Level of Detail in Coding

Diagnosis codes are to be used and reported at their highest number of digits available.

ICD-10-CM diagnosis codes are composed of codes with 3, 4, 5, 6 or 7 digits. Codes with three digits are included in ICD-10-CM as the heading of a category of codes that may be further subdivided by the use of fourth and/or fifth digits, which provide greater detail.

A three-digit code is to be used only if it is not further subdivided. A code is invalid if it has not been coded to the full number of characters required for that code, including the 7th character, if applicable.

3. Code or codes from A00.0 through T88.9, Z00-Z99.8

The appropriate code or codes from A00.0 through T88.9, Z00-Z99.8 must be used to identify diagnoses, symptoms, conditions, problems, complaints or other reason(s) for the encounter/visit.

4. Signs and symptoms

Codes that describe symptoms and signs, as opposed to diagnoses, are acceptable for reporting purposes when a related definitive diagnosis has not been established (confirmed) by the provider. Chapter 18 of ICD-10-CM, Symptoms, Signs, and Abnormal Clinical and Laboratory Findings, Not Elsewhere Classified (codes R00.0 - R99) contains many, but not all codes for symptoms.

5. Conditions that are an integral part of a disease process

Signs and symptoms that are associated routinely with a disease process should not be assigned as additional codes, unless otherwise instructed by the classification.

6. Conditions that are not an integral part of a disease process

Additional signs and symptoms that may not be associated routinely with a disease process should be coded when present.

7. Multiple coding for a single condition

In addition to the etiology/manifestation convention that requires two codes to fully describe a single condition that affects multiple body systems, there are

other single conditions that also require more than one code. “Use additional code” notes are found in the Tabular at codes that are not part of an etiology/manifestation pair where a secondary code is useful to fully describe a condition. The sequencing rule is the same as the etiology/manifestation pair, “use additional code” indicates that a secondary code should be added.

For example, for bacterial infections that are not included in chapter 1, a secondary code from category B95, Streptococcus, Staphylococcus, and Enterococcus, as the cause of diseases classified elsewhere, or B96, Other bacterial agents as the cause of diseases classified elsewhere, may be required to identify the bacterial organism causing the infection. A “use additional code” note will normally be found at the infectious disease code, indicating a need for the organism code to be added as a secondary code.

“Code first” notes are also under certain codes that are not specifically manifestation codes but may be due to an underlying cause. When there is a “code first” note and an underlying condition is present, the underlying condition should be sequenced first.

“Code, if applicable, any causal condition first”, notes indicate that this code may be assigned as a principal diagnosis when the causal condition is unknown or not applicable. If a causal condition is known, then the code for that condition should be sequenced as the principal or first-listed diagnosis.

Multiple codes may be needed for late effects, complication codes and obstetric codes to more fully describe a condition. See the specific guidelines for these conditions for further instruction.

8. Acute and Chronic Conditions

If the same condition is described as both acute (subacute) and chronic, and separate subentries exist in the Alphabetic Index at the same indentation level, code both and sequence the acute (subacute) code first.

9. Combination Code

A combination code is a single code used to classify:

Two diagnoses, or

A diagnosis with an associated secondary process (manifestation)

A diagnosis with an associated complication

Combination codes are identified by referring to subterm entries in the Alphabetic Index and by reading the inclusion and exclusion notes in the Tabular List.

Assign only the combination code when that code fully identifies the diagnostic conditions involved or when the Alphabetic Index so directs. Multiple coding should not be used when the classification provides a

combination code that clearly identifies all of the elements documented in the diagnosis. When the combination code lacks necessary specificity in describing the manifestation or complication, an additional code should be used as a secondary code.

10. **Late Effects (Sequela)**

A late effect is the residual effect (condition produced) after the acute phase of an illness or injury has terminated. There is no time limit on when a late effect code can be used. The residual may be apparent early, such as in cerebral infarction, or it may occur months or years later, such as that due to a previous injury. Coding of late effects generally requires two codes sequenced in the following order: The condition or nature of the late effect is sequenced first. The late effect code is sequenced second.

An exception to the above guidelines are those instances where the code for late effect is followed by a manifestation code identified in the Tabular List and title, or the late effect code has been expanded (at the fourth, fifth or sixth character levels) to include the manifestation(s). The code for the acute phase of an illness or injury that led to the late effect is never used with a code for the late effect.

See Section I.C.9. Sequelae of cerebrovascular disease

See Section I.C.15. Sequelae of complication of pregnancy, childbirth and the puerperium

See Section I.C.19. Code extensions

11. **Impending or Threatened Condition**

Code any condition described at the time of discharge as “impending” or “threatened” as follows:

If it did occur, code as confirmed diagnosis.

If it did not occur, reference the Alphabetic Index to determine if the condition has a subentry term for “impending” or “threatened” and also reference main term entries for “Impending” and for “Threatened.”

If the subterms are listed, assign the given code.

If the subterms are not listed, code the existing underlying condition(s) and not the condition described as impending or threatened.

12. **Reporting Same Diagnosis Code More than Once**

Each unique ICD-10-CM diagnosis code may be reported only once for an encounter. This applies to bilateral conditions **when there are no distinct codes identifying laterality** or two different conditions classified to the same ICD-10-CM diagnosis code.

13. **Laterality**

For bilateral sites, the final character of the codes in the ICD-10-CM indicates laterality. An unspecified side code is also provided should the side not be

identified in the medical record. If no bilateral code is provided and the condition is bilateral, assign separate codes for both the left and right side.

14. Documentation for BMI and Pressure Ulcer Stages

For the Body Mass Index (BMI) and pressure ulcer stage codes, code assignment may be based on medical record documentation from clinicians who are not the patient’s provider (i.e., physician or other qualified healthcare practitioner legally accountable for establishing the patient’s diagnosis), since this information is typically documented by other clinicians involved in the care of the patient (e.g., a dietitian often documents the BMI and nurses often documents the pressure ulcer stages). However, the associated diagnosis (such as overweight, obesity, or pressure ulcer) must be documented by the patient’s provider. If there is conflicting medical record documentation, either from the same clinician or different clinicians, the patient’s attending provider should be queried for clarification.

The BMI codes should only be reported as secondary diagnoses. As with all other secondary diagnosis codes, the BMI codes should only be assigned when they meet the definition of a reportable additional diagnosis (see Section III, Reporting Additional Diagnoses).

C. Chapter-Specific Coding Guidelines

In addition to general coding guidelines, there are guidelines for specific diagnoses and/or conditions in the classification. Unless otherwise indicated, these guidelines apply to all health care settings. Please refer to Section II for guidelines on the selection of principal diagnosis.

1. Chapter 1: Certain Infectious and Parasitic Diseases (A00-B99)

a. Human Immunodeficiency Virus (HIV) Infections

1) Code only confirmed cases

Code only confirmed cases of HIV infection/illness. This is an exception to the hospital inpatient guideline Section II, H.

In this context, “confirmation” does not require documentation of positive serology or culture for HIV; the provider’s diagnostic statement that the patient is HIV positive, or has an HIV-related illness is sufficient.

2) Selection and sequencing of HIV codes

(a) Patient admitted for HIV-related condition

If a patient is admitted for an HIV-related condition, the principal diagnosis should be B20, followed by additional diagnosis codes for all reported HIV-related conditions.

(b) Patient with HIV disease admitted for unrelated condition

If a patient with HIV disease is admitted for an unrelated condition (such as a traumatic injury), the code for the unrelated condition (e.g., the nature of injury code) should be the principal diagnosis. Other diagnoses would be B20 followed by additional diagnosis codes for all reported HIV-related conditions.

(c) Whether the patient is newly diagnosed

Whether the patient is newly diagnosed or has had previous admissions/encounters for HIV conditions is irrelevant to the sequencing decision.

(d) Asymptomatic human immunodeficiency virus

Z21, Asymptomatic human immunodeficiency virus [HIV] infection status, is to be applied when the patient without any documentation of symptoms is listed as being “HIV positive,” “known HIV,” “HIV test positive,” or similar terminology. Do not use this code if the term “AIDS” is used or if the patient is treated for any HIV-related illness or is described as having any condition(s) resulting from his/her HIV positive status; use B20 in these cases.

(e) Patients with inconclusive HIV serology

Patients with inconclusive HIV serology, but no definitive diagnosis or manifestations of the illness, may be assigned code R75, Inconclusive laboratory evidence of human immunodeficiency virus [HIV].

(f) Previously diagnosed HIV-related illness

Patients with any known prior diagnosis of an HIV-related illness should be coded to B20. Once a patient has developed an HIV-related illness, the patient should always be assigned code B20 on every subsequent admission/encounter. Patients previously

diagnosed with any HIV illness (B20) should never be assigned to R75 or Z21, Asymptomatic human immunodeficiency virus [HIV] infection status.

(g) HIV Infection in Pregnancy, Childbirth and the Puerperium

During pregnancy, childbirth or the puerperium, a patient admitted (or presenting for a health care encounter) because of an HIV-related illness should receive a principal diagnosis code of O98.7-, Human immunodeficiency [HIV] disease complicating pregnancy, childbirth and the puerperium, followed by B20 and the code(s) for the HIV-related illness(es). Codes from Chapter 15 always take sequencing priority.

Patients with asymptomatic HIV infection status admitted (or presenting for a health care encounter) during pregnancy, childbirth, or the puerperium should receive codes of O98.7- and Z21.

(h) Encounters for testing for HIV

If a patient is being seen to determine his/her HIV status, use code Z11.4, Encounter for screening for human immunodeficiency virus [HIV]. Use additional codes for any associated high risk behavior.

If a patient with signs or symptoms is being seen for HIV testing, code the signs and symptoms. An additional counseling code Z71.7, Human immunodeficiency virus [HIV] counseling, may be used if counseling is provided during the encounter for the test.

When a patient returns to be informed of his/her HIV test results and the test result is negative, use code Z71.7, Human immunodeficiency virus [HIV] counseling.

If the results are positive, see previous guidelines and assign codes as appropriate.

b. Infectious agents as the cause of diseases classified to other chapters

Certain infections are classified in chapters other than Chapter 1 and no organism is identified as part of the infection code. In these

instances, it is necessary to use an additional code from Chapter 1 to identify the organism. A code from category B95, Streptococcus, Staphylococcus, and Enterococcus as the cause of diseases classified to other chapters, B96, Other bacterial agents as the cause of diseases classified to other chapters, or B97, Viral agents as the cause of diseases classified to other chapters, is to be used as an additional code to identify the organism. An instructional note will be found at the infection code advising that an additional organism code is required.

c. Infections resistant to antibiotics

Many bacterial infections are resistant to current antibiotics. It is necessary to identify all infections documented as antibiotic resistant. Assign code Z16, Infection with drug resistant microorganisms, following the infection code for these cases.

d. Sepsis, Severe Sepsis, and Septic Shock

1) Coding of Sepsis and Severe Sepsis

(a) Sepsis

For a diagnosis of sepsis, assign the appropriate code for the underlying systemic infection. If the type of infection or causal organism is not further specified, assign code A41.9, Sepsis, unspecified.

A code from subcategory R65.2, Severe sepsis, should not be assigned unless severe sepsis or an associated acute organ dysfunction is documented.

(i) Negative or inconclusive blood cultures and sepsis

Negative or inconclusive blood cultures do not preclude a diagnosis of sepsis in patients with clinical evidence of the condition, however, the provider should be queried.

(ii) Urosepsis

The term urosepsis is a nonspecific term. It is not to be considered synonymous with sepsis. It has no default code in the Alphabetic Index. Should a provider use this term, he/she must be queried for clarification.

(iii) Sepsis with organ dysfunction

If a patient has sepsis and associated acute organ dysfunction or multiple organ dysfunction

(MOD), follow the instructions for coding severe sepsis.

- (iv) Acute organ dysfunction that is not clearly associated with the sepsis

If a patient has sepsis and an acute organ dysfunction, but the medical record documentation indicates that the acute organ dysfunction is related to a medical condition other than the sepsis, do not assign a code from subcategory R65.2, Severe sepsis. An acute organ dysfunction must be associated with the sepsis in order to assign the severe sepsis code. If the documentation is not clear as to whether an acute organ dysfunction is related to the sepsis or another medical condition, query the provider.

(b) Severe sepsis

The coding of severe sepsis requires a minimum of 2 codes: first a code for the underlying systemic infection, followed by a code from subcategory R65.2, Severe sepsis. If the causal organism is not documented, assign code A41.9, Sepsis, unspecified, for the infection. Additional code(s) for the associated acute organ dysfunction are also required.

Due to the complex nature of severe sepsis, some cases may require querying the provider prior to assignment of the codes.

2) Septic shock

Septic shock is circulatory failure associated with severe sepsis, and therefore, it represents a type of acute organ dysfunction. For all cases of septic shock, the code for the underlying systemic infection should be sequenced first, followed by code R65.21, Severe sepsis with septic shock. Any additional codes for the other acute organ dysfunctions should also be assigned.

Septic shock indicates the presence of severe sepsis. Code R65.21, Severe sepsis with septic shock, must be assigned if septic shock is documented in the medical record, even if the term severe sepsis is not documented.

3) Sequencing of severe sepsis

If severe sepsis is present on admission, and meets the definition of principal diagnosis, the underlying systemic infection should be assigned as principal diagnosis followed by the appropriate code from subcategory R65.2 as required by the sequencing rules in the Tabular List. A code from subcategory R65.2 can never be assigned as a principal diagnosis.

When severe sepsis develops during an encounter (it was not present on admission) the underlying systemic infection and the appropriate code from subcategory R65.2 should be assigned as secondary diagnoses.

Severe sepsis may be present on admission but the diagnosis may not be confirmed until sometime after admission. If the documentation is not clear whether severe sepsis was present on admission, the provider should be queried.

4) Sepsis and severe sepsis with a localized infection

If the reason for admission is both sepsis or severe sepsis and a localized infection, such as pneumonia or cellulitis, a code(s) for the underlying systemic infection should be assigned first and the code for the localized infection should be assigned as a secondary diagnosis. If the patient has severe sepsis, a code from subcategory R65.2 should also be assigned as a secondary diagnosis. If the patient is admitted with a localized infection, such as pneumonia, and sepsis/severe sepsis doesn't develop until after admission, the localized infection should be assigned first, followed by the appropriate sepsis/severe sepsis codes.

5) Sepsis due to a postprocedural infection

Sepsis resulting from a postprocedural infection is a complication of medical care. For such cases, the postprocedural infection code, such as, T80.2, Infections following infusion, transfusion, and therapeutic injection, T81.4, Infection following a procedure, T88.0, Infection following immunization, or O86.0, Infection of obstetric surgical wound, should be coded first, followed by the code for the specific infection. If the patient has severe sepsis the appropriate code from subcategory R65.2 should also be assigned with the additional code(s) for any acute organ dysfunction.

6) Sepsis and severe sepsis associated with a noninfectious process (condition)

In some cases a noninfectious process (condition), such as trauma, may lead to an infection which can result in sepsis or severe sepsis. If sepsis or severe sepsis is documented as associated with a noninfectious condition, such as a burn or serious injury, and this condition meets the definition for principal diagnosis, the code for the noninfectious condition should be sequenced first, followed by the code for the resulting infection. If severe sepsis is present a code from subcategory R65.2 should also be assigned with any associated organ dysfunction(s) codes. It is not necessary to assign a code from subcategory R65.1, Systemic inflammatory response syndrome (SIRS) of non-infectious origin, for these cases.

If the infection meets the definition of principal diagnosis it should be sequenced before the non-infectious condition. When both the associated non-infectious condition and the infection meet the definition of principal diagnosis either may be assigned as principal diagnosis.

Only one code from category R65, Symptoms and signs specifically associated with systemic inflammation and infection, should be assigned. Therefore, when a non-infectious condition leads to an infection resulting in severe sepsis, assign the appropriate code from subcategory R65.2, Severe sepsis. Do not additionally assign a code from subcategory R65.1, Systemic inflammatory response syndrome (SIRS) of non-infectious origin.

See Section I.C.18. SIRS due to non-infectious process

7) Sepsis and septic shock complicating abortion, pregnancy, childbirth, and the puerperium

See Section I.C.15. Sepsis and septic shock complicating abortion, pregnancy, childbirth and the puerperium

8) Newborn sepsis

See Section I.C.16. Newborn sepsis

2. Chapter 2: Neoplasms (C00-D49)

General guidelines

Chapter 2 of the ICD-10-CM contains the codes for most benign and all malignant neoplasms. Certain benign neoplasms, such as prostatic adenomas, may be found in the specific body system chapters. To properly code a neoplasm it is necessary to determine from the record if the neoplasm is benign, in-situ, malignant, or of uncertain histologic behavior. If malignant, any secondary (metastatic) sites should also be determined.

The neoplasm table in the Alphabetic Index should be referenced first. However, if the histological term is documented, that term should be referenced first, rather than going immediately to the Neoplasm Table, in order to determine which column in the Neoplasm Table is appropriate. For example, if the documentation indicates “adenoma,” refer to the term in the Alphabetic Index to review the entries under this term and the instructional note to “see also neoplasm, by site, benign.” The table provides the proper code based on the type of neoplasm and the site. It is important to select the proper column in the table that corresponds to the type of neoplasm. The Tabular should then be referenced to verify that the correct code has been selected from the table and that a more specific site code does not exist.

See Section I.C.21. Factors influencing health status and contact with health services, Status, for information regarding Z15.0, codes for genetic susceptibility to cancer.

a. Treatment directed at the malignancy

If the treatment is directed at the malignancy, designate the malignancy as the principal diagnosis.

The only exception to this guideline is if a patient admission/encounter is solely for the administration of chemotherapy, immunotherapy or radiation therapy, assign the appropriate Z51.-- code as the first-listed or principal diagnosis, and the diagnosis or problem for which the service is being performed as a secondary diagnosis.

b. Treatment of secondary site

When a patient is admitted because of a primary neoplasm with metastasis and treatment is directed toward the secondary site only, the secondary neoplasm is designated as the principal diagnosis even though the primary malignancy is still present.

c. Coding and sequencing of complications

Coding and sequencing of complications associated with the malignancies or with the therapy thereof are subject to the following guidelines:

1) Anemia associated with malignancy

When admission/encounter is for management of an anemia associated with the malignancy, and the treatment is only for anemia, the appropriate code for the malignancy is sequenced as the principal or first-listed diagnosis followed by code D63.0, Anemia in neoplastic disease).

2) Anemia associated with chemotherapy, immunotherapy and radiation therapy

When the admission/encounter is for management of an anemia associated with **an adverse effect of** chemotherapy, immunotherapy or radiotherapy and the only treatment is for the anemia, **the appropriate adverse effect code should be sequenced first, followed by the appropriate codes for the anemia and neoplasm.**

3) Management of dehydration due to the malignancy

When the admission/encounter is for management of dehydration due to the malignancy or the therapy, or a combination of both, and only the dehydration is being treated (intravenous rehydration), the dehydration is sequenced first, followed by the code(s) for the malignancy.

4) Treatment of a complication resulting from a surgical procedure

When the admission/encounter is for treatment of a complication resulting from a surgical procedure, designate the complication as the principal or first-listed diagnosis if treatment is directed at resolving the complication.

d. Primary malignancy previously excised

When a primary malignancy has been previously excised or eradicated from its site and there is no further treatment directed to that site and there is no evidence of any existing primary malignancy, a code from category Z85, Personal history of primary and secondary malignant neoplasm, should be used to indicate the former site of the malignancy. Any mention of extension, invasion, or metastasis to another site is coded as a secondary malignant neoplasm to that site. The secondary site may be the principal or first-listed with the Z85 code used as a secondary code.

e. Admissions/Encounters involving chemotherapy, immunotherapy and radiation therapy

1) Episode of care involves surgical removal of neoplasm

When an episode of care involves the surgical removal of a neoplasm, primary or secondary site, followed by adjunct chemotherapy or radiation treatment during the same episode of care, the neoplasm code should be assigned as principal or first-listed diagnosis, using codes in the C00-D49 series or where appropriate in the C83-C90 series.

2) Patient admission/encounter solely for administration of chemotherapy, immunotherapy and radiation therapy

If a patient admission/encounter is solely for the administration of chemotherapy, immunotherapy or radiation therapy assign code Z51.0, Encounter for antineoplastic radiation therapy, or Z51.11, Encounter for antineoplastic chemotherapy, or Z51.12, Encounter for antineoplastic immunotherapy as the first-listed or principal diagnosis. If a patient receives more than one of these therapies during the same admission more than one of these codes may be assigned, in any sequence.

The malignancy for which the therapy is being administered should be assigned as a secondary diagnosis.

3) Patient admitted for radiation therapy, chemotherapy or immunotherapy and develops complications

When a patient is admitted for the purpose of radiotherapy, immunotherapy or chemotherapy and develops complications such as uncontrolled nausea and vomiting or dehydration, the principal or first-listed diagnosis is Z51.0, Encounter for antineoplastic radiation therapy, or Z51.11, Encounter for antineoplastic chemotherapy, or Z51.12, Encounter for antineoplastic immunotherapy followed by any codes for the complications.

f. Admission/encounter to determine extent of malignancy

When the reason for admission/encounter is to determine the extent of the malignancy, or for a procedure such as paracentesis or thoracentesis, the primary malignancy or appropriate metastatic site is designated as the principal or first-listed diagnosis, even though chemotherapy or radiotherapy is administered.

g. Symptoms, signs, and abnormal findings listed in Chapter 18 associated with neoplasms

Symptoms, signs, and ill-defined conditions listed in Chapter 18 characteristic of, or associated with, an existing primary or secondary site malignancy cannot be used to replace the malignancy as principal or first-listed diagnosis, regardless of the number of admissions or encounters for treatment and care of the neoplasm.

See section I.C.21. Factors influencing health status and contact with health services, Encounter for prophylactic organ removal.

h. Admission/encounter for pain control/management

See Section I.C.6. for information on coding admission/encounter for pain control/management.

i. Malignancy in two or more noncontiguous sites

A patient may have more than one malignant tumor in the same organ. These tumors may represent different primaries or metastatic disease, depending on the site. Should the documentation be unclear, the provider should be queried as to the status of each tumor so that the correct codes can be assigned.

j. Disseminated malignant neoplasm, unspecified

Code C80.0, Disseminated malignant neoplasm, unspecified, is for use only in those cases where the patient has advanced metastatic disease and no known primary or secondary sites are specified. It should not be used in place of assigning codes for the primary site and all known secondary sites.

k. Malignant neoplasm without specification of site

Code C80.1, Malignant neoplasm, unspecified, equates to Cancer, unspecified. This code should only be used when no determination can be made as to the primary site of a malignancy. This code should rarely be used in the inpatient setting.

l. Sequencing of neoplasm codes

1) Encounter for treatment of primary malignancy

If the reason for the encounter is for treatment of a primary malignancy, assign the malignancy as the principal/first listed diagnosis. The primary site is to be sequenced first, followed by any metastatic sites.

2) Encounter for treatment of secondary malignancy

When an encounter is for a primary malignancy with metastasis and treatment is directed toward the metastatic

(secondary) site(s) only, the metastatic site(s) is designated as the principal/first listed diagnosis. The primary malignancy is coded as an additional code.

3) Malignant neoplasm in a pregnant patient

Codes from chapter 15, Pregnancy, childbirth, and the puerperium, are always sequenced first on a medical record. A code from subcategory O94.1-, Malignant neoplasm complicating pregnancy, childbirth, and the puerperium, should be used first, followed by the appropriate code from Chapter 2 to indicate the type of neoplasm.

4) Encounter for complication associated with a neoplasm

When an encounter is for management of a complication associated with a neoplasm, such as dehydration, and the treatment is only for the complication, the complication is coded first, followed by the appropriate code(s) for the neoplasm.

The exception to this guideline is anemia. When the admission/encounter is for management of an anemia associated with the malignancy, and the treatment is only for anemia, the appropriate code for the malignancy is sequenced as the principal or first-listed diagnosis followed by code D63.0, Anemia in neoplastic disease.

5) Complication from surgical procedure for treatment of a neoplasm

When an encounter is for treatment of a complication resulting from a surgical procedure performed for the treatment of the neoplasm, designate the complication as the principal/first listed diagnosis. See guideline regarding the coding of a current malignancy versus personal history to determine if the code for the neoplasm should also be assigned.

6) Pathologic fracture due to a neoplasm

When an encounter is for a pathological fracture due to a neoplasm, if the focus of treatment is the fracture, a code from subcategory M84.5, Pathological fracture in neoplastic disease, should be sequenced first, followed by the code for the neoplasm.

If the focus of treatment is the neoplasm with an associated pathological fracture, the neoplasm code should be sequenced

first, followed by a code from M84.5 for the pathological fracture. The “code also” note at M84.5 provides this sequencing instruction.

m. Current malignancy versus personal history of malignancy

When a primary malignancy has been excised but further treatment, such as an additional surgery for the malignancy, radiation therapy or chemotherapy is directed to that site, the primary malignancy code should be used until treatment is completed.

When a primary malignancy has been previously excised or eradicated from its site, there is no further treatment (of the malignancy) directed to that site, and there is no evidence of any existing primary malignancy, a code from category Z85, Personal history of primary and secondary malignant neoplasm, should be used to indicate the former site of the malignancy.

See Section I.C.21. Factors influencing health status and contact with health services, History (of)

n. Leukemia in remission versus personal history of leukemia

The categories for leukemia, and category C90, Multiple myeloma, have codes for in remission. There are also codes Z85.6, Personal history of leukemia, and Z85.79, Personal history of other malignant neoplasms of lymphoid, hematopoietic and related tissues. If the documentation is unclear, as to whether the patient is in remission, the provider should be queried.

See Section I.C.21. Factors influencing health status and contact with health services, History (of)

o. Aftercare following surgery for neoplasm

See Section I.C.21. Factors influencing health status and contact with health services, Aftercare

p. Follow-up care for completed treatment of a malignancy

See Section I.C.21. Factors influencing health status and contact with health services, Follow-up

q. Prophylactic organ removal for prevention of malignancy

See Section I.C. 21, Factors influencing health status and contact with health services, Prophylactic organ removal

- r. **Malignant neoplasm associated with transplanted organ**
A malignant neoplasm of a transplanted organ should be coded as a transplant complication. Assign first the appropriate code from category T86.-, Complications of transplanted organ, followed by code C80.2, Malignant neoplasm associated with transplanted organ. Use an additional code for the specific malignancy.

3. **Chapter 3: Disease of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)**

Reserved for future guideline expansion

4. **Chapter 4: Endocrine, Nutritional, and Metabolic Diseases (E00-E89)**

a. **Diabetes mellitus**

The diabetes mellitus codes are combination codes that include the type of DM, the body system affected, and the complications affecting that body system. As many codes within a particular category as are necessary to describe all of the complications of the disease may be used. They should be sequenced based on the reason for a particular encounter. Assign as many codes from categories E08 – E13 as needed to identify all of the associated conditions that the patient has.

1) **Type of diabetes**

The age of a patient is not the sole determining factor, though most type 1 diabetics develop the condition before reaching puberty. For this reason type 1 diabetes mellitus is also referred to as juvenile diabetes.

2) **Type of diabetes mellitus not documented**

If the type of diabetes mellitus is not documented in the medical record the default is E11.-, Type 2 diabetes mellitus.

3) **Diabetes mellitus and the use of insulin**

If the documentation in a medical record does not indicate the type of diabetes but does indicate that the patient uses insulin, code E11, Type 2 diabetes mellitus, should be assigned for type 2 patients who routinely use insulin, code Z79.4, Long-term (current) use of insulin, should also be assigned to indicate that the patient uses insulin. Code Z79.4 should not be assigned if insulin is given temporarily to bring a type 2 patient's blood sugar under control during an encounter.

4) Diabetes mellitus in pregnancy and gestational diabetes

See Section I.C.15. Diabetes mellitus in pregnancy.

See Section I.C.15. Gestational (pregnancy induced) diabetes

5) Complications due to insulin pump malfunction

(a) Underdose of insulin due insulin pump failure

An underdose of insulin due to an insulin pump failure should be assigned to a code from subcategory T85.6, Mechanical complication of other specified internal and external prosthetic devices, implants and grafts, that specifies the type of pump malfunction, as the principal or first listed code, followed by code T38.3x6-, Underdosing of insulin and oral hypoglycemic [antidiabetic] drugs. Additional codes for the type of diabetes mellitus and any associated complications due to the underdosing should also be assigned.

(b) Overdose of insulin due to insulin pump failure

The principal or first listed code for an encounter due to an insulin pump malfunction resulting in an overdose of insulin, should also be T85.6-, Mechanical complication of other specified internal and external prosthetic devices, implants and grafts, followed by code T38.3x1-, Poisoning by insulin and oral hypoglycemic [antidiabetic] drugs, accidental (unintentional).

6) Secondary Diabetes Mellitus

Codes under category E08, Diabetes mellitus due to underlying condition, and E09, Drug or chemical induced diabetes mellitus, identify complications/manifestations associated with secondary diabetes mellitus. Secondary diabetes is always caused by another condition or event (e.g., cystic fibrosis, malignant neoplasm of pancreas, pancreatectomy, adverse effect of drug, or poisoning).

(a) Secondary diabetes mellitus and the use of insulin

For patients who routinely use insulin, code Z79.4, Long-term (current) use of insulin, should also be assigned. Code Z79.4 should not be assigned if insulin is given temporarily to bring a patient's blood sugar under control during an encounter.

(b) **Assigning and sequencing secondary diabetes codes and its causes**

The sequencing of the secondary diabetes codes in relationship to codes for the cause of the diabetes is based on the tabular instructions for categories E08 and E09. For example, for category E08, Diabetes mellitus due to underlying condition, code first the underlying condition; for category E09, Drug or chemical induced diabetes mellitus, code first the drug or chemical (T36-T65).

(i) **Secondary diabetes mellitus due to pancreatectomy**

For postpancreatectomy diabetes mellitus (lack of insulin due to the surgical removal of all or part of the pancreas), assign code E89.1, Postsurgical hypoinsulinemia. Assign a code from category E08 and code Z79.4, Other acquired absence of organ, as additional codes.

(ii) **Secondary diabetes due to drugs**

Secondary diabetes may be caused by an adverse effect of correctly administered medications, poisoning or late effect of poisoning.

See section I.C.19.e for coding of adverse effects and poisoning, and section I.C.20 for external cause code reporting.

5. Chapter 5: Mental and behavioral disorders (F01 – F99)

a. Pain disorders related to psychological factors

Assign code F45.41, for pain that is exclusively psychological. Code F45.41, Pain disorder with related psychological factors, should be used following the appropriate code from category G89, Pain, not elsewhere classified, if there is documentation of a psychological component for a patient with acute or chronic pain.

See Section I.C.6. Pain

6. **Chapter 6: Diseases of Nervous System and Sense Organs (G00-G99)**

a. **Dominant/nondominant side**

Codes from category G81, Hemiplegia and hemiparesis, and subcategories, G83.1, Monoplegia of lower limb, G83.2, Monoplegia of upper limb, and G83.3, Monoplegia, unspecified, identify whether the dominant or nondominant side is affected. Should this information not be available in the record, **and the classification system does not indicate a default**, the default should be dominant. For ambidextrous patients, the default should also be dominant.

b. **Pain - Category G89**

1) **General coding information**

Codes in category G89, Pain, not elsewhere classified, may be used in conjunction with codes from other categories and chapters to provide more detail about acute or chronic pain and neoplasm-related pain, unless otherwise indicated below.

If the pain is not specified as acute or chronic, post-thoracotomy, postprocedural, or neoplasm-related, do not assign codes from category G89.

A code from category G89 should not be assigned if the underlying (definitive) diagnosis is known, unless the reason for the encounter is pain control/ management and not management of the underlying condition.

When an admission or encounter is for a procedure aimed at treating the underlying condition (e.g., spinal fusion, kyphoplasty), a code for the underlying condition (e.g., vertebral fracture, spinal stenosis) should be assigned as the principal diagnosis. No code from category G89 should be assigned.

(a) **Category G89 Codes as Principal or First-Listed Diagnosis**

Category G89 codes are acceptable as principal diagnosis or the first-listed code:

- When pain control or pain management is the reason for the admission/encounter (e.g., a patient with displaced intervertebral disc, nerve impingement and severe back pain presents for

injection of steroid into the spinal canal). The underlying cause of the pain should be reported as an additional diagnosis, if known.

- When a patient is admitted for the insertion of a neurostimulator for pain control, assign the appropriate pain code as the principal or first listed diagnosis. When an admission or encounter is for a procedure aimed at treating the underlying condition and a neurostimulator is inserted for pain control during the same admission/encounter, a code for the underlying condition should be assigned as the principal diagnosis and the appropriate pain code should be assigned as a secondary diagnosis.

(b) Use of Category G89 Codes in Conjunction with Site Specific Pain Codes

(i) Assigning Category G89 and Site-Specific Pain Codes

Codes from category G89 may be used in conjunction with codes that identify the site of pain (including codes from chapter 18) if the category G89 code provides additional information. For example, if the code describes the site of the pain, but does not fully describe whether the pain is acute or chronic, then both codes should be assigned.

(ii) Sequencing of Category G89 Codes with Site-Specific Pain Codes

The sequencing of category G89 codes with site-specific pain codes (including chapter 18 codes), is dependent on the circumstances of the encounter/admission as follows:

- If the encounter is for pain control or pain management, assign the code from category G89 followed by the code identifying the specific site of pain (e.g., encounter for pain management for acute neck pain from trauma is assigned code G89.11, Acute pain due to trauma, followed by code M54.2, Cervicalgia, to identify the site of pain).

- If the encounter is for any other reason except pain control or pain management, and a related definitive diagnosis has not been established (confirmed) by the provider, assign the code for the specific site of pain first, followed by the appropriate code from category G89.

2) Pain due to devices, implants and grafts

See Section I.C.19. Pain due to medical devices

3) Postoperative Pain

The provider's documentation should be used to guide the coding of postoperative pain, as well as *Section III. Reporting Additional Diagnoses* and *Section IV. Diagnostic Coding and Reporting in the Outpatient Setting*.

The default for post-thoracotomy and other postoperative pain not specified as acute or chronic is the code for the acute form.

Routine or expected postoperative pain immediately after surgery should not be coded.

(a) Postoperative pain not associated with specific postoperative complication

Postoperative pain not associated with a specific postoperative complication is assigned to the appropriate postoperative pain code in category G89.

(b) Postoperative pain associated with specific postoperative complication

Postoperative pain associated with a specific postoperative complication (such as painful wire sutures) is assigned to the appropriate code(s) found in Chapter 19, Injury, poisoning, and certain other consequences of external causes. If appropriate, use additional code(s) from category G89 to identify acute or chronic pain (G89.18 or G89.28).

4) Chronic pain

Chronic pain is classified to subcategory G89.2. There is no time frame defining when pain becomes chronic

pain. The provider's documentation should be used to guide use of these codes.

5) Neoplasm Related Pain

Code G89.3 is assigned to pain documented as being related, associated or due to cancer, primary or secondary malignancy, or tumor. This code is assigned regardless of whether the pain is acute or chronic.

This code may be assigned as the principal or first-listed code when the stated reason for the admission/encounter is documented as pain control/pain management. The underlying neoplasm should be reported as an additional diagnosis.

When the reason for the admission/encounter is management of the neoplasm and the pain associated with the neoplasm is also documented, code G89.3 may be assigned as an additional diagnosis. It is not necessary to assign an additional code for the site of the pain.

See Section I.C.2 for instructions on the sequencing of neoplasms for all other stated reasons for the admission/encounter (except for pain control/pain management).

6) Chronic pain syndrome

Central pain syndrome (G89.0) and chronic pain syndrome (G89.4) are different than the term "chronic pain," and therefore codes should only be used when the provider has specifically documented this condition.

See Section I.C.5. Pain disorders related to psychological factors

7. Chapter 7: Diseases of Eye and Adnexa (H00-H59)

Reserved for future guideline expansion

8. Chapter 8: Diseases of Ear and Mastoid Process (H60-H95)

Reserved for future guideline expansion

9. Chapter 9: Diseases of Circulatory System (I00-I99)

a. Hypertension

1) Hypertension with Heart Disease

Heart conditions classified to I50.- or I51.4-I51.9, are assigned to, a code from category I11, Hypertensive heart disease, when a causal relationship is stated (due to hypertension) or implied (hypertensive). Use an additional code from category I50, Heart failure, to identify the type of heart failure in those patients with heart failure.

The same heart conditions (I50.-, I51.4-I51.9) with hypertension, but without a stated causal relationship, are coded separately. Sequence according to the circumstances of the admission/encounter.

2) Hypertensive Chronic Kidney Disease

Assign codes from category I12, Hypertensive chronic kidney disease, when both hypertension and a condition classifiable to category N18, Chronic kidney disease (CKD), are present. Unlike hypertension with heart disease, ICD-10-CM presumes a cause-and-effect relationship and classifies chronic kidney disease with hypertension as hypertensive chronic kidney disease.

The appropriate code from category N18 should be used as a secondary code with a code from category I12 to identify the stage of chronic kidney disease.

See Section I.C.14. Chronic kidney disease.

If a patient has hypertensive chronic kidney disease **and acute** renal failure, an additional code for the acute renal failure is required.

3) Hypertensive Heart and Chronic Kidney Disease

Assign codes from combination category I13, Hypertensive heart and chronic kidney disease, when both hypertensive kidney disease and hypertensive heart disease are stated in the diagnosis. Assume a relationship between the hypertension and the chronic kidney disease, whether or not the condition is so designated. If heart failure is present, assign an additional code from category I50 to identify the type of heart failure.

The appropriate code from category N18, Chronic kidney disease, should be used as a secondary code with a code from category I13 to identify the stage of chronic kidney disease.

See Section I.C.14. Chronic kidney disease.

The codes in category I13, Hypertensive heart and chronic kidney disease, are combination codes that include hypertension, heart disease and chronic kidney disease. The Includes note at I13 specifies that the conditions included at I11 and I12 are included together in I13. If a patient has hypertension, heart disease and chronic kidney disease then a code from I13 should be used, not individual codes for hypertension, heart disease and chronic kidney disease, or codes from I11 or I12.

For patients with both acute renal failure and chronic kidney disease an additional code for acute renal failure is required.

4) Hypertensive Cerebrovascular Disease

For hypertensive cerebrovascular disease, first assign the appropriate code from categories I60-I69, followed by the appropriate hypertension code.

5) Hypertensive Retinopathy

Code H35.0, Hypertensive retinopathy, should be used with code I10, Essential (primary) hypertension, to include the systemic hypertension. The sequencing is based on the reason for the encounter.

6) Hypertension, Secondary

Secondary hypertension is due to an underlying condition. Two codes are required: one to identify the underlying etiology and one from category I15 to identify the hypertension. Sequencing of codes is determined by the reason for admission/encounter.

7) Hypertension, Transient

Assign code R03.0, Elevated blood pressure reading without diagnosis of hypertension, unless patient has an established diagnosis of hypertension. Assign code O13.-, Gestational [pregnancy-induced] hypertension without significant proteinuria, or O14.-, Gestational [pregnancy-induced] hypertension with significant proteinuria, for transient hypertension of pregnancy.

8) Hypertension, Controlled

This diagnostic statement usually refers to an existing state of hypertension under control by therapy. Assign code I10.

9) Hypertension, Uncontrolled

Uncontrolled hypertension may refer to untreated hypertension or hypertension not responding to current therapeutic regimen. In either case, assign code I10.

b. Atherosclerotic coronary artery disease and angina

ICD-10-CM has combination codes for atherosclerotic heart disease with angina pectoris. The subcategories for these codes are I25.11, Atherosclerotic heart disease of native coronary artery with angina pectoris and I25.7, Atherosclerosis of coronary artery bypass graft(s) and coronary artery of transplanted heart with angina pectoris.

When using one of these combination codes it is not necessary to use an additional code for angina pectoris. A causal relationship can be assumed in a patient with both atherosclerosis and angina pectoris, unless the documentation indicates the angina is due to something other than the atherosclerosis.

If a patient with coronary artery disease is admitted due to an acute myocardial infarction (AMI), the AMI should be sequenced before the coronary artery disease.

See Section I.C.9. Acute myocardial infarction (AMI)

c. Intraoperative and Postprocedural cerebrovascular accident

Medical record documentation should clearly specify the cause-and-effect relationship between the medical intervention and the cerebrovascular accident in order to assign a code for intraoperative or postprocedural cerebrovascular accident.

Proper code assignment depends on whether it was an infarction or hemorrhage and whether it occurred intraoperatively or postoperatively. If it was a cerebral hemorrhage, code assignment depends on the type of procedure performed.

d. Sequelae of Cerebrovascular Disease

1) Category I69, Sequelae of Cerebrovascular disease

Category I69 is used to indicate conditions classifiable to categories I60-I67 as the causes of late effects (neurologic

deficits), themselves classified elsewhere. These “late effects” include neurologic deficits that persist after initial onset of conditions classifiable to categories I60-I67. The neurologic deficits caused by cerebrovascular disease may be present from the onset or may arise at any time after the onset of the condition classifiable to categories I60-I67.

2) Codes from category I69 with codes from I60-I67

Codes from category I69 may be assigned on a health care record with codes from I60-I67, if the patient has a current cerebrovascular accident (CVA) and deficits from an old CVA.

3) Code Z86.73

Assign code Z86.73, Personal history of transient ischemic attack (TIA), and cerebral infarction without residual deficits (and not a code from category I69) as an additional code for history of cerebrovascular disease when no neurologic deficits are present.

e. Acute myocardial infarction (AMI)

1) ST elevation myocardial infarction (STEMI) and non ST elevation myocardial infarction (NSTEMI)

The ICD-10-CM codes for acute myocardial infarction (AMI) identify the site, such as anterolateral wall or true posterior wall. Subcategories I21.0-I21.2 and code I21.4 are used for ST elevation myocardial infarction (STEMI). Code I21.4, Non-ST elevation (NSTEMI) myocardial infarction, is used for non ST elevation myocardial infarction (NSTEMI) and nontransmural MIs.

2) Acute myocardial infarction, unspecified

Code I21.3, ST elevation (STEMI) myocardial infarction of unspecified site, is the default for the unspecified term acute myocardial infarction. If only STEMI or transmural MI without the site is documented, query the provider as to the site, or assign code I21.3.

3) AMI documented as nontransmural or subendocardial but site provided

If an AMI is documented as nontransmural or subendocardial, but the site is provided, it is still coded as a subendocardial AMI. If NSTEMI evolves to STEMI, assign the STEMI code.

If STEMI converts to NSTEMI due to thrombolytic therapy, it is still coded as STEMI.

See Section I.C.21.3 for information on coding status post administration of tPA in a different facility within the last 24 hours.

4) Subsequent acute myocardial infarction

A code from category I22, Subsequent ST elevation (STEMI) and non ST elevation (NSTEMI) myocardial infarction, is to be used when a patient who has suffered an AMI has a new AMI within the 4 week time frame of the initial AMI. A code from category I22 must be used in conjunction with a code from category I21.

The sequencing of the I22 and I21 codes depends on the circumstances of the encounter. Should a patient who is in the hospital due to an AMI have a subsequent AMI while still in the hospital code I21 would be sequenced first as the reason for admission, with code I22 sequenced as a secondary code. Should a patient have a subsequent AMI after discharge for care of an initial AMI, and the reason for admission is the subsequent AMI, the I22 code should be sequenced first followed by the I21. An I21 code must accompany an I22 code to identify the site of the initial AMI, and to indicate that the patient is still within the 4 week time frame of healing from the initial AMI.

The guidelines for assigning the correct I22 code are the same as for the initial AMI.

10. Chapter 10: Diseases of Respiratory System (J00-J99)

a. Chronic Obstructive Pulmonary Disease [COPD] and Asthma

1) Acute exacerbation of chronic obstructive bronchitis and asthma

The codes in categories J44 and J45 distinguish between uncomplicated cases and those in acute exacerbation. An acute exacerbation is a worsening or a decompensation of a chronic condition. An acute exacerbation is not equivalent to an infection superimposed on a chronic condition, though an exacerbation may be triggered by an infection.

b. Acute Respiratory Failure

1) Acute respiratory failure as principal diagnosis

Code J96.0, Acute respiratory failure, or code J96.2, Acute and chronic respiratory failure, may be assigned as a principal diagnosis when it is the condition established after study to be chiefly responsible for occasioning the admission to the hospital, and the selection is supported by the Alphabetic Index and Tabular List. However, chapter-specific coding guidelines (such as obstetrics, poisoning, HIV, newborn) that provide sequencing direction take precedence.

2) Acute respiratory failure as secondary diagnosis

Respiratory failure may be listed as a secondary diagnosis if it occurs after admission, or if it is present on admission, but does not meet the definition of principal diagnosis.

3) Sequencing of acute respiratory failure and another acute condition

When a patient is admitted with respiratory failure and another acute condition, (e.g., myocardial infarction, cerebrovascular accident, aspiration pneumonia), the principal diagnosis will not be the same in every situation. This applies whether the other acute condition is a respiratory or nonrespiratory condition. Selection of the principal diagnosis will be dependent on the circumstances of admission. If both the respiratory failure and the other acute condition are equally responsible for occasioning the admission to the hospital, and there are no chapter-specific sequencing rules, the guideline regarding two or more diagnoses that equally meet the definition for principal diagnosis (*Section II, C.*) may be applied in these situations.

If the documentation is not clear as to whether acute respiratory failure and another condition are equally responsible for occasioning the admission, query the provider for clarification.

c. Influenza due to *certain identified influenza influenza viruses*

Code only confirmed cases of avian influenza (**code J09.0-, Influenza due to identified avian influenza virus**) or novel H1N1 or swine flu, **code J09.1-**. This is an exception to the hospital inpatient guideline Section II, H. (Uncertain Diagnosis).

In this context, “confirmation” does not require documentation of positive laboratory testing specific for avian **or novel H1N1 (H1N1 or swine flu)** influenza. However, coding should be based on the provider’s diagnostic statement that the patient has avian influenza.

If the provider records “suspected or possible or probable avian influenza,” the appropriate influenza code from category J10, Influenza due to other influenza virus, should be assigned. **A code from category J09, Influenza due to certain identified influenza viruses, should not be assigned.**

11. Chapter 11: Diseases of Digestive System (K00-K94)

Reserved for future guideline expansion

12. Chapter 12: Diseases of Skin and Subcutaneous Tissue (L00-L99)

a. Pressure ulcer stage codes

1) Pressure ulcer stages

Codes from category L89, Pressure ulcer, are combination codes that identify the site of the pressure ulcer as well as the stage of the ulcer.

The ICD-10-CM classifies pressure ulcer stages based on severity, which is designated by stages 1-4, unspecified stage and unstageable .

Assign as many codes from category L89 as needed to identify all the pressure ulcers the patient has, if applicable.

2) Unstageable pressure ulcers

Assignment of the code for unstageable pressure ulcer (L89.--0) should be based on the clinical documentation. These codes are used for pressure ulcers whose stage cannot be clinically determined (e.g., the ulcer is covered by eschar or has been treated with a skin or muscle graft) and pressure ulcers that are documented as deep tissue injury but not documented as due to trauma. This code should not be confused with the codes for unspecified stage (L89.--9). When there is no documentation regarding the stage of the pressure ulcer, assign the appropriate code for unspecified stage (L89.--9).

- 3) Documented pressure ulcer stage**
Assignment of the pressure ulcer stage code should be guided by clinical documentation of the stage or documentation of the terms found in the index. For clinical terms describing the stage that are not found in the index, and there is no documentation of the stage, the provider should be queried.
- 4) Patients admitted with pressure ulcers documented as healed**
No code is assigned if the documentation states that the pressure ulcer is completely healed.
- 5) Patients admitted with pressure ulcers documented as healing**
Pressure ulcers described as healing should be assigned the appropriate pressure ulcer stage code based on the documentation in the medical record. If the documentation does not provide information about the stage of the healing pressure ulcer, assign the appropriate code for unspecified stage.

If the documentation is unclear as to whether the patient has a current (new) pressure ulcer or if the patient is being treated for a healing pressure ulcer, query the provider.
- 6) Patient admitted with pressure ulcer evolving into another stage during the admission**
If a patient is admitted with a pressure ulcer at one stage and it progresses to a higher stage, assign the code for the highest stage reported for that site.

13. Chapter 13: Diseases of the Musculoskeletal System and Connective Tissue (M00-M99)

a. Site and laterality

Most of the codes within Chapter 13 have site and laterality designations. The site represents either the bone, joint or the muscle involved. For some conditions where more than one bone, joint or muscle is usually involved, such as osteoarthritis, there is a “multiple sites” code available. For categories where no multiple site code is provided and more than one bone, joint or muscle is involved, multiple codes should be used to indicate the different sites involved.

1) Bone versus joint

For certain conditions, the bone may be affected at the upper or lower end, (e.g., avascular necrosis of bone, M87, Osteoporosis, M80, M81). Though the portion of the bone affected may be at the joint, the site designation will be the bone, not the joint.

b. Acute traumatic versus chronic or recurrent musculoskeletal conditions

Many musculoskeletal conditions are a result of previous injury or trauma to a site, or are recurrent conditions. Bone, joint or muscle conditions that are the result of a healed injury are usually found in chapter 13. Recurrent bone, joint or muscle conditions are also usually found in chapter 13. Any current, acute injury should be coded to the appropriate injury code from chapter 19. Chronic or recurrent conditions should generally be coded with a code from chapter 13. If it is difficult to determine from the documentation in the record which code is best to describe a condition, query the provider.

c. Coding of Pathologic Fractures

7th character A is for use as long as the patient is receiving active treatment for the fracture. Examples of active treatment are: surgical treatment, emergency department encounter, evaluation and treatment by a new physician. 7th character, D is to be used for encounters after the patient has completed active treatment. The other 7th characters, listed under each subcategory in the Tabular List, are to be used for subsequent encounters for treatment of problems associated with the healing, such as malunions, nonunions, and sequelae.

Care for complications of surgical treatment for fracture repairs during the healing or recovery phase should be coded with the appropriate complication codes.

See Section I.C.19. Coding of traumatic fractures.

d. Osteoporosis

Osteoporosis is a systemic condition, meaning that all bones of the musculoskeletal system are affected. Therefore, site is not a component of the codes under category M81, Osteoporosis without current pathological fracture. The site codes under category M80, Osteoporosis with current pathological fracture, identify the site of the fracture, not the osteoporosis.

1) Osteoporosis without pathological fracture

Category M81, Osteoporosis without current pathological fracture, is for use for patients with osteoporosis who do not

currently have a pathologic fracture due to the osteoporosis, even if they have had a fracture in the past. For patients with a history of osteoporosis fractures, status code Z87.31, Personal history of osteoporosis fracture, should follow the code from M81.

2) Osteoporosis with current pathological fracture

Category M80, Osteoporosis with current pathological fracture, is for patients who have a current pathologic fracture at the time of an encounter. The codes under M80 identify the site of the fracture. A code from category M80, not a traumatic fracture code, should be used for any patient with known osteoporosis who suffers a fracture, even if the patient had a minor fall or trauma, if that fall or trauma would not usually break a normal, healthy bone.

14. Chapter 14: Diseases of Genitourinary System (N00-N99)

a. Chronic kidney disease

1) Stages of chronic kidney disease (CKD)

The ICD-10-CM classifies CKD based on severity. The severity of CKD is designated by stages I-V. Stage II, code N18.2, equates to mild CKD; stage III, code N18.3, equates to moderate CKD; and stage IV, code N18.4, equates to severe CKD. Code N18.6, End stage renal disease (ESRD), is assigned when the provider has documented end-stage-renal disease (ESRD).

If both a stage of CKD and ESRD are documented, assign code N18.6 only.

2) Chronic kidney disease and kidney transplant status

Patients who have undergone kidney transplant may still have some form of **chronic kidney disease (CKD)** because the kidney transplant may not fully restore kidney function. Therefore, the presence of CKD alone does not constitute a transplant complication. Assign the appropriate N18 code for the patient's stage of CKD and code Z94.0, Kidney transplant status. If a transplant complication such as failure or rejection **or other transplant complication** is documented, see section I.C.19.g for information on coding complications of a kidney transplant. If the documentation is unclear as to whether the patient has a complication of the transplant, query the provider.

3) Chronic kidney disease with other conditions

Patients with CKD may also suffer from other serious conditions, most commonly diabetes mellitus and hypertension. The sequencing of the CKD code in relationship to codes for other contributing conditions is based on the conventions in the Tabular List.

See I.C.9. Hypertensive chronic kidney disease.

See I.C.19. Chronic kidney disease and kidney transplant complications.

15. Chapter 15: Pregnancy, Childbirth, and the Puerperium (O00-O9A)

a. General Rules for Obstetric Cases

1) Codes from chapter 15 and sequencing priority

Obstetric cases require codes from chapter 15, codes in the range O00-**O9A**, Pregnancy, Childbirth, and the Puerperium. Chapter 15 codes have sequencing priority over codes from other chapters. Additional codes from other chapters may be used in conjunction with chapter 15 codes to further specify conditions. Should the provider document that the pregnancy is incidental to the encounter, then code Z33.1, Pregnant state, incidental, should be used in place of any chapter 15 codes. It is the provider's responsibility to state that the condition being treated is not affecting the pregnancy.

2) Chapter 15 codes used only on the maternal record

Chapter 15 codes are to be used only on the maternal record, never on the record of the newborn.

3) Final character for trimester

The majority of codes in Chapter 15 have a final character indicating the trimester of pregnancy. The timeframes for the trimesters are indicated at the beginning of the chapter. If trimester is not a component of a code it is because the condition always occurs in a specific trimester, or the concept of trimester of pregnancy is not applicable. Certain codes have characters for only certain trimesters because the condition does not occur in all trimesters, but it may occur in more than just one.

Assignment of the final character for trimester should be based on the trimester for the current admission/encounter. This applies to the assignment of trimester for pre-existing conditions as well as those that develop during or are due to the pregnancy.

Whenever delivery occurs during the current admission, and there is an “in childbirth” option for the obstetric complication being coded, the “in childbirth” code should be assigned.

4) Selection of trimester for inpatient admissions *that encompass more than one trimesters*

In instances when a patient is admitted to a hospital for complications of pregnancy **during one trimester** and remains in the hospital **into a subsequent trimester, the trimester character** for the antepartum complication code should be assigned on the basis of the trimester when the complication developed, **not the trimester of the discharge. If the condition developed prior to the current admission/encounter or represents a pre-existing condition, the trimester character for the trimester at the time of the admission/encounter should be assigned.**

5) Unspecified trimester

Each category that includes codes for trimester has a code for “unspecified trimester.” The “unspecified trimester” code should rarely be used, such as when the documentation in the record is insufficient to determine the trimester and it is not possible to obtain clarification.

b. Selection of OB Principal or First-listed Diagnosis

1) Routine outpatient prenatal visits

For routine outpatient prenatal visits when no complications are present, a code from category Z34, Encounter for supervision of normal pregnancy, should be used as the first-listed diagnosis. These codes should not be used in conjunction with chapter 15 codes.

2) Prenatal outpatient visits for high-risk patients

For routine prenatal outpatient visits for patients with high-risk pregnancies, a code from category O09, Supervision of high-risk pregnancy, should be used as the first-listed diagnosis. Secondary chapter 15 codes may be used in conjunction with these codes if appropriate.

3) Episodes when no delivery occurs

In episodes when no delivery occurs, the principal diagnosis should correspond to the principal complication of the pregnancy which necessitated the encounter. Should more than one complication exist, all of which are treated or monitored, any of the complications codes may be sequenced first.

4) When a delivery occurs

When a delivery occurs, the principal diagnosis should correspond to the main circumstances or complication of the delivery. In cases of cesarean delivery, the selection of the principal **diagnosis should be the condition established after study that was responsible for the patient's admission. If the patient was admitted with a condition that resulted in the performance of a cesarean procedure, that condition should be selected as the principal diagnosis. If the reason for the admission/encounter was unrelated to the condition resulting in the cesarean delivery, the condition related to the reason for the admission/encounter should be selected as the principal diagnosis, even if a cesarean was performed.**

5) Outcome of delivery

A code from category Z37, Outcome of delivery, should be included on every maternal record when a delivery has occurred. These codes are not to be used on subsequent records or on the newborn record.

c. Pre-existing conditions versus conditions due to the pregnancy

Certain categories in Chapter 15 distinguish between conditions of the mother that existed prior to pregnancy (pre-existing) and those that are a direct result of pregnancy. When assigning codes from Chapter 15, it is important to assess if a condition was pre-existing prior to pregnancy or developed during or due to the pregnancy in order to assign the correct code.

Categories that do not distinguish between pre-existing and pregnancy-related conditions may be used for either. It is acceptable to use codes specifically for the puerperium with codes complicating pregnancy and childbirth if a condition arises postpartum during the delivery encounter.

d. Pre-existing hypertension in pregnancy

Category O10, Pre-existing hypertension complicating pregnancy, childbirth and the puerperium, includes codes for hypertensive heart and hypertensive chronic kidney disease. When assigning one of the O10 codes that includes hypertensive heart disease or hypertensive chronic kidney disease, it is necessary to add a secondary code from the appropriate hypertension category to specify the type of heart failure or chronic kidney disease.

See Section I.C.9. Hypertension.

e. Fetal Conditions Affecting the Management of the Mother

1) Codes from categories O35 and O36

Codes from categories O35, Maternal care for known or suspected fetal abnormality and damage, and O36, Maternal care for other fetal problems, are assigned only when the fetal condition is actually responsible for modifying the management of the mother, i.e., by requiring diagnostic studies, additional observation, special care, or termination of pregnancy. The fact that the fetal condition exists does not justify assigning a code from this series to the mother's record.

2) In utero surgery

In cases when surgery is performed on the fetus, a diagnosis code from category O35, Maternal care for known or suspected fetal abnormality and damage, should be assigned identifying the fetal condition. Assign the appropriate procedure code for the procedure performed.

No code from Chapter 16, the perinatal codes, should be used on the mother's record to identify fetal conditions. Surgery performed in utero on a fetus is still to be coded as an obstetric encounter.

f. HIV Infection in Pregnancy, Childbirth and the Puerperium

During pregnancy, childbirth or the puerperium, a patient admitted because of an HIV-related illness should receive a principal diagnosis from subcategory O98.7-, Human immunodeficiency [HIV] disease complicating pregnancy, childbirth and the puerperium, followed by the code(s) for the HIV-related illness(es).

Patients with asymptomatic HIV infection status admitted during pregnancy, childbirth, or the puerperium should receive codes of O98.7- and Z21, Asymptomatic human immunodeficiency virus [HIV] infection status.

g. Diabetes mellitus in pregnancy

Diabetes mellitus is a significant complicating factor in pregnancy. Pregnant women who are diabetic should be assigned a code O24, Diabetes mellitus in pregnancy, childbirth, and the puerperium, first, followed by the appropriate diabetes code(s) (E08-E13) from Chapter 4.

h. Long term use of insulin

Code Z79.4, Long-term (current) use of insulin, should also be assigned if the diabetes mellitus is being treated with insulin.

i. Gestational (pregnancy induced) diabetes

Gestational (pregnancy induced) diabetes can occur during the second and third trimester of pregnancy in women who were not diabetic prior to pregnancy. Gestational diabetes can cause complications in the pregnancy similar to those of pre-existing diabetes mellitus. It also puts the woman at greater risk of developing diabetes after the pregnancy. Codes for gestational diabetes are in subcategory O24.4, Gestational diabetes mellitus. No other code from category O24, Diabetes mellitus in pregnancy, childbirth, and the puerperium, should be used with a code from O24.4

The codes under subcategory O24.4 include diet controlled and insulin controlled. If a patient with gestational diabetes is treated with both diet and insulin, only the code for insulin-controlled is required. Code **Z79.4**, Long-term (current) use of insulin, should **not** be assigned **with codes from subcategory O24.4**.

An abnormal glucose tolerance in pregnancy is assigned a code from subcategory O99.81, Abnormal glucose complicating pregnancy, childbirth, and the puerperium.

j. Sepsis and septic shock complicating abortion, pregnancy, childbirth and the puerperium

When assigning a chapter 15 code for sepsis complicating abortion, pregnancy, childbirth, and the puerperium, a code for the specific type of infection should be assigned as an additional diagnosis. If severe sepsis is present, a code from subcategory R65.2, Severe sepsis, and code(s) for associated organ dysfunction(s) should also be assigned as additional diagnoses.

k. Puerperal sepsis

Code O85, Puerperal sepsis, should be assigned with a secondary code to identify the causal organism (e.g., for a bacterial infection, assign a code from category B95-B96, Bacterial infections in conditions classified elsewhere). A code from category A40, Streptococcal sepsis, or A41, Other sepsis, should not be used for puerperal sepsis. If applicable, use additional codes to identify severe sepsis (R65.2-) and any associated acute organ dysfunction.

l. Alcohol and tobacco use during pregnancy, childbirth and the puerperium

1) Alcohol use during pregnancy, childbirth and the puerperium

Codes under subcategory O99.31, Alcohol use complicating pregnancy, childbirth, and the puerperium, should be assigned for any pregnancy case when a mother uses alcohol during the pregnancy or postpartum. A secondary code from category F10, Alcohol related disorders, should also be assigned.

2) Tobacco use during pregnancy, childbirth and the puerperium

Codes under subcategory O99.33, Smoking (tobacco) complicating pregnancy, childbirth, and the puerperium, should be assigned for any pregnancy case when a mother uses any type of tobacco product during the pregnancy or postpartum. A secondary code from category F17, Nicotine dependence, or code Z72.0, Tobacco use, should also be assigned.

m. Poisoning, toxic effects, adverse effects and underdosing in a pregnant patient

A code from subcategory O9A.2, Injury, poisoning and certain other consequences of external causes complicating pregnancy, childbirth, and the puerperium, should be sequenced first, followed by the appropriate poisoning, toxic effect, adverse effect or underdosing code, and then the additional code(s) that specifies the condition caused by the poisoning, toxic effect, adverse effect or underdosing.

See Section I.C.19. Adverse effects, poisoning, underdosing and toxic effects.

n. Normal Delivery, Code O80

1) Encounter for full term uncomplicated delivery

Code O80 should be assigned when a woman is admitted for a full-term normal delivery and delivers a single, healthy infant without any complications antepartum, during the delivery, or postpartum during the delivery episode. Code O80 is always a principal diagnosis. It is not to be used if any other code from chapter 15 is needed to describe a current complication of the antenatal, delivery, or perinatal period. Additional codes from other chapters may be used with code O80 if they are not related to or are in any way complicating the pregnancy.

2) Uncomplicated delivery with resolved antepartum complication

Code O80 may be used if the patient had a complication at some point during the pregnancy, but the complication is not present at the time of the admission for delivery.

3) Outcome of delivery for O80

Z37.0, Single live birth, is the only outcome of delivery code appropriate for use with O80.

o. The Peripartum and Postpartum Periods

1) Peripartum and Postpartum periods

The postpartum period begins immediately after delivery and continues for six weeks following delivery. The peripartum period is defined as the last month of pregnancy to five months postpartum.

2) Peripartum and postpartum complication

A postpartum complication is any complication occurring within the six-week period.

3) Pregnancy-related complications after 6 week period

Chapter 15 codes may also be used to describe pregnancy-related complications after the peripartum or postpartum period if the provider documents that a condition is pregnancy related.

4) Admission for routine postpartum care following delivery outside hospital

When the mother delivers outside the hospital prior to admission and is admitted for routine postpartum care and no complications are noted, code Z39.0, Encounter for care and examination of mother immediately after delivery, should be assigned as the principal diagnosis.

5) Pregnancy associated cardiomyopathy

Pregnancy associated cardiomyopathy, code O90.3, is unique in that it may be diagnosed in the third trimester of pregnancy but may continue to progress months after delivery. For this reason, it is referred to as peripartum cardiomyopathy. Code O90.3 is only for use when the cardiomyopathy develops as a result of pregnancy in a woman who did not have pre-existing heart disease.

p. Code O94, Sequelae of complication of pregnancy, childbirth, and the puerperium

1) Code O94

Code O94, Sequelae of complication of pregnancy, childbirth, and the puerperium, is for use in those cases when an initial complication of a pregnancy develops a sequelae requiring care or treatment at a future date.

2) After the initial postpartum period

This code may be used at any time after the initial postpartum period.

3) Sequencing of Code O94

This code, like all late effect codes, is to be sequenced following the code describing the sequelae of the complication.

q. Abortions

1) Abortion with Liveborn Fetus

When an attempted termination of pregnancy results in a liveborn fetus assign a code from subcategory O60.1, Preterm labor with preterm delivery, **and a code from** category Z37, Outcome of Delivery. The procedure code for the attempted termination of pregnancy should also be assigned.

2) Retained Products of Conception following an abortion

Subsequent encounters for retained products of conception following a spontaneous abortion or elective termination of

pregnancy are assigned the appropriate code from category O03, Spontaneous abortion, or code Z33.2, Encounter for elective termination of pregnancy. This advice is appropriate even when the patient was discharged previously with a discharge diagnosis of complete abortion.

16. Chapter 16: Newborn (Perinatal) Guidelines (P00-P96)

For coding and reporting purposes the perinatal period is defined as before birth through the 28th day following birth. The following guidelines are provided for reporting purposes

a. General Perinatal Rules

1) Use of Chapter 16 Codes

Codes in this chapter are never for use on the maternal record. Codes from Chapter 15, the obstetric chapter, are never permitted on the newborn record. Chapter 16 code may be used throughout the life of the patient if the condition is still present.

2) Principal Diagnosis for Birth Record

When coding the birth episode in a newborn record, assign a code from category Z38, Liveborn according to place of birth and type of delivery, as the principal diagnosis. A code from category Z38 is assigned only once, to a newborn at the time of birth. If a newborn is transferred to another institution, a code from category Z38 should not be used at the receiving hospital.

A code from category Z38 is used only on the newborn record, not on the mother's record.

3) Use of Codes from other Chapters with Codes from Chapter 16

Codes from other chapters may be used with codes from chapter 16 if the codes from the other chapters provide more specific detail. Codes for signs and symptoms may be assigned when a definitive diagnosis has not been established. If the reason for the encounter is a perinatal condition, the code from chapter 16 should be sequenced first.

4) Use of Chapter 16 Codes after the Perinatal Period

Should a condition originate in the perinatal period, and continue throughout the life of the patient, the perinatal code should continue to be used regardless of the patient's age.

5) Birth process or community acquired conditions

If a newborn has a condition that may be either due to the birth process or community acquired and the documentation does not indicate which it is, the default is due to the birth process and the code from Chapter 16 should be used. If the condition is community-acquired, a code from Chapter 16 should not be assigned.

6) Code all clinically significant conditions

All clinically significant conditions noted on routine newborn examination should be coded. A condition is clinically significant if it requires:

- clinical evaluation; or
- therapeutic treatment; or
- diagnostic procedures; or
- extended length of hospital stay; or
- increased nursing care and/or monitoring; or
- has implications for future health care needs

Note: The perinatal guidelines listed above are the same as the general coding guidelines for “additional diagnoses”, except for the final point regarding implications for future health care needs. Codes should be assigned for conditions that have been specified by the provider as having implications for future health care needs.

b. Observation and Evaluation of Newborns for Suspected Conditions not Found

Assign a code from categories P00-P04 to identify those instances when a healthy newborn is evaluated for a suspected condition that is determined after study not to be present. Do not use a code from categories P00-P04 when the patient has identified signs or symptoms of a suspected problem; in such cases, code the sign or symptom.

c. Coding Additional Perinatal Diagnoses

1) Assigning codes for conditions that require treatment

Assign codes for conditions that require treatment or further investigation, prolong the length of stay, or require resource utilization.

2) Codes for conditions specified as having implications for future health care needs

Assign codes for conditions that have been specified by the provider as having implications for future health care needs.

Note: This guideline should not be used for adult patients.

d. Prematurity and Fetal Growth Retardation

Providers utilize different criteria in determining prematurity. A code for prematurity should not be assigned unless it is documented. Assignment of codes in categories P05, Disorders of newborn related to slow fetal growth and fetal malnutrition, and P07, Disorders of newborn related to short gestation and low birth weight, not elsewhere classified, should be based on the recorded birth weight and estimated gestational age. Codes from category P05 should not be assigned with codes from category P07.

When both birth weight and gestational age are available, two codes from category P07 should be assigned, with the code for birth weight sequenced before the code for gestational age.

e. Low birth weight and immaturity status

Codes from subcategory Z91.7, Low birth weight and immaturity status, are for use as personal status codes for a child or adult who was premature or had a low birth weight as a newborn and this is affecting the patient's current health status.

See Section I.C.21. Factors influencing health status and contact with health services, Status.

f. Bacterial Sepsis of Newborn

Category P36, Bacterial sepsis of newborn, includes congenital sepsis. If a perinate is documented as having sepsis without documentation of congenital or community acquired, the default is congenital and a code from category P36 should be assigned. If the P36 code includes the causal organism, an additional code from category B95, Streptococcus, Staphylococcus, and Enterococcus as the cause of diseases classified elsewhere, or B96, Other bacterial agents as the cause of diseases classified elsewhere, should **not** be assigned. If the P36 code does not include the causal organism, assign an additional code from category B96. If applicable, use additional codes to identify severe sepsis (R65.2-) and any associated acute organ dysfunction.

g. Stillbirth

Code P95, Stillbirth, is only for use in institutions that maintain separate records for stillbirths. No other code should be used with P95. Code P95 should not be used on the mother's record.

17. Chapter 17: Congenital malformations, deformations, and chromosomal abnormalities (Q00-Q99)

Assign an appropriate code(s) from categories Q00-Q99, Congenital malformations, deformations, and chromosomal abnormalities when a malformation/deformation or chromosomal abnormality is documented. A malformation/deformation/or chromosomal abnormality may be the principal/first listed diagnosis on a record or a secondary diagnosis.

When a malformation/deformation/or chromosomal abnormality does not have a unique code assignment, assign additional code(s) for any manifestations that may be present.

When the code assignment specifically identifies the malformation/deformation/or chromosomal abnormality, manifestations that are an inherent component of the anomaly should not be coded separately. Additional codes should be assigned for manifestations that are not an inherent component.

Codes from Chapter 17 may be used throughout the life of the patient. If a congenital malformation or deformity has been corrected, a personal history code should be used to identify the history of the malformation or deformity. Although present at birth, malformation/deformation/or chromosomal abnormality may not be identified until later in life. Whenever the condition

is diagnosed by the physician, it is appropriate to assign a code from codes Q00-Q99.

For the birth admission, the appropriate code from category Z38, Liveborn infants, according to place of birth and type of delivery, should be sequenced as the principal diagnosis, followed by any congenital anomaly codes, Q00-Q89.

18. Chapter 18: Symptoms, signs, and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Chapter 18 includes symptoms, signs, abnormal results of clinical or other investigative procedures, and ill-defined conditions regarding which no diagnosis classifiable elsewhere is recorded. Signs and symptoms that point to a **specific** diagnosis have been assigned to a category in other chapters of the classification.

a. Use of symptom codes

Codes that describe symptoms and signs are acceptable for reporting purposes when a related definitive diagnosis has not been established (confirmed) by the provider.

b. Use of a symptom code with a definitive diagnosis code

Codes for signs and symptoms may be reported in addition to a related definitive diagnosis when the sign or symptom is not routinely associated with that diagnosis, such as the various signs and symptoms associated with complex syndromes. The definitive diagnosis code should be sequenced before the symptom code.

Signs or symptoms that are associated routinely with a disease process should not be assigned as additional codes, unless otherwise instructed by the classification.

c. Combination codes that include symptoms

ICD-10-CM contains a number of combination codes that identify both the definitive diagnosis and common symptoms of that diagnosis. When using one of these combination codes, an additional code should not be assigned for the symptom.

d. Repeated falls

Code R29.6, Repeated falls, is for use for encounters when a patient has recently fallen and the reason for the fall is being investigated.

Code Z91.81, History of falling, is for use when a patient has fallen in the past and is at risk for future falls. When appropriate, both codes R29.6 and Z91.81 may be assigned together.

e. Glasgow coma scale

The Glasgow coma scale codes (R40.2-) can be used in conjunction with traumatic brain injury codes or sequelae of cerebrovascular accident codes. These codes are primarily for use by trauma registries, but they may be used in any setting where this information is collected. The coma scale codes should be sequenced after the diagnosis code(s).

These codes, one from each subcategory, are needed to complete the scale. The 7th character indicates when the scale was recorded. The 7th character should match for all three codes.

At a minimum, report the initial score documented on presentation at your facility. This may be a score from the emergency medicine technician (EMT) or in the emergency department. If desired, a facility may choose to capture multiple Glasgow coma scale scores.

f. Functional quadriplegia

Functional quadriplegia (code R53.2) is the lack of ability to use one's limbs or to ambulate due to extreme debility. It is not associated with neurologic deficit or injury, and code R53.2 should not be used for cases of neurologic quadriplegia. It should only be assigned if functional quadriplegia is specifically documented in the medical record.

g. SIRS due to Non-Infectious Process

The systemic inflammatory response syndrome (SIRS) can develop as a result of certain non-infectious disease processes, such as trauma, malignant neoplasm, or pancreatitis. When SIRS is documented with a noninfectious condition, and no subsequent infection is documented, the code for the underlying condition, such as an injury, should be assigned, followed by code R65.10, Systemic inflammatory response syndrome (SIRS) of non-infectious origin without acute organ dysfunction, or code R65.11, Systemic inflammatory response syndrome (SIRS) of non-infectious origin with acute organ dysfunction. If an associated acute organ dysfunction is documented, the appropriate code(s) for the specific type of organ dysfunction(s) should be assigned in addition to code R65.11. If acute organ dysfunction is documented, but it cannot be determined if the acute

organ dysfunction is associated with SIRS or due to another condition (e.g., directly due to the trauma), the provider should be queried.

g. Death NOS

Code R99, Ill-defined and unknown cause of mortality, is only for use in the very limited circumstance when a patient who has already died is brought into an emergency department or other healthcare facility and is pronounced dead upon arrival. It does not represent the discharge disposition of death.

19. Chapter 19: Injury, poisoning, and certain other consequences of external causes (S00-T88)

a. Code Extensions

Most categories in chapter 19 have 7th character extensions that are required for each applicable code. Most categories in this chapter have three extensions (with the exception of fractures): A, initial encounter, D, subsequent encounter and S, sequela.

Extension “A”, initial encounter is used while the patient is receiving active treatment for the injury. Examples of active treatment are: surgical treatment, emergency department encounter, and evaluation and treatment by a new physician.

Extension “D” subsequent encounter is used for encounters after the patient has received active treatment of the injury and is receiving routine care for the injury during the healing or recovery phase. Examples of subsequent care are: cast change or removal, removal of external **or** internal fixation device, medication adjustment, other aftercare and follow up visits following injury treatment.

The aftercare Z codes should not be used for aftercare for injuries. For aftercare of an injury, assign the acute injury code with the 7th character “D” (subsequent encounter).

Extension “S”, sequela, is for use for complications or conditions that arise as a direct result of an injury, such as scar formation after a burn. The scars are sequelae of the burn. When using extension “S”, it is necessary to use both the injury code that precipitated the sequela and the code for the sequela itself. The “S” is added only to the injury code, not the sequela code. The “S” extension identifies the injury responsible for the sequela. The specific type of sequela (e.g. scar) is sequenced first, followed by the injury code.

b. Coding of Injuries

When coding injuries, assign separate codes for each injury unless a combination code is provided, in which case the combination code is assigned. Multiple injury codes are provided in ICD-10-CM, but should not be assigned unless information for a more specific code is not available. These **traumatic injury** codes (S00-T14.9) are not to be used for normal, healing surgical wounds or to identify complications of surgical wounds.

The code for the most serious injury, as determined by the provider and the focus of treatment, is sequenced first.

1) Superficial injuries

Superficial injuries such as abrasions or contusions are not coded when associated with more severe injuries of the same site.

2) Primary injury with damage to nerves/blood vessels

When a primary injury results in minor damage to peripheral nerves or blood vessels, the primary injury is sequenced first with additional code(s) for injuries to nerves and spinal cord (such as category S04), and/or injury to blood vessels (such as category S15). When the primary injury is to the blood vessels or nerves, that injury should be sequenced first.

c. Coding of Traumatic Fractures

The principles of multiple coding of injuries should be followed in coding fractures. Fractures of specified sites are coded individually by site in accordance with both the provisions within categories S02, S12, S22, S32, S42, **S49**, S52, **S59**, S62, S72, **S79**, S82, **S89**, S92 and the level of detail furnished by medical record content.

A fracture not indicated as open or closed should be coded to closed. A fracture not indicated whether displaced or not displaced should be coded to displaced.

More specific guidelines are as follows:

1) Initial vs. Subsequent Encounter for Fractures

Traumatic fractures are coded using the appropriate 7th character extension for initial encounter (A, B, C) while the patient is receiving active treatment for the fracture. Examples of active treatment are: surgical treatment, emergency department encounter, and evaluation and treatment by a new physician.

Fractures are coded using the appropriate 7th character extension for subsequent care for encounters after the patient has completed active treatment of the fracture and is receiving routine care for the fracture during the healing or recovery phase. Examples of fracture aftercare are: cast change or removal, removal of external or internal fixation device, medication adjustment, and follow-up visits following fracture treatment.

Care for complications of surgical treatment for fracture repairs during the healing or recovery phase should be coded with the appropriate complication codes.

Care of complications of fractures, such as malunion and nonunion, should be reported with the appropriate 7th character extensions for subsequent care with nonunion (K, M, N,) or subsequent care with malunion (P, Q, R).

A code from category M80, not a traumatic fracture code, should be used for any patient with known osteoporosis who suffers a fracture, **even if the patient had a minor fall or trauma, if that fall or trauma would not usually break a normal, healthy bone.**

See Section I.C.13. Osteoporosis.

The aftercare Z codes should not be used for aftercare for injuries. For aftercare of an injury, assign the acute injury code with the 7th character “D” (subsequent encounter).

2) **Multiple fractures sequencing**

Multiple fractures are sequenced in accordance with the severity of the fracture. The provider should be asked to list the fracture diagnoses in the order of severity.

d. **Coding of Burns and Corrosions**

The ICD-10-CM distinguishes between burns and corrosions. The burn codes are for thermal burns, except sunburns, that come from a heat source, such as a fire or hot appliance. The burn codes are also for burns resulting from electricity and radiation. Corrosions are burns due to chemicals. The guidelines are the same for burns and corrosions.

Current burns (T20-T25) are classified by depth, extent and by agent (X code). Burns are classified by depth as first degree (erythema), second degree (blistering), and third degree (full-thickness

involvement). Burns of the eye and internal organs (T26-T28) are classified by site, but not by degree.

1) Sequencing of burn and related condition codes

Sequence first the code that reflects the highest degree of burn when more than one burn is present.

- a. When the reason for the admission or encounter is for treatment of external multiple burns, sequence first the code that reflects the burn of the highest degree.
- b. When a patient has both internal and external burns, the circumstances of admission govern the selection of the principal diagnosis or first-listed diagnosis.
- c. When a patient is admitted for burn injuries and other related conditions such as smoke inhalation and/or respiratory failure, the circumstances of admission govern the selection of the principal or first-listed diagnosis.

2) Burns of the same local site

Classify burns of the same local site (three-digit category level, T20-T28) but of different degrees to the subcategory identifying the highest degree recorded in the diagnosis.

3) Non-healing burns

Non-healing burns are coded as acute burns.
Necrosis of burned skin should be coded as a non-healed burn.

4) Infected Burn

For any documented infected burn site, use an additional code for the infection.

5) Assign separate codes for each burn site

When coding burns, assign separate codes for each burn site. Category T30, Burn and corrosion, body region unspecified is extremely vague and should rarely be used.

6) Burns and Corrosions Classified According to Extent of Body Surface Involved

Assign codes from category T31, Burns classified according to extent of body surface involved, or T32, Corrosions classified according to extent of body surface involved, when the site of the burn is not specified or when there is a need for additional data. It is advisable to use category T31 as additional coding when needed to provide data for evaluating burn mortality, such as that needed by burn units. It is also advisable to use category T31 as an additional code for reporting purposes

when there is mention of a third-degree burn involving 20 percent or more of the body surface.

Categories T31 and T32 are based on the classic “rule of nines” in estimating body surface involved: head and neck are assigned nine percent, each arm nine percent, each leg 18 percent, the anterior trunk 18 percent, posterior trunk 18 percent, and genitalia one percent. Providers may change these percentage assignments where necessary to accommodate infants and children who have proportionately larger heads than adults, and patients who have large buttocks, thighs, or abdomen that involve burns.

7) Encounters for treatment of late effects of burns

Encounters for the treatment of the late effects of burns or corruptions (i.e., scars or joint contractures) should be coded with a burn or corrosion code with the 7th character “S” or sequela.

8) Sequelae with a late effect code and current burn

When appropriate, both a code for a current burn or corrosion with 7th character extension “A” or “D” and a burn or corrosion code with extension “S” may be assigned on the same record (when both a current burn and sequelae of an old burn exist). Burns and corruptions do not heal at the same rate and a current healing wound may still exist with sequela of a healed burn or corrosion.

9) Use of an external cause code with burns and corruptions

An external cause code should be used with burns and corruptions to identify the source and intent of the burn, as well as the place where it occurred.

e. Adverse Effects, Poisoning , Underdosing and Toxic Effects

Codes in categories T36-T65 are combination codes that include the substances related to adverse effects, poisonings, toxic effects and underdosing, as well as the external cause. No additional external cause code is required for poisonings, toxic effects, adverse effects and underdosing codes.

A code from categories T36-T65 is sequenced first, followed by the code(s) that specify the nature of the adverse effect, poisoning, or toxic effect. **Note: This sequencing instruction does not apply to**

underdosing codes (fifth or sixth character “6”, for example T36.0x6-).

- 1) Do not code directly from the Table of Drugs**
Do not code directly from the Table of Drugs and Chemicals. Always refer back to the Tabular List.
- 2) Use as many codes as necessary to describe**
Use as many codes as necessary to describe completely all drugs, medicinal or biological substances.
- 3) If the same code would describe the causative agent**
If the same code would describe the causative agent for more than one adverse reaction, poisoning, toxic effect or underdosing, assign the code only once.
- 4) If two or more drugs, medicinal or biological substances**
If two or more drugs, medicinal or biological substances are reported, code each individually unless the combination code is listed in the Table of Drugs and Chemicals.
- 5) The occurrence of drug toxicity is classified in ICD-10-CM as follows:**
 - (a) Adverse Effect**
Assign the appropriate code for adverse effect (for example, T36.0x5-) when the drug was correctly prescribed and properly administered. Use additional code(s) for all manifestations of adverse effects. Examples of manifestations are tachycardia, delirium, gastrointestinal hemorrhaging, vomiting, hypokalemia, hepatitis, renal failure, or respiratory failure.
 - (b) Poisoning**
When coding a poisoning or reaction to the improper use of a medication (e.g., overdose, wrong substance given or taken in error, wrong route of administration), assign the appropriate code from categories T36-T50. Poisoning codes have an associated intent: accidental, intentional self-harm, assault and undetermined. Use additional code(s) for all manifestations of poisonings.

If there is also a diagnosis of **abuse or dependence on** the substance, the abuse or dependence is coded as an additional code.

Examples of poisoning include:

- (i) Error was made in drug prescription
Errors made in drug prescription or in the administration of the drug by provider, nurse, patient, or other person.
- (ii) Overdose of a drug intentionally taken
If an overdose of a drug was intentionally taken or administered and resulted in drug toxicity, it would be coded as a poisoning.
- (iii) Nonprescribed drug taken with correctly prescribed and properly administered drug
If a nonprescribed drug or medicinal agent was taken in combination with a correctly prescribed and properly administered drug, any drug toxicity or other reaction resulting from the interaction of the two drugs would be classified as a poisoning.
- (iv) Interaction of drug(s) and alcohol
When a reaction results from the interaction of a drug(s) and alcohol, this would be classified as poisoning.

See Section I.C.4. if poisoning is the result of insulin pump malfunctions.

(c) **Underdosing**

Underdosing refers to taking less of a medication than is prescribed by a **provider** or a manufacturer's instruction. For underdosing, assign the code from categories T36-T50 (**fifth or sixth character "6"**).

Codes for underdosing should never be assigned as principal or first-listed codes. If a patient has a relapse or exacerbation of the medical condition for which the drug is prescribed because of the reduction in dose, then the medical condition itself should be coded.

Noncompliance (Z91.12-, Z91.13-) or complication of care (Y63.61, Y63.8-Y63.9) codes are to be used with an underdosing code to indicate intent, if known.

(d) Toxic Effects

When a harmful substance is ingested or comes in contact with a person, this is classified as a toxic effect. The toxic effect codes are in categories T51-T65.

Toxic effect codes have an associated intent: accidental, intentional self-harm, assault and undetermined.

f. Adult and child abuse, neglect and other maltreatment

Sequence first the appropriate code from categories T74.- or T76.- for abuse, neglect and other maltreatment, followed by any accompanying mental health or injury code(s).

If the documentation in the medical record states abuse or neglect it is coded as confirmed. It is coded as suspected if it is documented as suspected.

For cases of confirmed abuse or neglect an external cause code from the assault section (X92-Y08) should be added to identify the cause of any physical injuries. A perpetrator code (Y07) should be added when the perpetrator of the abuse is known. For suspected cases of abuse or neglect, do not report external cause or perpetrator code.

If a suspected case of abuse, neglect or mistreatment is ruled out during an encounter code Z04.71, Suspected adult physical and sexual abuse, ruled out, or code Z04.72, Suspected child physical and sexual abuse, ruled out, should be used, not a code from T76.

g. Complications of care

1) Complications of care

(a) Documentation of complications of care

As with all procedural or postprocedural complications, code assignment is based on the provider's documentation of the relationship between the condition and the procedure.

2) **Pain due to medical devices**

Pain associated with devices, implants or grafts left in a surgical site (for example painful hip prosthesis) is assigned to the appropriate code(s) found in Chapter 19, Injury, poisoning, and certain other consequences of external causes. Specific codes for pain due to medical devices are found in the T code section of the ICD-10-CM. Use additional code(s) from category G89 to identify acute or chronic pain due to presence of the device, implant or graft (G89.18 or G89.28).

3) **Transplant complications**

(a) **Transplant complications other than kidney**

Codes under category T86, Complications of transplanted organs and tissues, are for use for both complications and rejection of transplanted organs. A transplant complication code is only assigned if the complication affects the function of the transplanted organ. Two codes are required to fully describe a transplant complication: the appropriate code from category T86 and a secondary code that identifies the complication.

Pre-existing conditions or conditions that develop after the transplant are not coded as complications unless they affect the function of the transplanted organs.

*See I.C.21.c.3 for transplant organ removal status
See I.C.2.r for malignant neoplasm associated with transplanted organ.*

(b) **Chronic kidney disease and kidney transplant complications**

Patients who have undergone kidney transplant may still have some form of chronic kidney disease (CKD) because the kidney transplant may not fully restore kidney function. Code T86.1- should be assigned for documented complications of a kidney transplant, such as transplant failure or rejection or other transplant complication. Code T86.1- should not be assigned for post kidney transplant patients who have chronic kidney (CKD) unless a transplant complication such as transplant failure or rejection is documented. If the

documentation is unclear as to whether the patient has a complication of the transplant, query the provider.

For patients with CKD following a kidney transplant, but who do not have a complication such as failure or rejection, *see section I.C.14. Chronic kidney disease and kidney transplant status.*

4) Complication codes that include the external cause

As with certain other T codes, some of the complications of care codes have the external cause included in the code. The code includes the nature of the complication as well as the type of procedure that caused the complication. No external cause code indicating the type of procedure is necessary for these codes.

5) Complications of care codes within the body system chapters

Intraoperative and postprocedural complication codes are found within the body system chapters with codes specific to the organs and structures of that body system. These codes should be sequenced first, followed by a code(s) for the specific complication, if applicable.

6) Ventilator associated pneumonia

(a) Documentation of Ventilator associated Pneumonia

As with all procedural or postprocedural complications, code assignment is based on the provider's documentation of the relationship between the condition and the procedure.

Code J95.851, Ventilator associated pneumonia, should be assigned only when the provider has documented ventilator associated pneumonia (VAP). An additional code to identify the organism (e.g., Pseudomonas aeruginosa, code B96.5) should also be assigned. Do not assign an additional code from categories J12-J18 to identify the type of pneumonia.

Code J95.851 should not be assigned for cases where the patient has pneumonia and is on a mechanical ventilator but the provider has not specifically stated that the pneumonia is ventilator-associated pneumonia.

If the documentation is unclear as to whether the patient has a pneumonia that is a complication attributable to the mechanical ventilator, query the provider.

- (b) **Patient admitted with pneumonia and develops VAP**
A patient may be admitted with one type of pneumonia (e.g., code J13, Pneumonia due to *Streptococcus pneumoniae*) and subsequently develop VAP. In this instance, the principal diagnosis would be the appropriate code from categories J12-J18 for the pneumonia diagnosed at the time of admission. Code J95.851, Ventilator associated pneumonia, would be assigned as an additional diagnosis when the provider has also documented the presence of ventilator associated pneumonia.

20. Chapter 20: External Causes of Morbidity (V01- Y99)

Introduction: These guidelines are provided for the reporting of external causes of morbidity codes in order that there will be standardization in the process. These codes are secondary codes for use in any health care setting.

External cause codes are intended to provide data for injury research and evaluation of injury prevention strategies. These codes capture how the injury or health condition happened (cause), the intent (unintentional or accidental; or intentional, such as suicide or assault), the place where the event occurred the activity of the patient at the time of the event, **and the person's status (e.g., civilian, military).**

a. General External Cause Coding Guidelines

1) Used with any code in the range of A00.0-T88.9, Z00-Z99

An external cause code may be used with any code in the range of A00.0-T88.9, Z00-Z99, classification that is a health condition due to an external cause. Though they are most applicable to injuries, they are also valid for use with such things as infections or diseases due to an external source, and other health conditions, such as a heart attack that occurs during strenuous physical activity.

2) External cause code used for length of treatment

Assign the external cause code, with the appropriate 7th character (initial encounter, subsequent encounter or sequela)

for each encounter for which the injury or condition is being treated.

3) Use the full range of external cause codes

Use the full range of external cause codes to completely describe the cause, the intent, the place of occurrence, **and if applicable**, the activity of the patient at the time of the event, **and the patient's status**, for all injuries, and other health conditions due to an external cause.

4) Assign as many external cause codes as necessary

Assign as many external cause codes as necessary to fully explain each cause. If only one external code can be recorded, assign the code most related to the principal diagnosis.

5) The selection of the appropriate external cause code

The selection of the appropriate external cause code is guided by the Index to External Causes, which is located after the Alphabetical Index to diseases and by Inclusion and Exclusion notes in the Tabular List.

6) External cause code can never be a principal diagnosis

An external cause code can never be a principal (first listed) diagnosis.

7) Combination external cause codes

Certain of the external cause codes are combination codes that identify sequential events that result in an injury, such as a fall which results in striking against an object. The injury may be due to either event or both. The combination external cause code used should correspond to the sequence of events regardless of which caused the most serious injury.

8) No external cause code needed in certain circumstances

No external cause code from Chapter 20 is needed if the external cause and intent are included in a code from another chapter (e.g. T360x1- Poisoning by penicillins, accidental (unintentional)).

b. Place of Occurrence Guideline

Codes from category Y92, Place of occurrence of the external cause, are secondary codes for use after other external cause codes to identify the location of the patient at the time of injury or other condition.

A place of occurrence code is used only once, at the initial encounter for treatment. No 7th characters are used for Y92. Only one code from Y92 should be recorded on a medical record. A place of occurrence code should be used in conjunction with an activity code, Y93.

Do not use place of occurrence code Y92.9 if the place is not stated or is not applicable.

c. Activity Code

Assign a code from category Y93, Activity code, to describe the activity of the patient at the time the injury or other health condition occurred.

An activity code is used only once, at the initial encounter for treatment. Only one code from Y93 should be recorded on a medical record. An activity code should be used in conjunction with a place of occurrence code, Y92.

If a patient is a student but is injured while performing an activity for income, use 7th character “2”, work related activity. A work related activity is any activity for which payment or income is received.

The activity codes are not applicable to poisonings, adverse effects, misadventures or late effects.

Do not assign Y93.9, Unspecified activity, if the activity is not stated.

d. Place of Occurrence, Activity, and Status Codes Used with other External Cause Code

When applicable, place of occurrence, activity, and external cause status codes are sequenced after the main external cause code(s). Regardless of the number of external cause codes assigned, there should be only one place of occurrence code, one activity code, and one external cause status code assigned to an encounter.

e. If the Reporting Format Limits the Number of External Cause Codes

If the reporting format limits the number of external cause codes that can be used in reporting clinical data, **report the code for the cause/intent** most related to the principal diagnosis. **If the format permits capture of additional external cause codes, the cause/intent, including medical misadventures, of the additional events should be reported rather than the codes for place, activity, or external status.**

f. Multiple External Cause Coding Guidelines

More than one external cause code is required to fully describe the external cause of an illness, injury or poisoning. The assignment of external cause codes should be sequenced in the following priority:

If two or more events cause separate injuries, an external cause code should be assigned for each cause. The first listed external cause code will be selected in the following order:

External codes for child and adult abuse take priority over all other external cause codes.

See Section I.C.19., Child and Adult abuse guidelines.

External cause codes for terrorism events take priority over all other external cause codes except child and adult abuse.

External cause codes for cataclysmic events take priority over all other external cause codes except child and adult abuse and terrorism.

External cause codes for transport accidents take priority over all other external cause codes except cataclysmic events, child and adult abuse and terrorism.

Activity and external cause status codes are assigned following all causal (intent) external cause codes.

The first-listed external cause code should correspond to the cause of the most serious diagnosis due to an assault, accident, or self-harm, following the order of hierarchy listed above.

g. Child and Adult Abuse Guideline

Adult and child abuse, neglect and maltreatment are classified as assault. Any of the assault codes may be used to indicate the external cause of any injury resulting from the confirmed abuse.

For confirmed cases of abuse, neglect and maltreatment, when the perpetrator is known, a code from Y07, Perpetrator of maltreatment and neglect, should accompany any other assault codes.

See Section I.C.19. Adult and child abuse, neglect and other maltreatment

h. Unknown or Undetermined Intent Guideline

If the intent (accident, self-harm, assault) of the cause of an injury or other condition is unknown or unspecified, code the intent as

accidental intent. All transport accident categories assume accidental intent.

1) Use of undetermined intent

External cause codes for events of undetermined intent are only for use if the documentation in the record specifies that the intent cannot be determined

i. Late Effects of External Cause Guidelines

1) Late effect external cause codes

Late effects are reported using the external cause code with the 7th character extension “S” for sequela. These codes should be used with any report of a late effect or sequela resulting from a previous injury.

2) Late effect external cause code with a related current injury

A late effect external cause code should never be used with a related current nature of injury code.

3) Use of late effect external cause codes for subsequent visits

Use a late effect external cause code for subsequent visits when a late effect of the initial injury is being treated. Do not use a late effect external cause code for subsequent visits for follow-up care (e.g., to assess healing, to receive rehabilitative therapy) of the injury or poisoning when no late effect of the injury has been documented.

j. Terrorism Guidelines

1) Cause of injury identified by the Federal Government (FBI) as terrorism

When the cause of an injury is identified by the Federal Government (FBI) as terrorism, the first-listed external cause code should be a code from category Y38, Terrorism. The definition of terrorism employed by the FBI is found at the inclusion note at the beginning of category Y38. Use additional code for place of occurrence (Y92.-). More than one Y38 code may be assigned if the injury is the result of more than one mechanism of terrorism.

2) Cause of an injury is suspected to be the result of terrorism

When the cause of an injury is suspected to be the result of terrorism a code from category Y38 should not be assigned. Suspected cases should be classified as assault.

3) Code Y38.9, Terrorism, secondary effects

Assign code Y38.9, Terrorism, secondary effects, for conditions occurring subsequent to the terrorist event. This code should not be assigned for conditions that are due to the initial terrorist act.

It is acceptable to assign code Y38.9 with another code from Y38 if there is an injury due to the initial terrorist event and an injury that is a subsequent result of the terrorist event.

k. External cause status

A code from category Y99, External cause status, should be assigned whenever any other external cause code is assigned for an encounter, including an Activity code, except for the events noted below. Assign a code from category Y99, External cause status, to indicate the work status of the person at the time the event occurred. The status code indicates whether the event occurred during military activity, whether a non-military person was at work, whether an individual including a student or volunteer was involved in a non-work activity at the time of the causal event.

A code from Y99, External cause status, should be assigned, when applicable, with other external cause codes, such as transport accidents and falls. The external cause status codes are not applicable to poisonings, adverse effects, misadventures or late effects.

Do not assign a code from category Y99 if no other external cause codes (cause, activity) are applicable for the encounter.

Do not assign code Y99.9, Unspecified external cause status, if the status is not stated.

21. Chapter 21: Factors influencing health status and contact with health services (Z00-Z99)

Note: The chapter specific guidelines provide additional information about the use of Z codes for specified encounters.

a. Use of Z codes in any healthcare setting

Z codes are for use in any healthcare setting. Z codes may be used as either a first listed (principal diagnosis code in the inpatient setting) or

secondary code, depending on the circumstances of the encounter. Certain **Z** codes may only be used as first listed or principal diagnosis.

b. Z Codes indicate a reason for an encounter

Z codes are not procedure codes. A corresponding procedure code must accompany a Z code to describe the procedure performed.

c. Categories of Z Codes

1) Contact/Exposure

Category Z20 indicates contact with, **and suspected** exposure to, communicable diseases. These codes are for patients who do not show any sign or symptom of a disease but **are suspected to** have been exposed to it by close personal contact with an infected individual or are in an area where a disease is epidemic.

Category Z77, indicates contact with and suspected exposures hazardous to health.

Contact/exposure codes may be used as a first listed code to explain an encounter for testing, or, more commonly, as a secondary code to identify a potential risk.

2) Inoculations and vaccinations

Code Z23 is for encounters for inoculations and vaccinations. It indicates that a patient is being seen to receive a prophylactic inoculation against a disease. Procedure codes are required to identify the actual administration of the injection and the type(s) of immunizations given. Code Z23 may be used as a secondary code if the inoculation is given as a routine part of preventive health care, such as a well-baby visit.

3) Status

Status codes indicate that a patient is either a carrier of a disease or has the sequelae or residual of a past disease or condition. This includes such things as the presence of prosthetic or mechanical devices resulting from past treatment. A status code is informative, because the status may affect the course of treatment and its outcome. A status code is distinct from a history code. The history code indicates that the patient no longer has the condition.

A status code should not be used with a diagnosis code from one of the body system chapters, if the diagnosis code includes the information provided by the status code. For example,

code Z94.1, Heart transplant status, should not be used with a code from subcategory T86.2, Complications of heart transplant. The status code does not provide additional information. The complication code indicates that the patient is a heart transplant patient.

For encounters for weaning from a mechanical ventilator, assign code J96.1, Chronic respiratory failure, followed by code Z99.11, Dependence on respirator [ventilator] status.

The status Z codes/categories are:

- Z14 Genetic carrier
Genetic carrier status indicates that a person carries a gene, associated with a particular disease, which may be passed to offspring who may develop that disease. The person does not have the disease and is not at risk of developing the disease.
- Z15 Genetic susceptibility to disease
Genetic susceptibility indicates that a person has a gene that increases the risk of that person developing the disease.

Codes from category Z15 should not be used as principal or first-listed codes. If the patient has the condition to which he/she is susceptible, and that condition is the reason for the encounter, the code for the current condition should be sequenced first. If the patient is being seen for follow-up after completed treatment for this condition, and the condition no longer exists, a follow-up code should be sequenced first, followed by the appropriate personal history and genetic susceptibility codes. If the purpose of the encounter is genetic counseling associated with procreative management, code Z31.5, Encounter for genetic counseling, should be assigned as the first-listed code, followed by a code from category Z15. Additional codes should be assigned for any applicable family or personal history.

- Z16 Infection with drug-resistant microorganisms
This code indicates that a patient has an infection that is resistant to drug treatment. Sequence the infection code first.
- Z17 Estrogen receptor status
- Z21 Asymptomatic HIV infection status
This code indicates that a patient has tested positive for HIV but has manifested no signs or symptoms of the disease.
- Z22 Carrier of infectious disease
Carrier status indicates that a person harbors the specific organisms of a disease without manifest symptoms and is capable of transmitting the infection.
- Z28.3 Underimmunization status**
- Z33.1 Pregnant state, incidental
This code is a secondary code only for use when the pregnancy is in no way complicating the reason for visit. Otherwise, a code from the obstetric chapter is required.
- Z66 Do not resuscitate
- Z67 Blood type
- Z68 Body mass index (BMI)
- Z74.01 Bed confinement status
- Z76.82 Awaiting organ transplant status
- Z78 Other specified health status**
- Z79 Long-term (current) drug therapy
Codes from this category indicate a patient's continuous use of a prescribed drug (including such things as aspirin therapy) for the long-term treatment of a condition or for prophylactic use. It is not for use for patients who have addictions to drugs. This subcategory is not for use of medications for detoxification or maintenance programs to prevent withdrawal symptoms in patients with drug dependence (e.g., methadone maintenance for opiate dependence). Assign the appropriate code for the drug dependence instead.

Assign a code from Z79 if the patient is receiving a medication for an extended period as a prophylactic measure (such as for the prevention of deep vein thrombosis) or as treatment of a chronic condition (such as arthritis) or a disease requiring a lengthy course of treatment (such as cancer). Do not assign a code from category Z79 for medication being administered for a brief period of time to treat an

- acute illness or injury (such as a course of antibiotics to treat acute bronchitis).
- Z88 Allergy status to drugs, medicaments and biological substances
Except: Z88.9, Allergy status to unspecified drugs, medicaments and biological substances status
- Z89 Acquired absence of limb
- Z90 Acquired absence of organs, not elsewhere classified
- Z91.0- Allergy status, other than to drugs and biological substances
- Z92.82 Status post administration of tPA (rtPA) in a different facility within the last 24 hours prior to admission to a current facility**

Assign code Z92.82, Status post administration of tPA (rtPA) in a different facility within the last 24 hours prior to admission to current facility, as a secondary diagnosis when a patient is received by transfer into a facility and documentation indicates they were administered tissue plasminogen activator (tPA) within the last 24 hours prior to admission to the current facility.

This guideline applies even if the patient is still receiving the tPA at the time they are received into the current facility.

The appropriate code for the condition for which the tPA was administered (such as cerebrovascular disease or myocardial infarction) should be assigned first.

Code Z92.82 is only applicable to the receiving facility record and not to the transferring facility record.

- Z93 Artificial opening status
- Z94 Transplanted organ and tissue status
- Z95 Presence of cardiac and vascular implants and grafts
- Z96 Presence of other functional implants
- Z97 Presence of other devices
- Z98 Other postprocedural states
- Assign code Z98.85, Transplanted organ removal status, to indicate that a transplanted organ has been previously removed. This code should not be assigned for the encounter in which the**

transplanted organ is removed. The complication necessitating removal of the transplant organ should be assigned for that encounter.

See section I.C19.g.3. for information on the coding of organ transplant complications.

Z99 Dependence on enabling machines and devices, not elsewhere classified
Note: Categories Z89-Z90 and Z93-Z99 are for use only if there are no complications or malfunctions of the organ or tissue replaced, the amputation site or the equipment on which the patient is dependent.

4) History (of)

There are two types of history Z codes, personal and family. Personal history codes explain a patient's past medical condition that no longer exists and is not receiving any treatment, but that has the potential for recurrence, and therefore may require continued monitoring.

Family history codes are for use when a patient has a family member(s) who has had a particular disease that causes the patient to be at higher risk of also contracting the disease.

Personal history codes may be used in conjunction with follow-up codes and family history codes may be used in conjunction with screening codes to explain the need for a test or procedure. History codes are also acceptable on any medical record regardless of the reason for visit. A history of an illness, even if no longer present, is important information that may alter the type of treatment ordered.

The history Z code categories are:

Z80	Family history of primary malignant neoplasm
Z81	Family history of mental and behavioral disorders
Z82	Family history of certain disabilities and chronic diseases (leading to disablement)
Z83	Family history of other specific disorders
Z84	Family history of other conditions
Z85	Personal history of malignant neoplasm
Z86	Personal history of certain other diseases
Z87	Personal history of other diseases and conditions
Z91.4-	Personal history of psychological trauma, not elsewhere classified
Z91.5	Personal history of self-harm

- Z91.8- Other specified personal risk factors, not elsewhere classified
- Z92 Personal history of medical treatment
 Except: Z92.0, Personal history of contraception
Except: Z92.82, Status post administration of tPA (rtPA) in a different facility within the last 24 hours prior to admission to a current facility

5) Screening

Screening is the testing for disease or disease precursors in seemingly well individuals so that early detection and treatment can be provided for those who test positive for the disease (e.g., screening mammogram).

The testing of a person to rule out or confirm a suspected diagnosis because the patient has some sign or symptom is a diagnostic examination, not a screening. In these cases, the sign or symptom is used to explain the reason for the test.

A screening code may be a first listed code if the reason for the visit is specifically the screening exam. It may also be used as an additional code if the screening is done during an office visit for other health problems. A screening code is not necessary if the screening is inherent to a routine examination, such as a pap smear done during a routine pelvic examination.

Should a condition be discovered during the screening then the code for the condition may be assigned as an additional diagnosis.

The Z code indicates that a screening exam is planned. A procedure code is required to confirm that the screening was performed.

The screening Z codes/categories:

- Z11 Encounter for screening for infectious and parasitic diseases
- Z12 Encounter for screening for malignant neoplasms
- Z13 Encounter for screening for other diseases and disorders
 Except: Z13.9, Encounter for screening, unspecified
- Z36 Encounter for antenatal screening for mother

6) **Observation**

There are two observation Z code categories. They are for use in very limited circumstances when a person is being observed for a suspected condition that is ruled out. The observation codes are not for use if an injury or illness or any signs or symptoms related to the suspected condition are present. In such cases the diagnosis/symptom code is used with the corresponding external cause code.

The observation codes are to be used as principal diagnosis only. Additional codes may be used in addition to the observation code but only if they are unrelated to the suspected condition being observed.

Codes from subcategory Z03.7 Encounter for suspected maternal and fetal conditions ruled out, may either be used as a first listed or as an additional code assignment depending on the case. They are for use in very limited circumstances on a maternal record when an encounter is for a suspected maternal or fetal condition that is ruled out during that encounter (for example, a maternal or fetal condition may be suspected due to an abnormal test result). These codes should not be used when the condition is confirmed. In those cases, the confirmed condition should be coded. In addition, these codes are not for use if an illness or any signs or symptoms related to the suspected condition or problem are present. In such cases the diagnosis/symptom code is used.

Additional codes may be used in addition to the code from subcategory Z03.7, but only if they are unrelated to the suspected condition being evaluated.

Codes from subcategory Z03.7 may not be used for encounters for antenatal screening of mother. See Section I.C.21.c.5, Screening.

For encounters for suspected fetal condition that are inconclusive following testing and evaluation, assign the appropriate code from category O35, O36, O40 or O41.

The observation Z code categories:

- Z03 Encounter for medical observation for suspected diseases and conditions ruled out
- Z04 Encounter for examination and observation for other reasons

Except: Z04.9, Encounter for examination and observation for unspecified reason

7) **Aftercare**

Aftercare visit codes cover situations when the initial treatment of a disease has been performed and the patient requires continued care during the healing or recovery phase, or for the long-term consequences of the disease. The aftercare Z code should not be used if treatment is directed at a current, acute disease. The diagnosis code is to be used in these cases.

Exceptions to this rule are codes Z51.0, Encounter for antineoplastic radiation therapy, and codes from subcategory Z51.1, Encounter for antineoplastic chemotherapy and immunotherapy. These codes are to be first listed, followed by the diagnosis code when a patient's encounter is solely to receive radiation therapy, chemotherapy, or immunotherapy for the treatment of a neoplasm. If the reason for the encounter is more than one type of antineoplastic therapy, code Z51.0 and a code from subcategory Z51.1 may be assigned together, in which case one of these codes would be reported as a secondary diagnosis.

The aftercare Z codes should also not be used for aftercare for injuries. For aftercare of an injury, assign the acute injury code with the 7th character "D" (subsequent encounter).

The aftercare codes are generally first listed to explain the specific reason for the encounter. An aftercare code may be used as an additional code when some type of aftercare is provided in addition to the reason for admission and no diagnosis code is applicable. An example of this would be the closure of a colostomy during an encounter for treatment of another condition.

Aftercare codes should be used in conjunction with other aftercare codes or diagnosis codes to provide better detail on the specifics of an aftercare encounter visit, unless otherwise directed by the classification. Should a patient receive multiple types of antineoplastic therapy during the same encounter, code Z51.0, Encounter for antineoplastic radiation therapy, and codes from subcategory Z51.1, Encounter for antineoplastic chemotherapy and immunotherapy, may be used together on a record. The sequencing of multiple aftercare codes **depends on the circumstances of the encounter.**

Certain aftercare Z code categories need a secondary diagnosis code to describe the resolving condition or sequelae. For others, the condition is **included** in the code title.

Additional Z code aftercare category terms include fitting and adjustment, and attention to artificial openings.

Status Z codes may be used with aftercare Z codes to indicate the nature of the aftercare. For example code Z95.1, Presence of aortocoronary bypass graft, may be used with code Z48.812, Encounter for surgical aftercare following surgery on the circulatory system, to indicate the surgery for which the aftercare is being performed. A status code should not be used when the aftercare code indicates the type of status, such as using Z43.0, Encounter for attention to tracheostomy, with Z93.0, Tracheostomy status.

The aftercare Z category/codes:

Z42	Encounter for plastic and reconstructive surgery following medical procedure or healed injury
Z43	Encounter for attention to artificial openings
Z44	Encounter for fitting and adjustment of external prosthetic device
Z45	Encounter for adjustment and management of implanted device
Z46	Encounter for fitting and adjustment of other devices
Z47	Orthopedic aftercare
Z48	Encounter for other postprocedural aftercare
Z49	Encounter for care involving renal dialysis
Z51	Encounter for other aftercare

8) **Follow-up**

The follow-up codes are used to explain continuing surveillance following completed treatment of a disease, condition, or injury. They imply that the condition has been fully treated and no longer exists. They should not be confused with aftercare codes, or injury codes with 7th character “D,” that explain ongoing care of a healing condition or its sequelae. Follow-up codes may be used in conjunction with history codes to provide the full picture of the healed condition and its treatment. The follow-up code is sequenced first, followed by the history code.

A follow-up code may be used to explain **multiple** visits. Should a condition be found to have recurred on the follow-up

visit, then the code for the condition should be assigned as an additional diagnosis.

The follow-up Z code categories:

- Z08 Encounter for follow-up examination after completed treatment for malignant neoplasm
- Z09 Encounter for follow-up examination after completed treatment for conditions other than malignant neoplasm
- Z39 Encounter for maternal postpartum care and examination

9) Donor

Codes in category Z52, Donors of organs and tissues, are used for living individuals who are donating blood or other body tissue. These codes are only for individuals donating for others, not for self-donations. They are not used to identify cadaveric donations.

10) Counseling

Counseling Z codes are used when a patient or family member receives assistance in the aftermath of an illness or injury, or when support is required in coping with family or social problems. They are not used in conjunction with a diagnosis code when the counseling component of care is considered integral to standard treatment.

The counseling Z codes/categories:

- Z30.0- Encounter for general counseling and advice on contraception
- Z31.5 Encounter for genetic counseling
- Z31.6- Encounter for general counseling and advice on procreation
- Z32.2 Encounter for childbirth instruction
- Z32.3 Encounter for childcare instruction
- Z69 Encounter for mental health services for victim and perpetrator of abuse
- Z70 Counseling related to sexual attitude, behavior and orientation
- Z71 Persons encountering health services for other counseling and medical advice, not elsewhere classified
- Z76.81 Expectant mother prebirth pediatrician visit

11) Encounters for Obstetrical and Reproductive Services

See Section I.C.15. Pregnancy, Childbirth, and the Puerperium, for further instruction on the use of these codes.

Z codes for pregnancy are for use in those circumstances when none of the problems or complications included in the codes from the Obstetrics chapter exist (a routine prenatal visit or postpartum care). Codes in category Z34, Encounter for supervision of normal pregnancy, are always first listed and are not to be used with any other code from the OB chapter.

The outcome of delivery, category Z37, should be included on all maternal delivery records. It is always a secondary code. Codes in category Z37 should not be used on the newborn record.

Z codes for family planning (contraceptive) or procreative management and counseling should be included on an obstetric record either during the pregnancy or the postpartum stage, if applicable.

Z codes/categories for obstetrical and reproductive services:

Z30	Encounter for contraceptive management
Z31	Encounter for procreative management
Z32.2	Encounter for childbirth instruction
Z32.3	Encounter for childcare instruction
Z33	Pregnant state
Z34	Encounter for supervision of normal pregnancy
Z36	Encounter for antenatal screening of mother
Z37	Outcome of delivery
Z39	Encounter for maternal postpartum care and examination
Z76.81	Expectant mother prebirth pediatrician visit

12) Newborns and Infants

See Section I.C.16. Newborn (Perinatal) Guidelines, for further instruction on the use of these codes.

Newborn Z codes/categories:

Z76.1	Encounter for health supervision and care of foundling
Z00.1-	Encounter for routine child health examination
Z38	Liveborn infants according to place of birth and type of delivery

13) Routine and administrative examinations

The Z codes allow for the description of encounters for routine examinations, such as, a general check-up, or, examinations for administrative purposes, such as, a pre-employment physical. The codes are not to be used if the examination is for diagnosis of a suspected condition or for treatment purposes. In such cases the diagnosis code is used. During a routine exam, should a diagnosis or condition be discovered, it should be coded as an additional code. Pre-existing and chronic conditions and history codes may also be included as additional codes as long as the examination is for administrative purposes and not focused on any particular condition.

Some of the codes for routine health examinations distinguish between “with” and “without” abnormal findings. Code assignment depends on the information that is known at the time the encounter is being coded. For example, if no abnormal findings were found during the examination, but the encounter is being coded before test results are back, it is acceptable to assign the code for “without abnormal findings.” When assigning a code for “with abnormal findings,” additional code(s) should be assigned to identify the specific abnormal finding(s).

Pre-operative examination and pre-procedural laboratory examination Z codes are for use only in those situations when a patient is being cleared for **a procedure or** surgery and no treatment is given.

The Z codes/categories for routine and administrative examinations:

Z00	Encounter for general examination without complaint, suspected or reported diagnosis
Z01	Encounter for other special examination without complaint, suspected or reported diagnosis
Z02	Encounter for administrative examination Except: Z02.9, Encounter for administrative examinations, unspecified
Z32.0-	Encounter for pregnancy test

14) Miscellaneous Z codes

The miscellaneous Z codes capture a number of other health care encounters that do not fall into one of the other categories. Certain of these codes identify the reason for the encounter; others are for use as additional codes that provide useful information on circumstances that may affect a patient’s care and treatment.

Prophylactic Organ Removal

For encounters specifically for prophylactic removal of an organ (such as prophylactic removal of breasts due to a genetic susceptibility to cancer or a family history of cancer), the principal or first listed code should be a code from category Z40, Encounter for prophylactic surgery, followed by the appropriate codes to identify the associated risk factor (such as genetic susceptibility or family history).

If the patient has a malignancy of one site and is having prophylactic removal at another site to prevent either a new primary malignancy or metastatic disease, a code for the malignancy should also be assigned in addition to a code from subcategory Z40.0, Encounter for prophylactic surgery for risk factors related to malignant neoplasms. A Z40.0 code should not be assigned if the patient is having organ removal for treatment of a malignancy, such as the removal of the testes for the treatment of prostate cancer.

Miscellaneous Z codes/categories:

Z28	Immunization not carried out Except: Z28.3, Underimmunization status
Z40	Encounter for prophylactic surgery
Z41	Encounter for procedures for purposes other than remedying health state Except: Z41.9, Encounter for procedure for purposes other than remedying health state, unspecified
Z53	Persons encountering health services for specific procedures and treatment, not carried out
Z55	Problems related to education and literacy
Z56	Problems related to employment and unemployment
Z57	Occupational exposure to risk factors
Z58	Problems related to physical environment
Z59	Problems related to housing and economic circumstances
Z60	Problems related to social environment
Z62	Problems related to upbringing
Z63	Other problems related to primary support group, including family circumstances
Z64	Problems related to certain psychosocial circumstances
Z65	Problems related to other psychosocial circumstances
Z72	Problems related to lifestyle
Z73	Problems related to life management difficulty

Z74	Problems related to care provider dependency Except: Z74.01, Bed confinement status
Z75	Problems related to medical facilities and other health care
Z76.0	Encounter for issue of repeat prescription
Z76.3	Healthy person accompanying sick person
Z76.4	Other boarder to healthcare facility
Z76.5	Malingerer [conscious simulation]
Z76.89	Persons encountering health services in other specified circumstances
Z91.1-	Patient's noncompliance with medical treatment and regimen
Z91.89	Other specified personal risk factors, not elsewhere classified

15) **Nonspecific Z codes**

Certain Z codes are so non-specific, or potentially redundant with other codes in the classification, that there can be little justification for their use in the inpatient setting. Their use in the outpatient setting should be limited to those instances when there is no further documentation to permit more precise coding. Otherwise, any sign or symptom or any other reason for visit that is captured in another code should be used.

Nonspecific Z codes/categories:

Z02.9	Encounter for administrative examinations, unspecified
Z04.9	Encounter for examination and observation for unspecified reason
Z13.9	Encounter for screening, unspecified
Z41.9	Encounter for procedure for purposes other than remedying health state, unspecified
Z52.9	Donor of unspecified organ or tissue
Z88.9	Allergy status to unspecified drugs, medicaments and biological substances status
Z92.0	Personal history of contraception

16) **Z Codes That May Only be Principal/First-Listed Diagnosis**

The following Z codes/categories may only be reported as the principal/first-listed diagnosis, except when there are multiple encounters on the same day and the medical records for the encounters are combined:

Z00	Encounter for general examination without complaint, suspected or reported diagnosis
-----	--

Z01	Encounter for other special examination without complaint, suspected or reported diagnosis
Z02	Encounter for administrative examination
Z03	Encounter for medical observation for suspected diseases and conditions ruled out
Z33.2	Encounter for elective termination of pregnancy
Z31.81	Encounter for male factor infertility in female patient
Z31.82	Encounter for Rh incompatibility status
Z31.83	Encounter for assisted reproductive fertility procedure cycle
Z31.84	Encounter for fertility preservation procedure
Z34	Encounter for supervision of normal pregnancy
Z39	Encounter for maternal postpartum care and examination
Z38	Liveborn infants according to place of birth and type of delivery
Z42	Encounter for plastic and reconstructive surgery following medical procedure or healed injury
Z51.0	Encounter for antineoplastic radiation therapy
Z51.1-	Encounter for antineoplastic chemotherapy and immunotherapy
Z52	Donors of organs and tissues Except: Z52.9, Donor of unspecified organ or tissue
Z76.1	Encounter for health supervision and care of foundling
Z76.2	Encounter for health supervision and care of other healthy infant and child
Z99.12	Encounter for respirator [ventilator] dependence during power failure

Section II. Selection of Principal Diagnosis

The circumstances of inpatient admission always govern the selection of principal diagnosis. The principal diagnosis is defined in the Uniform Hospital Discharge Data Set (UHDDS) as “that condition established after study to be chiefly responsible for occasioning the admission of the patient to the hospital for care.”

The UHDDS definitions are used by hospitals to report inpatient data elements in a standardized manner. These data elements and their definitions can be found in the July 31, 1985, Federal Register (Vol. 50, No, 147), pp. 31038-40.

Since that time the application of the UHDDS definitions has been expanded to include all non-outpatient settings (acute care, short term, long term care and psychiatric hospitals; home health agencies; rehab facilities; nursing homes, etc).

In determining principal diagnosis the coding conventions in the ICD-10-CM, Volumes I and II take precedence over these official coding guidelines.
(See Section I.A., *Conventions for the ICD-10-CM*)

The importance of consistent, complete documentation in the medical record cannot be overemphasized. Without such documentation the application of all coding guidelines is a difficult, if not impossible, task.

A. Codes for symptoms, signs, and ill-defined conditions

Codes for symptoms, signs, and ill-defined conditions from Chapter 18 are not to be used as principal diagnosis when a related definitive diagnosis has been established.

B. Two or more interrelated conditions, each potentially meeting the definition for principal diagnosis.

When there are two or more interrelated conditions (such as diseases in the same ICD-10-CM chapter or manifestations characteristically associated with a certain disease) potentially meeting the definition of principal diagnosis, either condition may be sequenced first, unless the circumstances of the admission, the therapy provided, the Tabular List, or the Alphabetic Index indicate otherwise.

C. Two or more diagnoses that equally meet the definition for principal diagnosis

In the unusual instance when two or more diagnoses equally meet the criteria for principal diagnosis as determined by the circumstances of admission, diagnostic workup and/or therapy provided, and the Alphabetic Index, Tabular List, or another coding guidelines does not provide sequencing direction, any one of the diagnoses may be sequenced first.

D. Two or more comparative or contrasting conditions.

In those rare instances when two or more contrasting or comparative diagnoses are documented as “either/or” (or similar terminology), they are coded as if the diagnoses were confirmed and the diagnoses are sequenced according to the circumstances of the admission. If no further determination can be made as to which diagnosis should be principal, either diagnosis may be sequenced first.

E. A symptom(s) followed by contrasting/comparative diagnoses

When a symptom(s) is followed by contrasting/comparative diagnoses, the symptom code is sequenced first. All the contrasting/comparative diagnoses should be coded as additional diagnoses.

F. Original treatment plan not carried out

Sequence as the principal diagnosis the condition, which after study occasioned the admission to the hospital, even though treatment may not have been carried out due to unforeseen circumstances.

G. Complications of surgery and other medical care

When the admission is for treatment of a complication resulting from surgery or other medical care, the complication code is sequenced as the principal diagnosis. If the complication is classified to the T80-T88 series and the code lacks the necessary specificity in describing the complication, an additional code for the specific complication should be assigned.

H. Uncertain Diagnosis

If the diagnosis documented at the time of discharge is qualified as “probable”, “suspected”, “likely”, “questionable”, “possible”, or “still to be ruled out”, or other similar terms indicating uncertainty, code the condition as if it existed or was established. The bases for these guidelines are the diagnostic workup, arrangements for further workup or observation, and initial therapeutic approach that correspond most closely with the established diagnosis.

Note: This guideline is applicable only to inpatient admissions to short-term, acute, long-term care and psychiatric hospitals.

I. Admission from Observation Unit

1. Admission Following Medical Observation

When a patient is admitted to an observation unit for a medical condition, which either worsens or does not improve, and is subsequently admitted as an inpatient of the same hospital for this same medical condition, the principal diagnosis would be the medical condition which led to the hospital admission.

2. Admission Following Post-Operative Observation

When a patient is admitted to an observation unit to monitor a condition (or complication) that develops following outpatient surgery, and then is subsequently admitted as an inpatient of the same hospital, hospitals should apply the Uniform Hospital Discharge Data Set (UHDDS) definition of principal diagnosis as "that condition established after study to be chiefly responsible for occasioning the admission of the patient to the hospital for care."

J. Admission from Outpatient Surgery

When a patient receives surgery in the hospital's outpatient surgery department and is subsequently admitted for continuing inpatient care at the same hospital, the following guidelines should be followed in selecting the principal diagnosis for the inpatient admission:

- If the reason for the inpatient admission is a complication, assign the complication as the principal diagnosis.

- If no complication, or other condition, is documented as the reason for the inpatient admission, assign the reason for the outpatient surgery as the principal diagnosis.
- If the reason for the inpatient admission is another condition unrelated to the surgery, assign the unrelated condition as the principal diagnosis.

Section III. Reporting Additional Diagnoses

GENERAL RULES FOR OTHER (ADDITIONAL) DIAGNOSES

For reporting purposes the definition for “other diagnoses” is interpreted as additional conditions that affect patient care in terms of requiring:

clinical evaluation; or
 therapeutic treatment; or
 diagnostic procedures; or
 extended length of hospital stay; or
 increased nursing care and/or monitoring.

The UHDDS item #11-b defines Other Diagnoses as “all conditions that coexist at the time of admission, that develop subsequently, or that affect the treatment received and/or the length of stay. Diagnoses that relate to an earlier episode which have no bearing on the current hospital stay are to be excluded.” UHDDS definitions apply to inpatients in acute care, short-term, long term care and psychiatric hospital setting. The UHDDS definitions are used by acute care short-term hospitals to report inpatient data elements in a standardized manner. These data elements and their definitions can be found in the July 31, 1985, Federal Register (Vol. 50, No, 147), pp. 31038-40.

Since that time the application of the UHDDS definitions has been expanded to include all non-outpatient settings (acute care, short term, long term care and psychiatric hospitals; home health agencies; rehab facilities; nursing homes, etc).

The following guidelines are to be applied in designating “other diagnoses” when neither the Alphabetic Index nor the Tabular List in ICD-10-CM provide direction. The listing of the diagnoses in the patient record is the responsibility of the attending provider.

A. Previous conditions

If the provider has included a diagnosis in the final diagnostic statement, such as the discharge summary or the face sheet, it should ordinarily be coded. Some providers include in the diagnostic statement resolved conditions or diagnoses and status-post procedures from previous admission that have no bearing on the current stay. Such conditions are not to be reported and are coded only if required by hospital policy.

However, history codes (categories Z80-Z87) may be used as secondary codes if the historical condition or family history has an impact on current care or influences treatment.

B. Abnormal findings

Abnormal findings (laboratory, x-ray, pathologic, and other diagnostic results) are not coded and reported unless the provider indicates their clinical significance. If the findings are outside the normal range and the attending provider has ordered other tests to evaluate the condition or prescribed treatment, it is appropriate to ask the provider whether the abnormal finding should be added.

Please note: This differs from the coding practices in the outpatient setting for coding encounters for diagnostic tests that have been interpreted by a provider.

C. Uncertain Diagnosis

If the diagnosis documented at the time of discharge is qualified as “probable”, “suspected”, “likely”, “questionable”, “possible”, or “still to be ruled out” or other similar terms indicating uncertainty, code the condition as if it existed or was established. The bases for these guidelines are the diagnostic workup, arrangements for further workup or observation, and initial therapeutic approach that correspond most closely with the established diagnosis.

Note: This guideline is applicable only to inpatient admissions to short-term, acute, long-term care and psychiatric hospitals.

Section IV. Diagnostic Coding and Reporting Guidelines for Outpatient Services

These coding guidelines for outpatient diagnoses have been approved for use by hospitals/providers in coding and reporting hospital-based outpatient services and provider-based office visits.

Information about the use of certain abbreviations, punctuation, symbols, and other conventions used in the ICD-10-CM Tabular List (code numbers and titles), can be found in Section IA of these guidelines, under “Conventions Used in the Tabular List.” Information about the correct sequence to use in finding a code is also described in Section I.

The terms encounter and visit are often used interchangeably in describing outpatient service contacts and, therefore, appear together in these guidelines without distinguishing one from the other.

Though the conventions and general guidelines apply to all settings, coding guidelines for outpatient and provider reporting of diagnoses will vary in a number of instances from those for inpatient diagnoses, recognizing that:

The Uniform Hospital Discharge Data Set (UHDDS) definition of principal diagnosis applies only to inpatients in acute, short-term, long-term care and psychiatric hospitals.

Coding guidelines for inconclusive diagnoses (probable, suspected, rule out, etc.) were developed for inpatient reporting and do not apply to outpatients.

A. Selection of first-listed condition

In the outpatient setting, the term first-listed diagnosis is used in lieu of principal diagnosis.

In determining the first-listed diagnosis the coding conventions of ICD-10-CM, as well as the general and disease specific guidelines take precedence over the outpatient guidelines.

Diagnoses often are not established at the time of the initial encounter/visit. It may take two or more visits before the diagnosis is confirmed.

The most critical rule involves beginning the search for the correct code assignment through the Alphabetic Index. Never begin searching initially in the Tabular List as this will lead to coding errors.

1. Outpatient Surgery

When a patient presents for outpatient surgery (same day surgery), code the reason for the surgery as the first-listed diagnosis (reason for the encounter), even if the surgery is not performed due to a contraindication.

2. Observation Stay

When a patient is admitted for observation for a medical condition, assign a code for the medical condition as the first-listed diagnosis.

When a patient presents for outpatient surgery and develops complications requiring admission to observation, code the reason for the surgery as the first reported diagnosis (reason for the encounter), followed by codes for the complications as secondary diagnoses.

B. Codes from A00.0 through T88.9, Z00-Z99

The appropriate code(s) from A00.0 through T88.9, Z00-Z99 must be used to identify diagnoses, symptoms, conditions, problems, complaints, or other reason(s) for the encounter/visit.

C. Accurate reporting of ICD-10-CM diagnosis codes

For accurate reporting of ICD-10-CM diagnosis codes, the documentation should describe the patient's condition, using terminology which includes specific diagnoses as well as symptoms, problems, or reasons for the encounter. There are ICD-10-CM codes to describe all of these.

D. Codes that describe symptoms and signs

Codes that describe symptoms and signs, as opposed to diagnoses, are acceptable for reporting purposes when a diagnosis has not been established (confirmed) by the provider. Chapter 18 of ICD-10-CM, Symptoms, Signs, and Abnormal Clinical and Laboratory Findings Not Elsewhere Classified (codes R00-R99) contain many, but not all codes for symptoms.

E. Encounters for circumstances other than a disease or injury

ICD-10-CM provides codes to deal with encounters for circumstances other than a disease or injury. The Factors Influencing Health Status and Contact with Health Services codes (Z00-99) is provided to deal with occasions when circumstances other than a disease or injury are recorded as diagnosis or problems.

See Section I.C.21. Factors influencing health status and contact with health services.

F. Level of Detail in Coding

1. ICD-10-CM codes with 3, 4, or 5 digits

ICD-10-CM is composed of codes with either 3, 4, 5, 6 or 7 digits. Codes with three digits are included in ICD-10-CM as the heading of a category of codes that may be further subdivided by the use of fourth fifth digits, sixth or seventh digits which provide greater specificity.

2. Use of full number of digits required for a code

A three-digit code is to be used only if it is not further subdivided. A code is invalid if it has not been coded to the full number of characters required for that code, including the 7th character extension, if applicable.

G. ICD-10-CM code for the diagnosis, condition, problem, or other reason for encounter/visit

List first the ICD-10-CM code for the diagnosis, condition, problem, or other reason for encounter/visit shown in the medical record to be chiefly responsible for the services provided. List additional codes that describe any coexisting conditions. In some cases the first-listed diagnosis may be a symptom when a diagnosis has not been established (confirmed) by the physician.

H. Uncertain diagnosis

Do not code diagnoses documented as “probable”, “suspected,” “questionable,” “rule out,” or “working diagnosis” or other similar terms indicating uncertainty. Rather, code the condition(s) to the highest degree of certainty for that encounter/visit, such as symptoms, signs, abnormal test results, or other reason for the visit.

Please note: This differs from the coding practices used by short-term, acute care, long-term care and psychiatric hospitals.

I. Chronic diseases

Chronic diseases treated on an ongoing basis may be coded and reported as many times as the patient receives treatment and care for the condition(s)

J. Code all documented conditions that coexist

Code all documented conditions that coexist at the time of the encounter/visit, and require or affect patient care treatment or management. Do not code conditions that were previously treated and no longer exist. However, history codes (categories Z80-Z87) may be used as secondary codes if the historical condition or family history has an impact on current care or influences treatment.

K. Patients receiving diagnostic services only

For patients receiving diagnostic services only during an encounter/visit, sequence first the diagnosis, condition, problem, or other reason for encounter/visit shown in the medical record to be chiefly responsible for the outpatient services provided during the encounter/visit. Codes for other diagnoses (e.g., chronic conditions) may be sequenced as additional diagnoses.

For encounters for routine laboratory/radiology testing in the absence of any signs, symptoms, or associated diagnosis, assign Z01.89, Encounter for other specified special examinations. If routine testing is performed during the same encounter as a test to evaluate a sign, symptom, or diagnosis, it is appropriate to assign both the V code and the code describing the reason for the non-routine test.

For outpatient encounters for diagnostic tests that have been interpreted by a physician, and the final report is available at the time of coding, code any confirmed or definitive diagnosis(es) documented in the interpretation. Do not code related signs and symptoms as additional diagnoses.

Please note: This differs from the coding practice in the hospital inpatient setting regarding abnormal findings on test results.

L. Patients receiving therapeutic services only

For patients receiving therapeutic services only during an encounter/visit, sequence first the diagnosis, condition, problem, or other reason for encounter/visit shown in the medical record to be chiefly responsible for the outpatient services provided during the encounter/visit. Codes for other diagnoses (e.g., chronic conditions) may be sequenced as additional diagnoses.

The only exception to this rule is that when the primary reason for the admission/encounter is chemotherapy or radiation therapy, the appropriate Z code for the service is listed first, and the diagnosis or problem for which the service is being performed listed second.

M. Patients receiving preoperative evaluations only

For patients receiving preoperative evaluations only, sequence first a code from subcategory Z01.81, Encounter for pre-procedural examinations, to describe the pre-op consultations. Assign a code for the condition to describe the reason for the surgery as an additional diagnosis. Code also any findings related to the pre-op evaluation.

N. Ambulatory surgery

For ambulatory surgery, code the diagnosis for which the surgery was performed. If the postoperative diagnosis is known to be different from the preoperative diagnosis at the time the diagnosis is confirmed, select the postoperative diagnosis for coding, since it is the most definitive.

O. Routine outpatient prenatal visits

See Section I.C.15. Routine outpatient prenatal visits.

P. Encounters for general medical examinations with abnormal findings

The subcategories for encounters for general medical examinations, Z00.0-, provide codes for with and without abnormal findings. Should a general medical examination result in an abnormal finding, the code for general medical examination with abnormal finding should be assigned as the first listed diagnosis. A secondary code for the abnormal finding should also be coded.

Q. Encounters for routine health screenings

See Section I.C.21. Factors influencing health status and contact with health services, Screening