

Potential Internal Medicine Preferred Specialty Measure Set

PQRS# CMS# GPRO#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
1 CMS122v2	0059	Claims, Registry, EHR, GPRO Web Interface, Measures Group (DM)	Effective Clinical Care	Diabetes: Hemoglobin A1c (HbA1c) Poor Control (>9%): Percentage of patients 18-75 years of age with diabetes who had hemoglobin A1c > 9.0% during the measurement period
5 CMS135v2	0081	Registry, EHR, Measures Group (HF)	Effective Clinical Care	Heart Failure: Angiotensin-Converting Enzyme (ACE) Inhibitor or Angiotensin Receptor Blocker (ARB) Therapy for Left Ventricular Systolic Dysfunction (LVSD): Percentage of patients aged 18 years and older with a diagnosis of heart failure (HF) with a current or prior left ventricular ejection fraction (LVEF) < 40% who were prescribed ACE inhibitor or ARB therapy either within a 12 month period when seen in the outpatient setting or at <u>each</u> hospital discharge
9 CMS128v2	0105	Registry, EHR	Effective Clinical Care	Anti-depressant Medication Management: Percentage of patients 18 years of age and older who were diagnosed with major depression and treated with antidepressant medication, and who remained on antidepressant medication treatment. Two rates are reported. • Effective Acute Phase Treatment: Percentage of patients who remained on an antidepressant medication for at least 84 days (12 weeks) • Effective Continuation Phase Treatment: Percentage of patients who remained on an antidepressant medication for at least 180 days (6 months)
40	0048	Registry	Effective Clinical Care	Osteoporosis: Management Following Fracture of Hip, Spine or Distal Radius for Men and Women Aged 50 Years and Older: Percentage of patients <u>aged 50 years and older</u> with fracture of the hip, spine, or distal radius who had a central dual-energy X-ray absorptiometry (DXA) measurement ordered or performed or pharmacologic therapy prescribed
41	0049	Claims, Registry	Effective Clinical Care	Osteoporosis: Pharmacologic Therapy for Men and Women Aged 50 Years and Older: Percentage of patients <u>aged 50 years and older</u> with a diagnosis of osteoporosis who were prescribed pharmacologic therapy within 12 months
50	0100	Registry	Person and Caregiver-Centered Experience and Outcomes	Urinary Incontinence: Plan of Care for Urinary Incontinence in Women Aged 65 Years and Older: Percentage of female patients aged 65 years and older with a diagnosis of urinary incontinence with a documented plan of care for urinary incontinence at least once within 12 months

Potential Internal Medicine Preferred Specialty Measure Set

PQRS# CMS# GPRO#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
109	0050	Registry	Person and Caregiver-Centered Experience and Outcomes	Osteoarthritis (OA): Function and Pain Assessment: Percentage of patient visits for patients aged 21 years and older with a diagnosis of osteoarthritis (OA) with assessment for function and pain
112 CMS125v2	0031	Registry, EHR, GPRO Web Interface, Measures Group (Prev Care)	Effective Clinical Care	Breast Cancer Screening: Percentage of women 50 through 74 years of age who had a mammogram to screen for breast cancer within 27 months
113 CMS130v2	0034	Registry, EHR, GPRO Web Interface, Measures Group (Prev Care)	Effective Clinical Care	Colorectal Cancer Screening: Percentage of patients 50 through 75 years of age who had appropriate screening for colorectal cancer
116	0058	Registry	Efficiency and Cost Reduction	Avoidance of Antibiotic Treatment in Adults With Acute Bronchitis: Percentage of adults 18 through 64 years of age with a diagnosis of acute bronchitis who were not prescribed or dispensed an antibiotic prescription on or 3 days after the episode
117 CMS131v2	0055	Registry, GPRO Web Interface, EHR, Measures Group (DM)	Effective Clinical Care	Diabetes: Eye Exam: Percentage of patients 18 through 75 years of age with a diagnosis of diabetes (type 1 and type 2) who had a retinal or dilated eye exam in the measurement period or a negative retinal or dilated eye exam (negative for retinopathy) in the year prior to the measurement period
134 CMS2v3	0418	Claims, Registry, EHR, GPRO Web Interface	Community/Population Health	Preventive Care and Screening: Screening for Clinical Depression and Follow-Up Plan: Percentage of patients aged 12 years and older screened for clinical depression on the date of the encounter using an age appropriate standardized depression screening tool AND if positive, a follow-up plan is documented on the date of the positive screen.
154	0101	Claims, Registry	Patient Safety	Falls: Risk Assessment: Percentage of patients aged 65 years and older with a history of falls who had a risk assessment for falls completed within 12 months
155	0101	Claims, Registry	Communication and Care Coordination	Falls: Plan of Care: Percentage of patients aged 65 years and older with a history of falls who had a plan of care for falls documented within 12 months
163 CMS123v2	0056	Registry, GPRO Web Interface, EHR, Measures Group (DM)	Effective Clinical Care	Diabetes: Foot Exam: Percentage of patients aged 18-75 years of age with diabetes who had a foot exam during the measurement period

Potential Internal Medicine Preferred Specialty Measure Set

PQRS# CMS# GPRO#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
181	N/A	Claims, Registry	Patient Safety	Elder Maltreatment Screen and Follow-Up Plan: Percentage of patients aged 65 years and older with a documented elder maltreatment screen using an Elder Maltreatment Screening Tool on the date of encounter AND a documented follow-up plan on the date of the positive screen
226 CMS138v2	0028	Claims, Registry, EHR, GPRO Web Interface, Measures Groups (CAD, COPD, HF, IBD, Prev Care, Oncology)	Community/Population Health	Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention: Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 24 months AND who received cessation counseling intervention if identified as a tobacco user.
236 CMS165v2	0018	Claims, Registry, EHR, GPRO Web Interface,	Effective Clinical Care	Controlling High Blood Pressure: Percentage of patients 18-85 years of age who had a diagnosis of hypertension and whose blood pressure was adequately controlled (<140/90 mmHg) during the measurement period.
242	N/A	Registry, GPRO Web Interface, Measures Group (CAD)	Effective Clinical Care	Coronary Artery Disease (CAD): Symptom Management: Percentage of patients aged 18 years and older with a diagnosis of coronary artery disease seen within a 12 month period with results of an evaluation of level of activity and an assessment of whether anginal symptoms are present or absent with appropriate management of anginal symptoms within a 12 month period
326	1525	Registry	Patient Safety	Atrial Fibrillation and Atrial Flutter: Chronic Anticoagulation Therapy: Percentage of patients aged 18 years and older with a diagnosis of nonvalvular atrial fibrillation (AF) or atrial flutter whose assessment of the specified thromboembolic risk factors indicate one or more high-risk factors or more than one moderate risk factor, as determined by CHADS2 risk stratification, who are prescribed warfarin OR another oral anticoagulant drug that is FDA approved for the prevention of thromboembolism
331	N/A	Registry	Effective Clinical Care	Adult Sinusitis: Antibiotic Prescribed for Acute Sinusitis (Appropriate Use): Percentage of patients, aged 18 years and older, with a diagnosis of acute sinusitis who were prescribed an antibiotic within 7 days of diagnosis or within 10 days after onset of symptoms

Potential Internal Medicine Preferred Specialty Measure Set

PQRS# CMS# GPRO#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
332	N/A	Registry	Effective Clinical Care	Adult Sinusitis: Appropriate Choice of Antibiotic: Amoxicillin Prescribed for Patients with Acute Bacterial Sinusitis: Percentage of patients aged 18 years and older with a diagnosis of acute bacterial sinusitis that were prescribed amoxicillin, with or without clavulanate, as a first line antibiotic at the time of diagnosis
333	N/A	Registry	Efficiency and Cost Reduction	Adult Sinusitis: Computerized Tomography (CT) for Acute Sinusitis (Overuse): Percentage of patients, aged 18 years and older, with a diagnosis of acute sinusitis who had a computerized tomography (CT) scan of the paranasal sinuses ordered at the time of diagnosis or received within 28 days after date of diagnosis
334	N/A	Registry	Efficiency and Cost Reduction	Adult Sinusitis: More than One Computerized Tomography (CT) Scan Within 90 Days for Chronic Sinusitis (Overuse): Percentage of patients aged 18 years and older with a diagnosis of chronic sinusitis who had more than one CT scan of the paranasal sinuses ordered or received within 90 days after date of diagnosis
2015 PQRS Proposed New Measure	N/A	Registry	Effective Clinical Care	Annual Hepatitis C Virus (HCV) Screening for Patients who are Active Injection Drug Users: Percentage of patients regardless of age who are active injection drug users who received a hepatitis C virus (HCV) antibody test or HCV ribonucleic acid (RNA) test within the 12 month reporting period
2015 PQRS Proposed New Measure	0710	Registry	Person and Caregiver-Centered Experience and Outcomes	Depression Response at Twelve Months-Progress Towards Remission: Adult patients age 18 and older with major depression or dysthymia and an initial PHQ-9 score > 9 who demonstrate a response to treatment at twelve months defined as a PHQ-9 score that is reduced by 50% or greater from the initial PHQ-9 score. This measure applies to both patients with newly diagnosed and existing depression identified during the defined measurement period whose current PHQ-9 score indicates a need for treatment

Potential Internal Medicine Preferred Specialty Measure Set

PQRS# CMS# GPRO#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
2015 PQRS Proposed New Measure	N/A	Registry	Person and Caregiver- Centered Experience and Outcomes	Discussion and Shared Decision Making Surrounding Treatment Options: Percentage of patients aged 18 years and older with a diagnosis of hepatitis C with whom a physician or other clinician reviewed the range of treatment options appropriate to their genotype and demonstrated a shared decision making approach with the patient. To meet the measure, there must be documentation in the patient record of a discussion between the physician/clinician and the patient that includes all of the following: •Treatment choices appropriate to genotype •Risks and benefits •Evidence of effectiveness •Patient preferences toward the outcome of the treatment
2015 PQRS Proposed New Measure	N/A	Registry	Community/Population Health	Screening for Hepatitis C Virus (HCV) for Patients at High Risk: Percentage of patients with one or more of the following: a history of injection drug use, patients who received blood transfusions prior to 1992, OR patients who were born in the years 1945–1965 who received a one-time hepatitis C virus (HCV) antibody test
2015 PQRS Proposed New Measure	N/A	Registry	Effective Clinical Care	Screening for Hepatocellular Carcinoma (HCC) in patients with Hepatitis C Cirrhosis: Percentage of patients aged 18 years and older with a diagnosis of chronic hepatitis C cirrhosis who were screened with either ultrasound, triple-contrast CT or triple-contrast MRI for hepatocellular carcinoma (HCC) at least once within the 12 month reporting period

This table is a resource of potential measures that may be applicable to eligible professionals providing clinical services for Internal Medicine.