## ICD-10 MS-DRGs Version 37.2 Effective August 01, 2020

In response to the COVID-19 pandemic, the Centers for Medicare & Medicaid Services (CMS) is implementing 12 new procedure codes to describe the introduction or infusion of therapeutics, including remdesivir and convalescent plasma, into the International Classification of Diseases, Tenth Revision, Procedure Coding System (ICD-10-PCS), effective August 01, 2020.

The ICD-10 MS-DRG Grouper assigns each case into an MS-DRG based on the reported diagnosis and procedure codes and demographic information (age, sex, and discharge status).

The ICD-10 Medicare code editor (MCE) Version 37.2 software uses edits to detect and report errors in the claims data for the ICD-10 codes reported to validate correct coding on claims for discharges on or after August 01, 2020.

The ICD-10 MS-DRG Grouper software package to accommodate these new codes, Version 37.2, is effective for discharges on or after August 01, 2020. As indicated in the table below, these codes will not affect the MS-DRG assignment.

Procedure	Description	*O.R.	MDC	MS-DRG
Code				
XW013F5	Introduction of Other New Technology Therapeutic	Ν		
	Substance into Subcutaneous Tissue, Percutaneous			
	Approach, New Technology Group 5			
XW033E5	Introduction of Remdesivir Anti-infective into	Ν		
	Peripheral Vein, Percutaneous Approach, New			
	Technology Group 5			
XW033F5	Introduction of Other New Technology Therapeutic	Ν		
	Substance into Peripheral Vein, Percutaneous			
	Approach, New Technology Group 5			
XW033G5	Introduction of Sarilumab into Peripheral Vein,	Ν		
	Percutaneous Approach, New Technology Group 5			
XW033H5	Introduction of Tocilizumab into Peripheral Vein,	Ν		
	Percutaneous Approach, New Technology Group 5			
XW043E5	Introduction of Remdesivir Anti-infective into Central	Ν		
	Vein, Percutaneous Approach, New Technology			
	Group 5			
XW043F5	Introduction of Other New Technology Therapeutic	Ν		
	Substance into Central Vein, Percutaneous Approach,			
	New Technology Group 5			
XW043G5	Introduction of Sarilumab into Central Vein,	Ν		
	Percutaneous Approach, New Technology Group 5			
XW043H5	Introduction of Tocilizumab into Central Vein,	N		
	Percutaneous Approach, New Technology Group 5			

Assignment of the new ICD-10-PCS procedure codes is as follows:

XW0DXF5	Introduction of Other New Technology Therapeutic Substance into Mouth and Pharynx, External Approach, New Technology Group 5	N	
XW13325	Transfusion of Convalescent Plasma (Nonautologous) into Peripheral Vein, Percutaneous Approach, New Technology Group 5	N	
XW14325	Transfusion of Convalescent Plasma (Nonautologous) into Central Vein, Percutaneous Approach, New Technology Group 5	N	

\*As the procedure codes are designated as non-O.R. procedures, there is no assigned MDC or MS-DRG. The ICD-10 MS-DRG assignment is dependent on the reported principal diagnosis, any secondary diagnoses defined as a complication or comorbidity (CC) or major complication or comorbidity (MCC), procedures or services performed, age, sex, and discharge status.

The ICD-10 MS-DRG V37.2 Grouper Software, Definitions Manual Table of Contents and the Definitions of Medicare Code Edits V37.2 manual will be available at <u>https://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/AcuteInpatientPPS/MS-DRG-Classifications-and-Software.html</u>

The Code Tables, Index and related Addenda files for the 12 new procedure codes are available at: <u>https://www.cms.gov/Medicare/Coding/ICD10/2020-ICD-10-PCS</u>