

## GG0130C. Toileting Hygiene

**C. Toileting hygiene:** The ability to maintain perineal hygiene, adjust clothes before and after voiding or having a bowel movement. If managing an ostomy, include wiping the opening but not managing the equipment.

### Assessment Timepoints

#### Admission:

- Admission performance.
- Discharge goal.

#### Discharge:

- Discharge performance.



### Toileting Hygiene Inclusions and Exclusions for Coding

#### Included:

- Managing undergarments, clothing, and incontinence products.
- Performing perineal/perianal cleansing before and after:
  - Voiding.
  - Bowel movements.
- Wiping the opening of an ostomy.
- Perineal cleansing around a catheter.

#### Excluded:

- Managing ostomy equipment.
- Managing catheter equipment.
- Transferring on/off the toilet, bedside commode, and/or bedpan.


### Coding Tips

Toileting hygiene takes place before and after voiding and/or having a bowel movement. Consider the patient's need for both managing clothing and perineal/perianal cleansing.

## Activity Performance Codes and Examples

06 – Independent	03 – Partial/moderate assistance
05 – Setup or clean-up assistance	02 – Substantial/maximal assistance
04 – Supervision or touching assistance	01 – Dependent

Each series of arrows below describes a toileting scenario. The green arrows in the second and fourth columns identify key elements of each scenario that inform the coding of **GG0130C. Toileting hygiene**.



## Additional Clinical Considerations

- If the patient does not typically use undergarments, code **GG0130C** based on the need for assistance to manage lower body clothing and perineal/perianal hygiene.
- If the patient has an indwelling urinary catheter, code **GG0130C** based on the type and amount of assistance needed before and after moving their bowels. This may necessarily include the need to perform perineal hygiene to the indwelling urinary catheter site after the bowel movement.
- If the patient completes a bowel toileting program in bed, code **GG0130C** based on the need for assistance to manage clothing and perineal/perianal cleansing.
- The toileting hygiene activity can be assessed and coded regardless of the patient's need to void or have a bowel movement.
- When the patient requires different levels of assistance to perform toileting hygiene after voiding vs. after a bowel movement, code based on the type and amount of assistance required to complete the ENTIRE activity.