

ICD-10-PCS to ICD-9-CM Procedure Version 23.0 Code Map

APPLICATION:

The ICD-10-PCS to ICD-9-CM map is used to facilitate converting ICD-10-PCS procedure codes to ICD-9-CM procedure codes. Every ICD-10-PCS code is mapped to one or more ICD-9-CM codes. The map contains possible options from which the appropriate ICD-9-CM code can be chosen, depending on the use to which the map is put.

FILES and FORMATS:

There is 1 file in this distribution for ICD-10-PCS to ICD-9-CM Code Mapping.

FILE #1 : GENERAL ICD-10-PCS to ICD-9-CM Mapping File

Filename: Map_pcsi9_all.txt

FIELD	POSITION	LENGTH
ICD-10-PCS Code	1 – 7	7
Filler	8	1
ICD-9-CM Procedure Code	9 – 12	4

ICD-9-CM Procedure Version 23.0 to ICD-10-PCS Code Map

APPLICATION:

The ICD-9-CM to ICD-10-PCS map is used to facilitate converting ICD-9-CM procedure codes to ICD-10-PCS procedure codes.

FILES and FORMATS:

There are 2 categories of mapping files for ICD-9-CM to ICD-10-PCS mapping:

1. General: The ICD-9-CM code can be mapped for a specific use with one ICD-10-PCS code. The map contains possible options from which the appropriate ICD-10-PCS code can be chosen, depending on the use to which the map is put. ICD-9-CM codes designated "No Equivalent" are not mapped to an ICD-10-PCS code. Reason indicators specify why the ICD-9-CM code is not mapped.
2. Combination: The ICD-9-CM code defines more than one procedure, and can only be fully mapped using more than one ICD-10-PCS code. The map contains lists of possible options from which the appropriate ICD-10-PCS codes can be chosen, depending on the use to which the map is put.

There are 3 files in this distribution for ICD-9-CM to ICD-10-PCS Code Mapping.

FILE #1 : GENERAL ICD-9-CM to ICD-10-PCS Mapping File

Filename: Map_i9pcs_all.txt

FIELD	POSITION	LENGTH
ICD-9-CM Procedure Code	1 – 4	4
Filler	5	1
ICD-10-PCS Code	6 – 12	7
Filler	13	1
"No Equivalent" Value	14 – 15	2

No Equivalent	Advice
1	Not Reportable: ICD-9-CM code considered not reportable for hospital inpatient procedure coding and therefore not mapped to any ICD-10-PCS code.
2	Component Code: ICD-9-CM code designated in the tabular list as <u>always</u> paired with a primary procedure code, and identified as a character (approach, device) in a single ICD-10-PCS code for the primary procedure, is therefore not mapped separately to an ICD-10-PCS code.
10	Procedure Not Performed: ICD-9-CM code for procedure determined as not performed or no longer performed and therefore not mapped to any ICD-10-PCS code.

FILE #2 i9.idmap: COMBINATION ICD-9-CM to LIST-ID Mapping File (Part 1 of 2)

Filename: Map_i9pcs_i9id.txt

FIELD	POSITION	LENGTH
ICD-9-CM Procedure Code	1 – 4	4
Filler	5	1
LIST ID's (up to 8)	6 – 45	40
LIST-ID #1	6 – 10	4
Filler (Blank)	11	1
LIST-ID #2	12 – 15	4
Filler (Blank)	16	1
LIST-ID #3	17 – 20	4
Filler (Blank)	21	1
LIST-ID #4	22 – 25	4
Filler (Blank)	26	1
LIST-ID #5	27 – 30	4
Filler (Blank)	31	1
LIST-ID #6	32 – 35	4
Filler (Blank)	36	1
LIST-ID #7	37 – 40	4
Filler (Blank)	41	1
LIST-ID #8	42 – 45	4

FILE #3 id.pcs.map: COMBINATION LIST-ID to ICD-10-PCS Mapping File (Part 2 of 2)

Filename: Map_i9pcs_idpcs.txt

FIELD	POSITION	LENGTH
LIST-ID	1 – 4	4
Filler	5	1
ICD-10-PCS Code	6 – 12	7

How to use the COMBINATION Mapping Files

For the ICD-9-CM procedures in the Combination category, more than one ICD-10-PCS procedure code is required to satisfy the equivalent meaning of a single ICD-9-CM procedure code.

The two files used are:

- The Combination ICD-9-CM to LIST-ID map [Map_i9pcs_i9id.txt]
This file contains one or more entries (lines) for the set of ICD-9-CM procedure codes that require a combination ICD-10-PCS map. If there is more than one entry for an ICD-9-CM procedure code, then meeting the criteria below for any one of the entries will satisfy the map.
 - Each entry will have from one to eight LIST-ID's. Each entry requires one ICD-10-PCS code from each of the LIST-ID's in order to satisfy the map.
 - If there is only one LIST-ID associated with the entry, then any ICD-10-PCS code on that list will satisfy the map.
 - If there is more than one LIST-ID associated with the entry, then one code from each of the LIST-ID's will satisfy the map.
- The Combination LIST-ID to ICD-10-PCS map [Map_i9pcs_idpcs.txt]
This file contains the ICD-10-PCS procedure codes for each LIST-ID.

EXAMPLE:

688 Pelvic Evisceration

There are two entries in the Combination ICD-9-CM to LIST-ID map file as follows:

I9CM	LIST-ID#1	LIST-ID#2	LIST-ID#3	LIST-ID#4	LIST-ID#5	LIST-ID#6	LIST-ID#7	LIST-ID#8
688	L324	L325	L326	L327	L328	L329		
688	L330	L331	L332	L333	L334	L335	L336	L337

The map would need:

EITHER :

one code from LIST-ID# L324
plus
one code from LIST-ID# L325
plus
one code from LIST-ID# L326
plus
one code from LIST-ID# L327
plus
one code from LIST-ID# L328
plus
one code from LIST-ID# L329

OR:

one code from LIST-ID# L330
plus
one code from LIST-ID# L331
plus
one code from LIST-ID# L332
plus
one code from LIST-ID# L333
plus
one code from LIST-ID# L334
plus
one code from LIST-ID# L335
plus
one code from LIST-ID# L336
plus
one code from LIST-ID# L337