


Figure 45: Distribution of single dwell volumes for adult peritoneal dialysis patients, by modality, October 2004–March 2005. 2005 ESRD CPM Project.




*Value suppressed because n < 11.

Source:
 2005 Annual Report
 ESRD Clinical Performance Measures Project




Figure 46: Distribution of 24-hour total infused dialysate volumes for adult peritoneal dialysis patients, by modality, October 2004–March 2005. 2005 ESRD CPM Project.



Source:
 2005 Annual Report
 ESRD Clinical Performance Measures Project



Figure 47: Distribution of the mean number of total exchanges for adult peritoneal dialysis patients, by modality, October 2004–March 2005. 2005 ESRD CPM Project.




*Value suppressed because n < 11.

Source:
2005 Annual Report
ESRD Clinical Performance Measures Project




Figure 48: Percent of adult peritoneal dialysis patients meeting 1997 NKF-DOQI guidelines for weekly Kt/V_{urea} and weekly creatinine clearance (PD Adequacy CPM III). 2005 ESRD CPM Project.



Source:
 2005 Annual Report
 ESRD Clinical Performance Measures Project




Figure 49: Distribution of mean hemoglobin values for adult peritoneal dialysis patients in the U.S., by race, October 2004–March 2005. 2005 ESRD CPM Project.



Source:
2005 Annual Report
ESRD Clinical Performance Measures Project


Figure 50: Percent of adult peritoneal dialysis patients with mean hemoglobin ≥ 11 g/dL, by race, October 2004–March 2005 compared to previous study periods. 2005 ESRD CPM Project.



Source:
2005 Annual Report
ESRD Clinical Performance Measures Project



Figure 51: Mean weekly epoetin dose (units/kg/week) by hemoglobin category for adult peritoneal dialysis patients prescribed epoetin, October 2004–March 2005 compared to previous study periods. 2005 ESRD CPM Project.




Note: Route of administration was not collected in 1998.

Source:
2005 Annual Report
ESRD Clinical Performance Measures Project




Figure 52: Percent of adult peritoneal dialysis patients with specific anemia management indicators, October 2004–March 2005 compared to selected previous study periods. 2005 ESRD CPM Project.



Source:
 2005 Annual Report
 ESRD Clinical Performance Measures Project



Figure 53: Percent of adult peritoneal dialysis patients with mean serum albumin $\geq 4.0/3.7$ g/dL (BCG/BCP) and $\geq 3.5/3.2$ g/dL (BCG/BCP), October 2004–March 2005 compared to previous study periods. 2005 ESRD CPM Project.*



*Note: BCG/BCP = bromcresol green/bromcresol purple laboratory methods.

Source:
2005 Annual Report
ESRD Clinical Performance Measures Project

