

Opioid Morphine Equivalent Conversion Factors¹

Type of Opioid	MME Conversion Factor
Buprenorphine patch ²	12.6
Buprenorphine tab or film	10
Butorphanol	7
Codeine	0.15
Dihydrocodeine	0.25
Fentanyl buccal or SL tablets, or lozenge/troche ³	0.13
Fentanyl film or oral spray ⁴	0.18
Fentanyl nasal spray ⁵	0.16
Fentanyl patch ⁶	7.2
Hydrocodone	1
Hydromorphone	4
Levorphanol tartrate	11
Meperidine hydrochloride	0.1
Methadone	3
Morphine	1
Nalbuphine	1
Opium	1
Oxycodone	1.5
Oxymorphone	3
Pentazocine	0.37
Tapentadol	0.4
Tramadol	0.1

¹The MME conversion factor is intended only for analytic purposes where prescription data is used to calculate daily MME. This value does not constitute clinical guidance or recommendations for converting patients from one form of opioid analgesic to another. Please consult the manufacturer's full prescribing information for such guidance. Use of this file for the purposes of any clinical decision-making warrants caution. Centers for Disease Control and Prevention, Atlanta, GA, May 2015. For more information, send an email to Mbohm@cdc.gov.

²The MME conversion factor for buprenorphine patches is based on the assumption that one milligram of parenteral buprenorphine is equivalent to 75 milligrams of oral morphine and that one patch delivers the dispensed micrograms per hour over a 24 hour day. Example: 5 ug/hr buprenorphine patch * 24 hrs = 120 ug/day buprenorphine = 0.12 mg/day buprenorphine = 9 mg/day oral morphine milligram equivalent. In other words, the conversion factor not accounting for days of use would be 9/5 or 1.8. However, since the buprenorphine patch remains in place for 7 days, we have multiplied the conversion factor by 7 (1.8 X 7 = 12.6). In this example, MME/day for four 5 ug/hr buprenorphine patches dispensed for use over 28 days would work out as follows: Example: 5 ug/hr buprenorphine patch * (4 patches/28 days) * 12.6 = 9 MME/day.

³The MME conversion factor for fentanyl buccal tablets, sublingual tablets, and lozenges/troche is 0.13. This conversion factor should be multiplied by the number of micrograms in a given lozenge/troche.

⁴The MME conversion factor for fentanyl film and oral spray is 0.18. This reflects a 40% greater bioavailability for films compared to lozenges/tablets and 38% greater bioavailability for oral sprays compared to lozenges/tablets.

⁵The MME conversion factor for fentanyl nasal spray is 0.16, which reflects a 20% greater bioavailability for sprays compared to lozenges/tablets.

⁶The MME conversion factor for fentanyl patches is based on the assumption that one milligram of parenteral fentanyl is equivalent to 100 milligrams of oral morphine and that one patch delivers the dispensed micrograms per hour over a 24 hour day. Example: 25 ug/hr fentanyl patch * 24 hrs = 600 ug/day fentanyl = 60 mg/day oral morphine milligram equivalent. In other words, the conversion factor not accounting for days of use would be 60/25 or 2.4. However, since the fentanyl patch remains in place for 3 days, we have multiplied the conversion factor by 3 (2.4 X 3 = 7.2). In this example, MME/day for ten 25 ug/hr fentanyl patches dispensed for use over 30 days would work out as follows: Example: 25 ug/hr fentanyl patch * (10 patches/30 days) * 7.2 = 60 MME/day.