

Home Health Quality Reporting Program M1311 and M1313 PRACTICE SCENARIO SHEETS



May 3–4, 2017
Radisson Hotel Baltimore
Downtown-Inner Harbor
Baltimore, MD

M1311 Practice Scenario (1)

- At SOC, the patient has three small Stage 2 pressure ulcers in her sacral area.
- Upon discharge, the sacrum is assessed.
- Two of the Stage 2 pressure ulcers have merged, and the third ulcer has increased to Stage 3.

How would you respond to M1311?

ITEM	SOC/ROC	DISCHARGE
M1311A1. Number of Stage 2 pressure ulcers		
M1311A2. Number of these Stage 2 pressure ulcers that were present at most recent SOC/ROC		
M1311B1. Number of Stage 3 pressure ulcers		
M1311B2. Number of these Stage 3 pressure ulcers that were present at most recent SOC/ROC		

Refer to Presentation Slides 72–80.



M1311 Practice Scenario (2)

- At the SOC, Mrs. J is admitted with four Stage 2 pressure ulcers and one Stage 3 pressure ulcer.
- During Mrs. J's home health quality episode, three of the Stage 2 pressure ulcers healed. However, Mrs. J develops an additional Stage 3 pressure ulcer.

How would you respond to M1311?

ITEM	SOC/ROC	DISCHARGE
M1311A1. Number of Stage 2 pressure ulcers		
M1311A2. Number of these Stage 2 pressure ulcers that were present at most recent SOC/ROC		
M1311B1. Number of Stage 3 pressure ulcers		
M1311B2. Number of these Stage 3 pressure ulcers that were present at most recent SOC/ROC		

Refer to Presentation Slides 81–89.

M1311 Practice Scenario (3)

- At the SOC, Ms. P was admitted with a diagnosis of cerebrovascular accident with right hemiparesis and assessed to have a 1 cm × 1 cm × < 0.1 cm Stage 2 pressure ulcer on her coccyx.
- Ms. P continued to decline at home, with decreased appetite, frequent transient ischemic attacks, and a wish not to be hospitalized again. After a palliative care consult, the patient and family agreed to hospice care.
- Upon discharge from home care, Ms. P was noted to have a pressure ulcer completely covered with eschar on her left heel and a Stage 3 pressure ulcer 3 cm × 2 cm × 0.4 cm on her coccyx.

How would you respond to M1311?

ITEM	SOC/ROC	DISCHARGE
M1311A1. Number of Stage 2 pressure ulcers		
M1311A2. Number of these Stage 2 pressure ulcers that were present at most recent SOC/ROC		
M1311B1. Number of Stage 3 pressure ulcers		
M1311B2. Number of these Stage 3 pressure ulcers that were present at most recent SOC/ROC		
M1311C1. Number of Stage 4 pressure ulcers		
M1311C2. Number of these Stage 4 pressure ulcers that were present at most recent SOC/ROC		
M1311D1. Number of Unstageable pressure ulcers: Non-removable dressing		
M1311D2. Number of these unstageable pressure ulcers that were present at most recent SOC/ROC		

M1311 Practice Scenario (3) continued

How would you respond to M1311?

ITEM	SOC/ROC	DISCHARGE
M1311E1. Number of Unstageable pressure ulcers: Slough and/or eschar		
M1311E2. Number of these unstageable pressure ulcers that were present at most recent SOC/ROC		
M1311F1. Number of Unstageable pressure ulcers: Deep tissue injury		
M1311F2. Number of these unstageable pressure ulcers that were present at most recent SOC/ROC		

Refer to Presentation Slides 90–110.

M1311 & M1313 Practice Scenario (1)

- A patient was admitted to home care from the acute care hospital with two Stage 2 pressure ulcers with clear fluid-filled blisters, one on each heel.
- After a few days, the left heel Stage 2 blister had ruptured and presented as a shallow ulcer with a pink wound bed. The right heel continued to evolve, having a blood-filled blister, and matured in color from red to a maroon/purple color, with the area surrounding the blister being boggy, painful, and warm.
- The patient stated she was moving and going to stay with her daughter who lives 2 hours away. She was discharged to another provider.
- Upon discharge, the left heel had a shallow ulcer with a pink wound bed. The right heel was covered in firm dark eschar with slight bogginess around the eschar.

How would you respond to M1311 and M1313?

ITEM	SOC/ROC	DISCHARGE
M1311A1. Number of Stage 2 pressure ulcers		
M1311A2. Number of these Stage 2 pressure ulcers present at most recent SOC/ROC		
M1311B1. Number of Stage 3 pressure ulcers		
M1311B2. Number of these Stage 3 pressure ulcers that were present at most recent SOC/ROC		
M1313A. Worsening in Pressure Ulcer Status Since SOC/ROC – Stage 2		
M1313B. Worsening in Pressure Ulcer Status Since SOC/ROC – Stage 3		
M1313C. Worsening in Pressure Ulcer Status Since SOC/ROC – Stage 4		
M1313D. Worsening in Pressure Ulcer Status Since SOC/ROC – Unstageable - Non-removable dressing		
M1313E. Worsening in Pressure Ulcer Status Since SOC/ROC – Unstageable - Slough and/or Eschar		
M1313F. Worsening in Pressure Ulcer Status Since SOC/ROC – Unstageable - Deep tissue injury		

Refer to Presentation Slides 131–145.

M1311 and M1313 Practice Scenario (2)

- At SOC, Mr. Y is noted to have a Stage 3 pressure ulcer on his right hip.
- During the home health quality episode, the pressure ulcer identified at SOC closes, but Mr. Y develops a new Stage 3 pressure ulcer on his left hip.
- At discharge, Mr. Y has one Stage 3 pressure ulcer on the left hip.

How would you respond to M1311 & M1313?

ITEM	SOC/ROC	DISCHARGE
M1311A1. Number of Stage 2 pressure ulcers		
M1311A2. Number of these Stage 2 pressure ulcers present at most recent SOC/ROC		
M1311B1. Number of Stage 3 pressure ulcers		
M1311B2. Number of these Stage 3 pressure ulcers that were present at most recent SOC/ROC		
M1313A. Worsening in Pressure Ulcer Status Since SOC/ROC – Stage 2		
M1313B. Worsening in Pressure Ulcer Status Since SOC/ROC – Stage 3		
M1313C. Worsening in Pressure Ulcer Status Since SOC/ROC – Stage 4		
M1313D. Worsening in Pressure Ulcer Status Since SOC/ROC – Unstageable - Non-removable dressing		
M1313E. Worsening in Pressure Ulcer Status Since SOC/ROC – Unstageable - Slough and/or Eschar		
M1313F. Worsening in Pressure Ulcer Status Since SOC/ROC – Unstageable - Deep tissue injury		

Refer to Presentation Slides 146–161.

M1311 & M1313 Practice Scenario (3)

- A patient was admitted to home care with two Stage 2 pressure ulcers.
- A mutual decision was made to transition the patient to hospice care.
- By the time of discharge, the two pressure ulcers had merged and increased to Stage 3.

Polling Question: How would you complete M1311 at Discharge?

A. M1311A1. Stage 2 = 2
M1311A2 = 2

B. M1311B1. Stage 3 = 1
M1311B2 = 0

C. M1311B1. Stage 3 = 1
M1311B2 = 1

Use the polling device on your table to make your response selection.

Polling Question: How would you complete M1313?

A. M1313A. Stage 2 = 1

B. M1313B. Stage 3 = 1

C. M1313C. Stage 4 = 1

Use the polling device on your table to make your response selection.

Refer to Presentation Slides 162–165.

M1311 & M1313 Practice Scenario (4)

- A patient developed a Stage 3 pressure ulcer during their home health quality episode.
- The wound bed was subsequently covered with slough; hence, the pressure ulcer became unstageable.
- The patient record indicates that wound debridement was performed on the Stage 3 pressure ulcer 2 weeks prior to discharge.
- During the discharge assessment, the wound bed was observed and numerically staged as a Stage 3.

Polling Question: How would you complete M1311 at Discharge?

A. M1311B1. Stage 3 = 1
M1311B2 = 1

B. M1311B1. Stage 3 = 1
M1311B2 = 0

C. M1311E1. Unstageable due to slough/eschar = 1
M1311E2 = 0

Use the polling device on your table to make your response selection.

Polling Question: How would you complete M1313?

A. M1313B. Stage 3 = 1

B. M1313C. Stage 4 = 1

C. M1313E. Unstageable due to slough/eschar = 1

Use the polling device on your table to make your response selection.

Refer to Presentation Slides 166–169.