

**HOME HEALTH AGENCY QUALITY MEASURES:
TECHNICAL DOCUMENTATION OF
POTENTIALLY AVOIDABLE EVENT MEASURES**

OASIS-D

January 2019

Department of Health and Human Services
Centers for Medicare & Medicaid Services

CONTENTS

| | <u>Page</u> |
|--|-------------|
| Introduction | 1 |
| Quality Episodes of Care for OASIS Quality Measures | 1 |
| Home Health Potentially Avoidable Event Measures | 3 |
| Assumptions and Conventions | 3 |
| Data Transformation Pseudo-code | 5 |
| Emergent Care for Potentially Avoidable Cause | 5 |
| Emergent Care for Improper Medication Administration or Medication Side Effects | 5 |
| Emergent Care for Hypo/Hyperglycemia | 5 |
| Discharged with New, Worsening, or Unresolved Health Problem | 5 |
| Development of Urinary Tract Infection | 5 |
| Increase in Number of Pressure Ulcers..... | 6 |
| Substantial Decline in 3 or more Activities of Daily Living | 6 |
| Substantial Decline in Management of Oral Medications | 8 |
| Discharged to the Community with an Unhealed Stage II Pressure Ulcer | 8 |
| Discharged with Unmet Needs | 9 |
| Discharged to the Community Needing Wound Care or Medication Assistance .. | 9 |
| Discharge to the Community Needing Toileting Assistance | 10 |
| Discharge to the Community with Behavioral Problems | 10 |

HOME HEALTH AGENCY QUALITY MEASURES: TECHNICAL DOCUMENTATION OF POTENTIALLY AVOIDABLE EVENT MEASURES

INTRODUCTION

This document provides definitions of measures that appear in Home Health Agency Potentially Avoidable Event Reports, including detailed specifications for calculating these measures based on OASIS-D assessment data.

Quality Episodes of Care for OASIS Quality Measures

A quality episode of care is the unit of analysis for OASIS-based measures. This section provides the steps to construct a home health quality episode of care from OASIS assessments. The following fields are needed from each assessment to construct home health quality episodes of care:

- M0100_ASSMT_REASON: Reason for assessment
- FAC_INT_ID: Facility internal ID
- RES_INT_ID: Resident internal ID
- STATE_CD: State abbreviation
- EFCTV_DT: Effective date of the assessment¹
- ASSESSMENT_ID: Unique identifier for each assessment

OASIS assessments are transmitted to the National Submissions Database residing on a secure database server maintained by CMS. Episodes of care are constructed by matching up assessments for each individual served by a home health agency, sorting those assessments by effective date, then pairing up assessments that mark the beginning and end of an episode of care. Episodes of care for which either the beginning or end assessment is missing, or for which assessments are out of sequence, are not included.

When working with assessment records from the National Submissions Database, the unique home health agency ID is a combination of the two position STATE_CD and FAC_INT_ID, and the unique patient ID is a combination of STATE_CD and RES_INT_ID.

The process of building episode-level records is as follows:

¹ Effective date depends on the reason for assessment:

- If M0100_ASSMT_REASON = 01, effective date = M0030_START_CARE_DT.
- If M0100_ASSMT_REASON = 03, effective date = M0032_ROC_DATE.
- If M0100_ASSMT_REASON = 04 or 05, effective date = M0090_INFO_COMPLETET_DT.
- If M0100_ASSMT_REASON = 06, 07, 08, or 09, effective date = M0906_DC_TRAN_DTH_DT.

1. Keep only assessments related to start of care (SOC), resumption of care (ROC) after an inpatient facility stay, or end of care (EOC) discharge from home health care, including discharge due to death, or admission to inpatient facility for 24 hours or more:
 - a. SOC: M0100_ASSMT_REASON = 01
 - b. ROC: M0100_ASSMT_REASON = 03
 - c. EOC: M0100_ASSMT_REASON = 06, 07, 08, or 09
2. Sort assessments by FAC_INT_ID, RES_INT_ID, STATE_CD in descending order based on EFCTV_DT (i.e., latest to earliest assessment).
3. For each set of assessments having the same combination of FAC_INT_ID, RES_INT_ID, STATE_CD, step through the assessments to find the latest assessment with M0100_ASSMT_REASON = 06, 07, 08, or 09.
4. If an assessment with M0100_ASSMT_REASON = 06, 07, 08, or 09 is found before an assessment with M0100_ASSMT_REASON = 01 or 03 is found, discard the episode being built and start over with step #3.
5. If no assessment with M0100_ASSMT_REASON = 01 or 03 is found before the assessments for this patient are exhausted, discard the episode being built.
6. When an assessment with M0100_ASSMT_REASON = 01 or 03 is found, the episode is complete.
7. Create an episode of care record with the following attributes:
 - FAC_INT_ID
 - RES_INT_ID
 - STATE_CD
 - ASSESSMENT_ID_1: Assessment ID of the SOC or ROC assessment
 - ASSESSMENT_ID_2: Assessment ID of the EOC assessment
 - EFCTV_DT_1: Effective date of the SOC or ROC assessment
 - EFCTV_DT_2: Effective date of the EOC assessment

Home Health Potentially Avoidable Event Measures

Measures are referred to as follows:

| <u>Measure Domain & Descriptive Measure Name</u> | <u>Short Measure Name</u> |
|---|---------------------------|
| Emergent Care for Potentially Avoidable Cause | |
| Emergent Care for Improper Medication Administration or Medication Side Effects | ED_Use_Medications |
| Emergent Care for Hypo/Hyperglycemia | ED_Use_Blood_Sugar |
| Discharged with New, Worsening, or Unresolved Health Problem | |
| Development of Urinary Tract Infection | New_UTI |
| Increase in Number of Pressure Ulcers | More_Pressure_Ulcers |
| Substantial Decline in 3 or more Activities of Daily Living | ADL_Decline |
| Substantial Decline in Management of Oral Medications | Oral_Med_Decline |
| Discharged to the Community with an Unhealed Stage II Pressure Ulcer | DC_w_Stg_II_Press_Ulc |
| Discharged with Unmet Needs | |
| Discharged to the Community Needing Wound Care or Medication Assistance | DC_Need_Wound_Care |
| Discharged to the Community Needing Toileting Assistance | DC_Need_Toilet_Assist |
| Discharged to the Community with Behavioral Problems | DC_Behavior_Problem |

Assumptions and Conventions

The following transformations result in measures with numeric values. For process and outcome measures, including potentially avoidable event measures, "1" indicates that the outcome event occurred or the process was followed, while "0" indicates the outcome event did not occur or the process was not followed. When the instructions indicate that a measure is to be set to "MISSING", the case is excluded from analysis for that particular measure, either because the measure is not applicable or because a data item on which that measure is based has been indicated as "unknown".

OASIS data items are referred to in this documentation using field names specified in OASIS Data Submission Specifications published by CMS. "[1]" is appended to the field name if the value is taken from the beginning of episode assessment (Start or Resumption of Care), and "[2]" is appended if the value is to be taken from the assessment conducted at the end of the episode (Discharge, Transfer, or Death). The

expression VALUE(FieldName) indicates conversion of a one or two character text field to a numeric value. The expression DATEVALUE(FieldName) indicates conversion of an eight character text field to a numeric value representing a date (number of days since 12/31/1899). Starting with OASIS D, dash '-' may be used as a response for certain items.

Data Transformation Pseudo-code

Emergent Care for Potentially Avoidable Cause

Emergent Care for Improper Medication Administration or Medication Side Effects

```
IF M2310_ECR_MEDICATION[2] = 1
THEN
    ED_USE_MEDICATIONS = 1
ELSEIF M2301_EMER_USE_AFTR_LAST_ASMT[2] = 00 OR
((M2301_EMER_USE_AFTR_LAST_ASMT[2] =01 OR
M2301_EMER_USE_AFTR_LAST_ASMT[2] =02) AND
M2310_ECR_UNKNOWN[2] = 0)
THEN
    ED_USE_MEDICATIONS = 0
ELSE
    ED_USE_MEDICATIONS = MISSING
END IF
```

Emergent Care for Hypo/Hyperglycemia

```
IF M2310_ECR_HYPOGLYC[2] = 1
THEN
    ED_USE_BLOOD_SUGAR = 1
ELSEIF M2301_EMER_USE_AFTR_LAST_ASMT[2] = 00 OR
((M2301_EMER_USE_AFTR_LAST_ASMT[2] =01 OR
M2301_EMER_USE_AFTR_LAST_ASMT[2] =02) AND
M2310_ECR_UNKNOWN[2] = 0)
THEN
    ED_USE_BLOOD_SUGAR = 0
ELSE
    ED_USE_BLOOD_SUGAR = MISSING
END IF
```

Discharged with New, Worsening, or Unresolved Health Problem

Development of Urinary Tract Infection

```
IF M0100_ASSMT_REASON[2] = 09
THEN
    IF M1600_UTI[1] = 00 AND M1600_UTI[2] = 01
    THEN
        NEW_UTI = 1
    ELSEIF M1600_UTI[1] = 00 AND M1600_UTI[2] = 00
    THEN
        NEW_UTI = 0
```

```

ELSE
    NEW_UTI = MISSING
END IF
ELSE
    NEW_UTI = MISSING
END IF

```

Increase in Number of Pressure Ulcers

```

IF M0100_ASSMT_REASON[2] = 09
THEN
    IF M1306_UNHLD_STG2_PRSR_ULCR[2] = 1
    THEN
        IF M1306_UNHLD_STG2_PRSR_ULCR[1] = 0
        THEN
            More_Pressure_Ulcers = 1
        ELSEIF (VALUE(M13011_NBR_PRSULC_STG2[1]) +
            VALUE(M13011_NBR_PRSULC_STG3[1]) +
            VALUE(M1311_NBR_PRSULC_STG4[1]) +
            VALUE(M1311_NSTG_DRSG[1]) +
            VALUE(M1311_NSTG_CVRG[1]) +
            VALUE(M1311_NSTG_DEEP_TISUE[1])) <
            (VALUE(M1311_NBR_PRSULC_STG2[2]) +
            VALUE(M1311_NBR_PRSULC_STG3[2]) +
            VALUE(M1311_NBR_PRSULC_STG4[2]) +
            VALUE(M1311_NSTG_DRSG[2]) +
            VALUE(M1311_NSTG_CVRG[2]) +
            VALUE(M1311_NSTG_DEEP_TISUE[2]))
        THEN
            More_Pressure_Ulcers = 1
        ELSE
            More_Pressure_Ulcers = 0
        END IF
    ELSE
        More_Pressure_Ulcers = 0
    END IF
ELSE
    More_Pressure_Ulcers = MISSING
END IF

```

Substantial Decline in 3 or more Activities of Daily Living

```

ADL_Temp1 = 0
IF M1800_CRNT_GROOMING[1] = 03 OR M1800_CRNT_GROOMING[1] = 02
THEN

```

```

    ADL_Temp1 = ADL_Temp1 + 1
END IF
IF M1830_CRNT_BATHG[1] = 06 OR M1830_CRNT_BATHG[1] = 05
THEN
    ADL_Temp1 = ADL_Temp1 + 1
END IF
IF M1840_CRNT_TOILTG[1] = 04 OR M1840_CRNT_TOILTG[1] = 03
THEN
    ADL_Temp1 = ADL_Temp1 + 1
END IF
IF M1845_CRNT_TOILTG_HYGN[1] = 03 OR M1845_CRNT_TOILTG_HYGN[1] = 02
THEN
    ADL_Temp1 = ADL_Temp1 + 1
END IF
IF M1850_CRNT_TRNSFRNG[1] = 05 OR M1850_CRNT_TRNSFRNG[1] = 04
THEN
    ADL_Temp1 = ADL_Temp1 + 1
END IF
IF M1860_CRNT_AMBLTN[1] = 06 OR M1860_CRNT_AMBLTN[1] = 05
THEN
    ADL_Temp1 = ADL_Temp1 + 1
END IF
IF M1870_CRNT_FEEDING[1] = 05 OR M1870_CRNT_FEEDING[1] = 04
THEN
    ADL_Temp1 = ADL_Temp1 + 1
END IF
IF M0100_ASSMT_REASON[2] = 09 AND ADL_Temp1 <5
THEN
    ADL_Temp2 = 0
    IF VALUE(M1800_CRNT_GROOMING[2]) > VALUE(M1800_CRNT_GROOMING[1]) + 1
    THEN
        ADL_Temp2 = ADL_Temp2 + 1
    END IF
    IF VALUE(M1830_CRNT_BATHG[2]) > VALUE(M1830_CRNT_BATHG[1]) + 1
    THEN
        ADL_Temp2 = ADL_Temp2 + 1
    END IF
    IF VALUE(M1840_CRNT_TOILTG[2]) > VALUE(M1840_CRNT_TOILTG[1]) + 1
    THEN
        ADL_Temp2 = ADL_Temp2 + 1
    END IF
    IF VALUE(M1845_CRNT_TOILTG_HYGN[2]) >
        VALUE(M1845_CRNT_TOILTG_HYGN[1]) + 1

```

```

THEN
    ADL_Temp2 = ADL_Temp2 + 1
END IF
IF VALUE(M1850_CRNT_TRNSFRNG[2]) > VALUE(M1850_CRNT_TRNSFRNG[1]) + 1
THEN
    ADL_Temp2 = ADL_Temp2 + 1
END IF
IF VALUE(M1860_CRNT_AMBLTN[2]) > VALUE(M1860_CRNT_AMBLTN[1]) + 1
THEN
    ADL_Temp2 = ADL_Temp2 + 1
END IF
IF VALUE(M1870_CRNT_FEEDING[2]) > VALUE(M1870_CRNT_FEEDING[1]) + 1
THEN
    ADL_Temp2 = ADL_Temp2 + 1
END IF
IF ADL_Temp2 >= 3
THEN
    ADL_Decline = 1
ELSE
    ADL_Decline = 0
END IF
ELSE
    ADL_Decline = MISSING
END IF

```

Substantial Decline in Management of Oral Medications

```

IF M0100_ASSMT_REASON[2] <> 09 OR
    M2020_CRNT_MGMT_ORAL_MDCTN[2] = NA OR
    M2001_DRUG_RGMN_RVW[1] = 9 OR
    M2020_CRNT_MGMT_ORAL_MDCTN[1] <> 00
THEN
    Oral_Med_Decline = MISSING
ELSE
    IF M2020_CRNT_MGMT_ORAL_MDCTN[2] = 03 AND
        M2020_CRNT_MGMT_ORAL_MDCTN[1] = 00
    THEN
        Oral_Med_Decline = 1
    ELSE
        Oral_Med_Decline = 0
    END IF
END IF

```

Discharged to the Community with an Unhealed Stage II Pressure Ulcer

```

IF M0100_ASSMT_REASON[2] = 09

```

```

THEN
  IF M2420_DSCHRG_DISP[2] = 01 AND
    (M1306_UNHLD_STG2_PRSR_ULCR[2] = 1 AND
    ((M1307_OLDST_STG2_AT_DSCHRG[2] = 02 AND
    DATEVALUE(M0906_DC_TRAN_DTH_DT[2]) -
    DATEVALUE(M1307_OLDST_STG2_ONST_DT[2]) > 30) OR
    (M1307_OLDST_STG2_AT_DSCHRG[2] = 01 AND
    M0100_ASSMT_REASON[1] = 01 AND
    DATEVALUE(M0906_DC_TRAN_DTH_DT[2]) -
    DATEVALUE(M0030_START_CARE_DT[1]) > 30) OR
    (M1307_OLDST_STG2_AT_DSCHRG[2] = 01 AND
    M0100_ASSMT_REASON[1] = 03 AND
    DATEVALUE(M0906_DC_TRAN_DTH_DT[2]) -
    DATEVALUE(M0032_ROC_DT[1]) > 30)
  THEN
    DC_w_Stg_II_Press_Ulc = 1
  ELSE
    DC_w_Stg_II_Press_Ulc = 0
  END IF
ELSE
  DC_w_Stg_II_Press_Ulc = MISSING
END IF

```

Discharged with Unmet Needs

Discharged to the Community Needing Wound Care or Medication Assistance

```

IF M0100_ASSMT_REASON[2] = 09
THEN
  IF M2420_DSCHRG_DISP[2] = 01 AND (VALUE(M2102_CARE_TYPE_SRC_MDCTN[2])
  > 2 OR VALUE(M2102_CARE_TYPE_SRC_PRCDR[2]) > 2) AND
  (M1710_WHEN_CONFUSED[2] = NA OR VALUE(M1710_WHEN_CONFUSED[2]) >
  2) AND (M1324_STG_PRBLM_ULCER[2] = 03 OR
  M1324_STG_PRBLM_ULCER[2] = 04 OR M1334_STUS_PRBLM_STAS_ULCR[2] = 03
  OR M1342_STUS_PRBLM_SRGCL_WND[2] = 03 OR
  M2020_CRNT_MGMT_ORAL_MDCTN[2] = 03)
  THEN
    DC_Need _Wound_Care = 1
  ELSE
    DC_Need _Wound_Care = 0
  END IF
ELSE
  DC_Need _Wound_Care = MISSING
END IF

```

Discharge to the Community Needing Toileting Assistance

```
IF M0100_ASSMT_REASON[2] = 09
THEN
  IF M2420_DSCHRG_DISP[2] = 01 AND (VALUE(M2102_CARE_TYPE_SRC_ADL[2]) > 2)
    AND (VALUE(M1840_CRNT_TOILTG[2]) = 4 OR
    VALUE(M1845_CRNT_TOILTG_HYGN[2] = 3))
  THEN
    DC_Need_Toilet_Assist = 1
  ELSE
    DC_Need_Toilet_Assist = 0
  END IF
ELSE
  DC_Need_Toilet_Assist = MISSING
END IF
```

Discharge to the Community with Behavioral Problems

```
IF M0100_ASSMT_REASON[2] = 09
THEN
  IF M2420_DSCHRG_DISP[2] = 01 AND
    VALUE(M2102_CARE_TYPE_SRC_SPRVSN[2]) > 2 AND
    (VALUE(M1740_BD_MEM_DEFICIT[2]) +
    VALUE(M1740_BD_IMP_DECISN[2]) +
    VALUE(M1740_BD_VERBAL[2]) +
    VALUE(M1740_BD_PHYSICAL[2]) +
    VALUE(M1740_BD_SOC_INAPPRO[2]) +
    VALUE(M1740_BD_DELUSIONS[2])) > 1
  THEN
    DC_Behavior_Problem = 1
  ELSE
    DC_Behavior_Problem = 0
  END IF
ELSE
  DC_Behavior_Problem = MISSING
END IF
```