

Proposed 2016 OB/GYN Preferred Specialty Measure Set

PQRS# CMS# GPRO#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
39	0046	Claims, Registry Measures Group (Prev Care)	Effective Clinical Care	Screening or Therapy for Osteoporosis for Women Aged 65 Years and Older: Percentage of female patients aged 65 years and older who have a central dual-energy X-ray absorptiometry (DXA) measurement ordered or performed at least once since age 60 or pharmacologic therapy prescribed within 12 months
41	N/A	Claims, Registry	Effective Clinical Care	Osteoporosis: Pharmacologic Therapy for Men and Women Aged 50 Years and Older: Percentage of patients aged 50 years and older with a diagnosis of osteoporosis who were prescribed pharmacologic therapy within 12 months
48	N/A	Claims, Registry Measures Group (Prev Care)	Effective Clinical Care	Urinary Incontinence: Assessment of Presence or Absence of Urinary Incontinence in Women Aged 65 Years and Older: Percentage of female patients aged 65 years and older who were assessed for the presence or absence of urinary incontinence within 12 months
50	N/A	Claims, Registry	Person and Caregiver-Centered Experience and Outcomes	Urinary Incontinence: Plan of Care for Urinary Incontinence in Women Aged 65 Years and Older: Percentage of female patients aged 65 years and older with a diagnosis of urinary incontinence with a documented plan of care for urinary incontinence at least once within 12 months
112 CMS125v2	0031	Claims, Registry, EHR, GPRO Web Interface, Measures Group (Prev Care)	Effective Clinical Care	Breast Cancer Screening: Percentage of women 50 through 74 years of age who had a mammogram to screen for breast cancer within 27 months
204	0068	Claims, Registry, EHR, GPRO (Web Interface), Measures Group (Million Hearts)	Effective Clinical Care	Ischemic Vascular Disease (IVD): Use of Aspirin or Another Antithrombotic: Percentage of patients 18 years of age and older who were discharged alive for acute myocardial infarction (AMI), coronary artery bypass graft (CABG) or percutaneous coronary interventions (PCI) in the 12 months prior to the measurement period, or who had an active diagnosis of ischemic vascular disease (IVD) during the measurement period and who had documentation of use of aspirin or another antithrombotic during the measurement period

Proposed 2016 OB/GYN Preferred Specialty Measure Set

PQRS# CMS# GPRO#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
226	0028	Claims, Registry, EHR, GPRO (Web Interface), Measures Groups	Community/ Population Health	Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention: Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 24 months AND who received cessation counseling intervention if identified as a tobacco user
236	0018	Claims, Registry, EHR, GPRO (Web Interface), Measures Group (Million Hearts)	Effective Clinical Care	Controlling High Blood Pressure: Percentage of patients 18-85 years of age who had a diagnosis of hypertension and whose blood pressure was adequately controlled (<140/90 mmHg) during the measurement period
265	N/A	Registry	Communication and Care Coordination	Biopsy Follow-Up: Percentage of patients whose biopsy results have been reviewed and communicated to the primary care/referring physician and patient by the performing physician
309 CMS124v2	0032	EHR, Registry	Effective Clinical Care	Cervical Cancer Screening: Percentage of women 21-64 years of age, who received one or more Pap tests to screen for cervical cancer
310 CMS153	0033	EHR	Community/Population Health	Chlamydia Screening: Percentage of women 16-24 years of age who were identified as sexually active and who had at least one test for chlamydia during the measurement period
317	N/A	Claims, Registry, EHR, GPRO (Web Interface), Measures Groups	Community/ Population Health	Preventive Care and Screening: Screening for High Blood Pressure and Follow-Up Documented: Percentage of patients aged 18 years and older seen during the reporting period who were screened for high blood pressure AND a recommended follow-up plan is documented based on the current blood pressure (BP) reading as indicated
Proposed New for 2016 PQRS	N/A	Claims, Registry	Patient Safety	Performing Cystoscopy at the time of Hysterectomy for Pelvic Organ Prolapse to Detect Lower Urinary Tract Injury: Percentage of patients who undergo cystoscopy to evaluate for lower urinary tract injury at the time of hysterectomy for pelvic organ prolapse
Proposed New for 2016 PQRS	N/A	Claims, Registry	Patient Safety	Proportion of Patients Sustaining a Bladder Injury at the Time of any Pelvic Organ Prolapse Repair: Percentage of patients undergoing any surgery to repair pelvic organ prolapse who sustains an injury to the bladder recognized either during or within 1 month after surgery

Proposed 2016 OB/GYN Preferred Specialty Measure Set

PQRS# CMS# GPRO#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
Proposed New for 2016 PQRS	N/A	Claims, Registry	Patient Safety	Proportion of Patients Sustaining a Major Viscus Injury at the Time of any Pelvic Organ Prolapse Repair: Percentage of patients undergoing surgical repair of pelvic organ prolapse that is complicated by perforation of a major viscus at the time of index surgery that is recognized intraoperative or within 1 month after surgery
Proposed New for 2016 PQRS	N/A	Claims, Registry	Patient Safety	Proportion of Patients Sustaining A Ureter Injury at the Time of any Pelvic Organ Prolapse Repair: Percentage of patients undergoing a pelvic organ prolapse repair who sustain an injury to the ureter recognized either during or within 1 month after surgery
Proposed New for 2016 PQRS	0053	Claims, Registry	Effective Clinical Care	Osteoporosis Management in Women Who Had a Fracture: The percentage of women age 50-85 who suffered a fracture and who either had a bone mineral density test or received a prescription for a drug to treat osteoporosis

This table is a resource of potential measures that may be applicable to OB/GYN. Please note that in addition to this table, a 2016 Proposed Cross-Cutting Measures list is also available for reporting.