

Proposed 2016 General Practice/Family Practice Preferred Specialty Measure Set

PQRS# CMS# GPRO#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
1 CMS122v2	0059	Claims, Registry, EHR, GPRO Web Interface/ACO, Measures Group (DM)	Effective Clinical Care	Diabetes: Hemoglobin A1c (HbA1c) Poor Control (>9%): Percentage of patients 18-75 years of age with diabetes who had hemoglobin A1c > 9.0% during the measurement period
5 CMS135v2	0081	Registry, EHR, Measures Group (HF)	Effective Clinical Care	Heart Failure: Angiotensin-Converting Enzyme (ACE) Inhibitor or Angiotensin Receptor Blocker (ARB) Therapy for Left Ventricular Systolic Dysfunction (LVSD): Percentage of patients aged 18 years and older with a diagnosis of heart failure (HF) with a current or prior left ventricular ejection fraction (LVEF) < 40% who were prescribed ACE inhibitor or ARB therapy either within a 12 month period when seen in the outpatient setting or at each hospital discharge
41	0049	Claims, Registry	Effective Clinical Care	Osteoporosis: Pharmacologic Therapy for Men and Women Aged 50 Years and Older: Percentage of patients aged 50 years and older with a diagnosis of osteoporosis who were prescribed pharmacologic therapy within 12 months
50	0100	Claims, Registry	Person and Caregiver-Centered Experience and Outcomes	Urinary Incontinence: Plan of Care for Urinary Incontinence in Women Aged 65 Years and Older: Percentage of female patients aged 65 years and older with a diagnosis of urinary incontinence with a documented plan of care for urinary incontinence at least once within 12 months
65 CMS154v2	0069	Registry, EHR	Efficiency and Cost Reduction	Appropriate Treatment for Children with Upper Respiratory Infection (URI): Percentage of children 3 months-18 years of age who were diagnosed with upper respiratory infection (URI) and were not dispensed an antibiotic prescription on or three days after the episode
66 CMS146v2	0002	Registry, EHR	Efficiency and Cost Reduction	Appropriate Testing for Children with Pharyngitis: Percentage of children 2-18 years of age who were diagnosed with pharyngitis, ordered an antibiotic and received a group A streptococcus (strep) test for the episode
93	0654	Claims, Registry, Measures Group (AOE)	Communication and Care Coordination	Acute Otitis Externa (AOE): Systemic Antimicrobial Therapy – Avoidance of Inappropriate Use: Percentage of patients aged 2 years and older with a diagnosis of AOE who were not prescribed systemic antimicrobial therapy

Proposed 2016 General Practice/Family Practice Preferred Specialty Measure Set

PQRS# CMS# GPRO#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
109	0050	Registry	Person and Caregiver-Centered Experience and Outcomes	Osteoarthritis (OA): Function and Pain Assessment: Percentage of patient visits for patients aged 21 years and older with a diagnosis of osteoarthritis (OA) with assessment for function and pain
112 CMS125v2	0031	Claims, Registry, EHR, GPRO Web Interface, Measures Group (Prev Care)	Effective Clinical Care	Breast Cancer Screening: Percentage of women 50 through 74 years of age who had a mammogram to screen for breast cancer within 27 months
113 CMS130v2	2372	Claims, Registry, EHR, GPRO Web Interface, Measures Group (Prev Care)	Effective Clinical Care	Colorectal Cancer Screening: Percentage of patients 50 through 75 years of age who had appropriate screening for colorectal cancer
116	0058	Registry	Efficiency and Cost Reduction	Avoidance of Antibiotic Treatment in Adults With Acute Bronchitis: Percentage of adults 18 through 64 years of age with a diagnosis of acute bronchitis who were not prescribed or dispensed an antibiotic prescription on or 3 days after the episode
117 CMS131v2	0055	Claims, Registry, GPRO Web Interface/ACO, EHR, Measures Group (DM)	Effective Clinical Care	Diabetes: Eye Exam: Percentage of patients 18 through 75 years of age with a diagnosis of diabetes (type 1 and type 2) who had a retinal or dilated eye exam by an eye care professional in the measurement period or a negative retinal or dilated eye exam (negative for retinopathy) in the year prior to the measurement period
154	0101	Claims, Registry, Measures Group (AOE)	Patient Safety	Falls: Risk Assessment: Percentage of patients aged 65 years and older with a history of falls who had a risk assessment for falls completed within 12 months
155	0101	Claims, Registry, Measures Group (AOE)	Communication and Care Coordination	Falls: Plan of Care: Percentage of patients aged 65 years and older with a history of falls who had a plan of care for falls documented within 12 months
163 CMS123v2	0056	EHR	Effective Clinical Care	Diabetes: Foot Exam: Percentage of patients aged 18-75 years of age with diabetes who had a foot exam during the measurement period

Proposed 2016 General Practice/Family Practice Preferred Specialty Measure Set

PQRS# CMS# GPRO#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
181	N/A	Claims, Registry	Patient Safety	Elder Maltreatment Screen and Follow-Up Plan: Percentage of patients aged 65 years and older with a documented elder maltreatment screen using an Elder Maltreatment Screening Tool on the date of encounter AND a documented follow-up plan on the date of the positive screen
204	0068	Claims, Registry, EHR, GPRO (Web Interface), Measures Group (Million Hearts)	Effective Clinical Care	Ischemic Vascular Disease (IVD): Use of Aspirin or Another Antithrombotic: Percentage of patients 18 years of age and older who were discharged alive for acute myocardial infarction (AMI), coronary artery bypass graft (CABG) or percutaneous coronary interventions (PCI) in the 12 months prior to the measurement period, or who had an active diagnosis of ischemic vascular disease (IVD) during the measurement period and who had documentation of use of aspirin or another antithrombotic during the measurement period
236	0018	Claims, Registry, EHR, GPRO (Web Interface), Measures Group (Million Hearts)	Effective Clinical Care	Controlling High Blood Pressure: Percentage of patients 18-85 years of age who had a diagnosis of hypertension and whose blood pressure was adequately controlled (<140/90 mmHg) during the measurement period
242	N/A	Measures Group (CAD)	Effective Clinical Care	Coronary Artery Disease (CAD): Symptom Management: Percentage of patients aged 18 years and older with a diagnosis of coronary artery disease seen within a 12 month period with results of an evaluation of level of activity and an assessment of whether anginal symptoms are present or absent with appropriate management of anginal symptoms within a 12 month period
317	N/A	Claims, Registry, EHR, GPRO (Web Interface), Measures Groups	Community/ Population Health	Preventive Care and Screening: Screening for High Blood Pressure and Follow-Up Documented: Percentage of patients aged 18 years and older seen during the reporting period who were screened for high blood pressure AND a recommended follow-up plan is documented based on the current blood pressure (BP) reading as indicated

Proposed 2016 General Practice/Family Practice Preferred Specialty Measure Set

PQRS# CMS# GPRO#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
326	1525	Claims, Registry	Effective Clinical Care	Atrial Fibrillation and Atrial Flutter: Chronic Anticoagulation Therapy: Percentage of patients aged 18 years and older with a diagnosis of nonvalvular atrial fibrillation (AF) or atrial flutter whose assessment of the specified thromboembolic risk factors indicate one or more high-risk factors or more than one moderate risk factor, as determined by CHADS2 risk stratification, who are prescribed warfarin OR another oral anticoagulant drug that is FDA approved for the prevention of thromboembolism
331	N/A	Registry, Measures Group (Sinusitis)	Efficiency and Cost Reduction	Adult Sinusitis: Antibiotic Prescribed for Acute Sinusitis (Appropriate Use): Percentage of patients, aged 18 years and older, with a diagnosis of acute sinusitis who were prescribed an antibiotic within 7 days of diagnosis or within 10 days after onset of symptoms
332	N/A	Registry, Measures Group (Sinusitis)	Efficiency and Cost Reduction	Adult Sinusitis: Appropriate Choice of Antibiotic: Amoxicillin Prescribed for Patients with Acute Bacterial Sinusitis (Appropriate Use): Percentage of patients aged 18 years and older with a diagnosis of acute bacterial sinusitis that were prescribed amoxicillin, with or without clavulanate, as a first line antibiotic at the time of diagnosis
333	N/A	Registry, Measures Group (Sinusitis)	Efficiency and Cost Reduction	Adult Sinusitis: Computerized Tomography (CT) for Acute Sinusitis (Overuse): Percentage of patients, aged 18 years and older, with a diagnosis of acute sinusitis who had a computerized tomography (CT) scan of the paranasal sinuses ordered at the time of diagnosis or received within 28 days after date of diagnosis
334	N/A	Registry	Efficiency and Cost Reduction	Adult Sinusitis: More than One Computerized Tomography (CT) Scan Within 90 Days for Chronic Sinusitis (Overuse): Percentage of patients aged 18 years and older with a diagnosis of chronic sinusitis who had more than one CT scan of the paranasal sinuses ordered or received within 90 days after date of diagnosis

Proposed 2016 General Practice/Family Practice Preferred Specialty Measure Set

PQRS# CMS# GPRO#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
337	N/A	Registry	Effective Clinical Care	Tuberculosis Prevention for Psoriasis, Psoriatic Arthritis and Rheumatoid Arthritis Patients on a Biological Immune Response Modifier: Percentage of patients whose providers are ensuring active tuberculosis prevention either through yearly negative standard tuberculosis screening tests or are reviewing the patient's history to determine if they have had appropriate management for a recent or prior positive test
338	2082	Measures Group (HIV/AIDS)	Effective Clinical Care	HIV Viral Load Suppression: The percentage of patients, regardless of age, with a diagnosis of HIV with a HIV viral load less than 200 copies/mL at last viral load test during the measurement year
342	0209	Registry	Person and Caregiver-Centered Experience and Outcomes	Pain Brought Under Control Within 48 Hours: Patients aged 18 and older who report being uncomfortable because of pain at the initial assessment (after admission to palliative care services) who report pain was brought to a comfortable level within 48 hours
343	N/A	Registry	Effective Clinical Care	Screening Colonoscopy Adenoma Detection Rate: The percentage of patients age 50 years or older with at least one adenoma or other colorectal cancer precursor or colorectal cancer detected during screening colonoscopy
383	1879	Registry	Patient Safety	Adherence to Antipsychotic Medications for Individuals with Schizophrenia: The percentage of individuals 18 years of age or greater as of the beginning of the measurement period with schizophrenia or schizoaffective disorder who are prescribed an antipsychotic medication, with adherence to the antipsychotic medication [defined as a Proportion of Days Covered (PDC)] of at least 0.8 during the measurement period (12 consecutive months)
387	N/A	Registry	Effective Clinical Care	Annual Hepatitis C Virus (HCV) Screening for Patients who are Active Injection Drug Users: Percentage of patients regardless of age who are active injection drug users who received a hepatitis C virus (HCV) antibody test or HCV ribonucleic acid (RNA) test within the 12 month reporting period

Proposed 2016 General Practice/Family Practice Preferred Specialty Measure Set

PQRS# CMS# GPRO#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
390	N/A	Registry, Measures Group (Hepatitis C)	Person and Caregiver-Centered Experience and Outcomes	Discussion and Shared Decision Making Surrounding Treatment Options: Percentage of patients aged 18 years and older with a diagnosis of hepatitis C with whom a physician or other clinician reviewed the range of treatment options appropriate to their genotype and demonstrated a shared decision making approach with the patient. To meet the measure, there must be documentation in the patient record of a discussion between the physician/clinician and the patient that includes all of the following: •Treatment choices appropriate to genotype •Risks and benefits •Evidence of effectiveness •Patient preferences toward the outcome of the treatment
391	0576	Registry	Communication and Care Coordination	Follow-up After Hospitalization for Mental Illness: The percentage of discharges for patients 6 years of age and older who were hospitalized for treatment of selected mental illness diagnoses and who had an outpatient visit, an intensive outpatient encounter or partial hospitalization with a mental health practitioner. Two rates are reported: - The percentage of discharges for which the patient received follow-up within 30 days of discharge - The percentage of discharges for which the patient received follow-up within 7 days of discharge
394	1407	Registry	Community/Population Health	Immunizations for Adolescents: The percentage of adolescents 13 years of age who had the recommended immunizations by their 13th birthday
398	N/A	Registry	Person and Caregiver-Centered Experience and Outcomes	Optimal Asthma Control: Patients ages 5-50 (pediatrics ages 5-17) whose asthma is well-controlled as demonstrated by one of three age appropriate patient reported outcome tools
400	N/A	Registry	Community/Population Health	Hepatitis C: One-Time Screening for Hepatitis C Virus (HCV) for Patients at Risk: Percentage of patients with one or more of the following: a history of injection drug use, receipt of a blood transfusion prior to 1992, receiving maintenance hemodialysis OR birthdate in the years 1945-1965 who received a one-time screening for HCV infection

Proposed 2016 General Practice/Family Practice Preferred Specialty Measure Set

PQRS# CMS# GPRO#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
402	N/A	Registry, Measures Group (Asthma)	Community/Population Health	Tobacco Use and Help with Quitting Among Adolescents: Percentage of adolescents 13 to 20 years of age with a primary care visit during the measurement period for whom tobacco use status was documented and received help quitting if identified as a tobacco user
Proposed New for 2016 PQRS	1880	Registry	Patient Safety	Adherence to Mood Stabilizers for Individuals with Bipolar I Disorder: The measure calculates the percentage of individuals aged 18 years and older with bipolar I disorder who are prescribed a mood stabilizer medication, with adherence to the mood stabilizer medication [defined as a Proportion of Days Covered (PDC)] of at least 0.8 during the measurement period (12 consecutive months)
Proposed New for 2016 PQRS	N/A	Registry	Effective Clinical Care	Documentation of a Health Care Proxy for Patients with Cognitive Impairment: The percentage of patients with a diagnosis of dementia or a positive result on a standardized tool for assessment of cognitive impairment, with documentation of a designated health care proxy during the measurement period
Proposed New for 2016 PQRS	N/A	Claims, Registry	Effective Clinical Care	HIV Screening of STI patients: Percentage of patients diagnosed with an acute STI who were tested for HIV
Proposed New for 2016 PQRS	N/A	Claims, Registry	Effective Clinical Care	Statin Therapy for the Prevention and Treatment of Cardiovascular Disease: Percentage of high-risk adult patients aged ≥ 21 years who were previously diagnosed with or currently have an active diagnosis of clinical atherosclerotic cardiovascular disease (ASCVD); OR adult patients aged ≥ 21 years with a fasting or direct Low-Density Lipoprotein Cholesterol (LDL-C) level ≥ 190 mg/dL; OR patients aged 40-75 years with a diagnosis of diabetes with a fasting or direct Low-Density Lipoprotein Cholesterol (LDL-C) level of 70-189 mg/dL; who were prescribed or are already on statin medication therapy during the measurement period.
Proposed New for 2016 PQRS	0053	Claims, Registry	Effective Clinical Care	Osteoporosis Management in Women Who Had a Fracture: The percentage of women age 50-85 who suffered a fracture and who either had a bone mineral density test or received a prescription for a drug to treat osteoporosis

This table is a resource of potential measures that may be applicable to eligible professionals providing clinical services for General Practice/Family Practice. Please note that in addition to this table, a 2016 Proposed Cross-Cutting Measures list is also available for reporting.