

Proposed 2016 Urology Preferred Specialty Measure Set

| PQRS# CMS# GPRO# | NQF# | Reporting Method | National Quality Strategy Domain | Measure Title: Description48 |
|------------------------|------|---|---|--|
| 48 | N/A | Claims, Registry, Measure Groups (Prev Care) | Effective Clinical Care | Urinary Incontinence: Assessment of Presence or Absence of Urinary Incontinence in Women Aged 65 Years and Older: Percentage of female patients aged 65 years and older who were assessed for the presence or absence of urinary incontinence within 12 months |
| 50 | N/A | Claims, Registry | Person and Caregiver-Centered Experience and Outcomes | Urinary Incontinence: Plan of Care for Urinary Incontinence in Women Aged 65 Years and Older: Percentage of female patients aged 65 years and older with a diagnosis of urinary incontinence with a documented plan of care for urinary incontinence at least once within 12 months |
| 102 | 0389 | Registry, EHR | Efficiency and Cost Reduction | Prostate Cancer: Avoidance of Overuse of Bone Scan for Staging Low Risk Prostate Cancer Patients: Percentage of patients, regardless of age, with a diagnosis of prostate cancer at low risk of recurrence receiving interstitial prostate brachytherapy, OR external beam radiotherapy to the prostate, OR radical prostatectomy, OR cryotherapy who did not have a bone scan performed at any time since diagnosis of prostate cancer |
| 104 | 0390 | Registry | Effective Clinical Care | Prostate Cancer: Adjuvant Hormonal Therapy for High Risk Prostate Cancer Patients: Percentage of patients, regardless of age, with a diagnosis of prostate cancer at high or very high risk of recurrence receiving external beam radiotherapy to the prostate who were prescribed adjuvant hormonal therapy (GnRH [gonadotropin-releasing hormone] agonist or antagonist) |
| 119 | 0062 | Claims, Registry, EHR, Measure Groups (Diabetes Mellitus) | Effective Clinical Care | Diabetes: Medical Attention for Nephropathy: The percentage of patients 18-75 years of age with diabetes who had a nephropathy screening test or evidence of nephropathy during the measurement period |
| 173 | N/A | Registry, Measures Group (Prev Care) | Community/ Population Health | Preventive Care and Screening: Unhealthy Alcohol Use – Screening: Percentage of patients aged 18 years and older who were screened for unhealthy alcohol use at least once within 24 months using a systematic screening method |

Proposed 2016 Urology Preferred Specialty Measure Set

| PQRS# CMS# GPRO# | NQF# | Reporting Method | National Quality Strategy Domain | Measure Title: Description ⁴⁸ |
|------------------------|------|--|---|--|
| 194 | 0386 | Registry | Effective Clinical Care | Oncology: Cancer Stage Documented: Percentage of patients, regardless of age, with a diagnosis of cancer who are seen in the ambulatory setting who have a baseline American Joint Committee on Cancer (AJCC) cancer stage or documentation that the cancer is metastatic in the medical record at least once during the 12 month reporting period |
| 265 | N/A | Registry | Communication and Care Coordination | Biopsy Follow-Up: Percentage of new patients whose biopsy results have been reviewed and communicated to the primary care/referring physician and patient by the performing physician |
| 358 | N/A | Registry, Measure Groups (General Surgery) | Person and Caregiver-Centered Experience and Outcomes | Patient-Centered Surgical Risk Assessment and Communication: Percentage of patients who underwent a non-emergency surgery who had their personalized risks of postoperative complications assessed by their surgical team prior to surgery using a clinical data-based, patient-specific risk calculator and who received personal discussion of those risks with the surgeon |

This table is a resource of potential measures that may be applicable to eligible professionals providing clinical services for Urology. Please note that in addition to this table, a 2016 Proposed Cross-Cutting Measures list is also available for reporting.