

Potential OB/GYN Preferred Specialty Measure Set

PQRS# CMS# GPRO#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
39	0046	Claims, Registry Measures Group (Prev Care)	Effective Clinical Care	Screening or Therapy for Osteoporosis for Women Aged 65 Years and Older. Percentage of female patients aged 65 years and older who have a central dual-energy X- ray absorptiometry (DXA) measurement ordered or performed at least once since age 60 or pharmacologic therapy prescribed within 12 months
40	0048	Claims, Registry	Effective Clinical Care	Osteoporosis: Management Following Fracture of Hip, Spine or Distal Radius for Men and Women Aged 50 Years and Older. Percentage of patients aged 50 years and older with fracture of the hip, spine, or distal radius who had a central dual-energy X-ray absorptiometry (DXA) measurement ordered or performed or pharmacologic therapy prescribed
41	N/A	Claims, Registry	Effective Clinical Care	Osteoporosis: Pharmacologic Therapy for Men and Women Aged 50 Years and Older. Percentage of patients aged 50 years and older with a diagnosis of osteoporosis who were prescribed pharmacologic therapy within 12 months
48	N/A	Claims, Registry Measures Group (Prev Care)	Effective Clinical Care	Urinary Incontinence: Assessment of Presence or Absence of Urinary Incontinence in Women Aged 65 Years and Older. Percentage of female patients aged 65 years and older who were assessed for the presence or absence of urinary incontinence within 12 months
50	N/A	Claims, Registry	Person and Caregiver-Centered Experience and Outcomes	Urinary Incontinence: Plan of Care for Urinary Incontinence in Women Aged 65 Years and Older. Percentage of female patients aged 65 years and older with a diagnosis of urinary incontinence with a documented plan of care for urinary incontinence at least once within 12 months
112 CMS125v2	0031	Claims, Registry, EHR, GPRO Web Interface, Measures Group (Prev Care)	Effective Clinical Care	Breast Cancer Screening. Percentage of women 50 through 74 years of age who had a mammogram to screen for breast cancer within 27 months
265	N/A	Registry	Communication and Care Coordination	Biopsy Follow-Up. Percentage of patients whose biopsy results have been reviewed and communicated to the primary care/referring physician and patient by the performing physician
309 CMS124v2	0032	EHR	Effective Clinical Care	Cervical Cancer Screening: Percentage of women 21-64 years of age, who received one or more Pap tests to screen for cervical cancer

Potential OB/GYN Preferred Specialty Measure Set

PQRS# CMS# GPRO#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
310 CMS153	0033	EHR	Community/Population Health	Chlamydia Screening: Percentage of women 16-24 years of age who were identified as sexually active and who had at least one test for chlamydia during the measurement period
335	N/A	Registry	Patient Safety	Maternity Care: Elective Delivery or Early Induction Without Medical Indication at ≥ 37 and < 39 Weeks. Percentage of patients, regardless of age, who gave birth during a 12-month period who delivered a live singleton at ≥ 37 and < 39 weeks of gestation completed who had elective deliveries or early inductions without medical indication
336	N/A	Registry	Communication and Care Coordination	Maternity Care: Post-Partum Follow-Up and Care Coordination. Percentage of patients, regardless of age, who gave birth during a 12-month period who were seen for post-partum care within 8 weeks of giving birth who received a breast feeding evaluation and education, post-partum depression screening, post-partum glucose screening for gestational diabetes patients, and family and contraceptive planning

This table is a resource of potential measures that may be applicable to OB/GYN.