

## 2016 OB/GYN Preferred Specialty Measure Set

| PQRS# | NQF# | Reporting Method   | National Quality Strategy Domain                      | Measure Title: Description   |
|-------|------|--|---|--|
| 39    | 0046 | Claims, Registry Measures Groups                             | Effective Clinical Care                               | <b>Screening for Osteoporosis for Women Aged 65-85 Years of Age:</b> Percentage of female patients aged 65-85 years of age who ever had a central dual-energy X-ray absorptiometry (DXA) to check for osteoporosis   |
| 41    | N/A  | Claims, Registry   | Effective Clinical Care                               | <b>Osteoporosis: Pharmacologic Therapy for Men and Women Aged 50 Years and Older:</b> Percentage of patients aged 50 years and older with a diagnosis of osteoporosis who were prescribed pharmacologic therapy within 12 months   |
| 48    | N/A  | Claims, Registry Measures Groups                             | Effective Clinical Care                               | <b>Urinary Incontinence: Assessment of Presence or Absence of Urinary Incontinence in Women Aged 65 Years and Older:</b> Percentage of female patients aged 65 years and older who were assessed for the presence or absence of urinary incontinence within 12 months  |
| 50    | N/A  | Claims, Registry   | Person and Caregiver-Centered Experience and Outcomes | <b>Urinary Incontinence: Plan of Care for Urinary Incontinence in Women Aged 65 Years and Older:</b> Percentage of female patients aged 65 years and older with a diagnosis of urinary incontinence with a documented plan of care for urinary incontinence at least once within 12 months   |
| 112   | 2372 | Claims, Registry, EHR, GPRO (Web Interface), Measures Groups | Effective Clinical Care                               | <b>Breast Cancer Screening:</b> Percentage of women 50 through 74 years of age who had a mammogram to screen for breast cancer within 27 months  |
| 204   | 0068 | Claims, Registry, EHR, GPRO (Web Interface), Measures Groups | Effective Clinical Care                               | <b>Ischemic Vascular Disease (IVD): Use of Aspirin or Another Antithrombotic:</b> Percentage of patients 18 years of age and older who were discharged alive for acute myocardial infarction (AMI), coronary artery bypass graft (CABG) or percutaneous coronary interventions (PCI) in the 12 months prior to the measurement period, or who had an active diagnosis of ischemic vascular disease (IVD) during the measurement period and who had documentation of use of aspirin or another antithrombotic during the measurement period |

## 2016 OB/GYN Preferred Specialty Measure Set

| PQRS# | NQF# | Reporting Method   | National Quality Strategy Domain    | Measure Title: Description   |
|-------|------|--|-------------------------------------|--|
| 226   | 0028 | Claims, Registry, EHR, GPRO (Web Interface), Measures Groups | Community/Population Health         | <b>Preventive Care and Screening: Tobacco Use: Screening and Cessation</b><br><b>Intervention:</b> Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 24 months AND who received cessation counseling intervention if identified as a tobacco user                                |
| 236   | 0018 | Claims, Registry, EHR, GPRO (Web Interface), Measures Groups | Effective Clinical Care             | <b>Controlling High Blood Pressure:</b> Percentage of patients 18-85 years of age who had a diagnosis of hypertension and whose blood pressure was adequately controlled (<140/90 mmHg) during the measurement period  |
| 265   | N/A  | Registry   | Communication and Care Coordination | <b>Biopsy Follow-Up:</b> Percentage of patients whose biopsy results have been reviewed and communicated to the primary care/referring physician and patient by the performing physician   |
| 309   | 0032 | EHR  | Effective Clinical Care             | <b>Cervical Cancer Screening:</b> Percentage of women 21-64 years of age, who received one or more Pap tests to screen for cervical cancer   |
| 310   | 0033 | EHR  | Community/Population Health         | <b>Chlamydia Screening for Women:</b> Percentage of women 16-24 years of age who were identified as sexually active and who had at least one test for chlamydia during the measurement period  |
| 317   | N/A  | Claims, Registry, EHR, GPRO (Web Interface), Measures Groups | Community/ Population Health        | <b>Preventive Care and Screening: Screening for High Blood Pressure and Follow-Up Documented:</b> Percentage of patients aged 18 years and older seen during the reporting period who were screened for high blood pressure AND a recommended follow-up plan is documented based on the current blood pressure (BP) reading as indicated |
| 418   | 0053 | Claims, Registry   | Effective Clinical Care             | <b>Osteoporosis Management in Women Who Had a Fracture:</b> The percentage of women age 50-85 who suffered a fracture and who either had a bone mineral density test or received a prescription for a drug to treat osteoporosis   |
| 422   | 2063 | Claims, Registry   | Patient Safety                      | <b>Performing Cystoscopy at the time of Hysterectomy for Pelvic Organ Prolapse to Detect Lower Urinary Tract Injury:</b> Percentage of patients who undergo cystoscopy to evaluate for lower urinary tract injury at the time of hysterectomy for pelvic organ prolapse  |

## 2016 OB/GYN Preferred Specialty Measure Set

| PQRS# | NQF# | Reporting Method | National Quality Strategy Domain | Measure Title: Description   |
|-------|------|------------------|----------------------------------|--|
| 432   | N/A  | Registry         | Patient Safety                   | <b>Proportion of Patients Sustaining a Bladder Injury at the Time of any Pelvic Organ Prolapse Repair:</b> Percentage of patients undergoing any surgery to repair pelvic organ prolapse who sustains an injury to the bladder recognized either during or within 1 month after surgery  |
| 433   | N/A  | Registry         | Patient Safety                   | <b>Proportion of Patients Sustaining a Major Viscus Injury at the Time of any Pelvic Organ Prolapse Repair:</b> Percentage of patients undergoing surgical repair of pelvic organ prolapse that is complicated by perforation of a major viscus at the time of index surgery that is recognized intraoperative or within 1 month after surgery |
| 434   | N/A  | Registry         | Patient Safety                   | <b>Proportion of Patients Sustaining A Ureter Injury at the Time of any Pelvic Organ Prolapse Repair:</b> Percentage of patients undergoing a pelvic organ prolapse repair who sustain an injury to the ureter recognized either during or within 1 month after surgery  |

This table is a resource of potential measures that may be applicable to OB/GYN. Please note that in addition to this table, a cross cutting measures list is also available for reporting at the following link: [2016 Cross Cutting Measures List](#)