

From: Clinician Outreach and Communication Activity (CDC)
Sent: Tuesday, December 15, 2009 10:46 AM
To: Clinician Outreach and Communication Activity (CDC)
Subject: CDC Health Update - Non-Safety Related Voluntary Recall of Certain Lots of Sanofi Pasteur H1N1 Pediatric Vaccine



This is an official CDC Health Update

Distributed via Health Alert Network
December 15, 2009, 10:04 EST (10:04 AM EST)
CDCHAN-00303-09-12-15-ADV-N

Non-Safety Related Voluntary Recall of Certain Lots of Sanofi Pasteur H1N1 Pediatric (0.25 mL, for 6-35 month olds) Vaccine in Pre-Filled Syringes

Summary

As part of its quality assurance program, Sanofi Pasteur, Inc., performs additional routine, ongoing testing of influenza vaccines after the vaccine has been distributed to health care providers to ensure that vaccines continue to meet required specifications. In recent testing of the amount of antigen in its influenza A (H1N1) monovalent vaccine, Sanofi Pasteur found four distributed lots of single-dose, pre-filled syringe pediatric (0.25 mL.) vaccine with antigen content lower than required potency levels. The manufacturer is conducting a non-safety related voluntary recall of these affected lots of vaccine.

Background

After performing these tests, Sanofi Pasteur notified the Centers for Disease Control and Prevention (CDC) and the Food and Drug Administration (FDA) that the antigen content in one lot of pediatric syringes that had been distributed to providers was later found to have dropped below a pre-specified limit. As a result of this finding, Sanofi Pasteur tested additional lots and found that three other lots that had been distributed also had an antigen content that had fallen below pre-specified limits. This means that doses from these four vaccine lots no longer meet the specifications for antigen content.

Recommendations

While the antigen content of these lots is now below the specification limit for the product, CDC and FDA are in agreement that the small decrease in antigen content is unlikely to result in a clinically significant reduction in immune response among persons who have received the vaccine. For this reason, there is no need to revaccinate persons who have received vaccine from these lots.

Providers are being asked to return any vaccine to the manufacturer in the following lots that remains unused to the manufacturer:

- 0.25 mL pre-filled syringes, 10-packs (NDC # 49281-650-25, sometimes coded as 49281-0650-25):
UT023DA
UT028DA
UT028CB
- 0.25 mL pre-filled syringes, 25-packs (NDC # 49281-650-70, sometimes coded as 49281-0650-70):

UT030CA

These lots were shipped in November and are intended for children 6 months through 35 months of age. Sanofi Pasteur will send directions for returning unused vaccine from these lots to providers.

All vaccines are thoroughly tested prior to release and shipping to determine that they meet all manufacturer and FDA standards for purity, potency and safety. The affected vaccine met all specifications at the time of release. CDC and FDA have determined that there are no safety concerns for children who have received this vaccine. Sanofi Pasteur has discontinued distribution of the 0.25 mL syringes of H1N1 pediatric vaccines.

The drop in antigen content below the required specification that is described here is specific to Sanofi Pasteur's pediatric H1N1 monovalent vaccine in 0.25 mL pre-filled syringes. The same vaccine packaged in other forms, such as 0.5 mL pre-filled syringes for older children and adults and multi-dose vials, continue to meet specifications.

The antigen content in the affected lots of vaccine is only slightly below the specification limit. The slightly reduced concentration of vaccine antigen found in retesting these lots is still expected to be effective in stimulating a protective response. There is no need to re-administer a dose to those who received vaccine from these lots. However, as is recommended for all 2009 H1N1 vaccines, all children less than 10 years old should get the recommended two doses of H1N1 vaccine approximately a month apart for the optimal immune response. So, children less than 10 years old who have only received one dose of vaccine thus far should still receive a second dose of 2009 H1N1 vaccine.

For children 6 months of age and older, vaccine is available in multidose vials. The vaccine in multidose vials is safe and effective vaccine for children. One difference between vaccine in pre-filled syringes and the multidose vials is that the multidose vials contain a preservative (thimerosal) to prevent potential contamination after the vial is opened. The standard dose for this preparation for administration to infants 6-35 months old is the same as for the pre-filled syringes, 0.25 mL. For healthy children at least 2 years of age, the nasal spray (live, attenuated influenza vaccine) is also an option. The nasal spray vaccine is produced in single units that do not contain thimerosal.

For More Information:

- For Questions and Answers related to the withdrawn vaccine see http://www.cdc.gov/h1n1flu/vaccination/syringes_qa.htm
- Call CDC's toll-free information line, 800-CDC-INFO (800-232-4636) TTY: (888) 232-6348, which is available 24 hours a day, every day.

Categories of Health Alert messages:

Health Alert conveys the highest level of importance; warrants immediate action or attention.

Health Advisory provides important information for a specific incident or situation; may not require immediate action.

Health Update provides updated information regarding an incident or situation; unlikely to require immediate action.

##This Message was distributed to **State and Local Health Officers, Public Information Officers, Epidemiologists and HAN Coordinators as well as Clinician organizations##**

You have received this message based upon the information contained within our emergency notification data base. If you have a different or additional e-mail or fax address that you would like us to use please contact your State-based Health Alert Network program at your State or local health department.

The CDC and HHS logos are the exclusive property of the Department of Health and Human Services and may not be used for any purpose without prior express written permission. Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.


Links to non-Federal organizations are provided solely as a service to our users. Links do not constitute an endorsement of any organization by CDC or the Federal Government, and none should be inferred. The CDC is not responsible for the content of the individual organizations.

Please send us your feedback or comments: <http://www.cdc.gov/flu/coca/feedback.htm>.

Our Clinician Communication Team is committed to excellence in reporting our weekly updates. Please e-mail coca@cdc.gov should you note any written errors or discrepancies.

If you need to unsubscribe or update your information, please go to our website: <http://www.bt.cdc.gov/clinregistry>.

If you need further information or technical help, please send an e-mail message to: coca@cdc.gov.



Communicating With Clinicians

CDC currently employs several mechanisms to communicate with clinicians on topics of urgent concern and interest.

[Download this flyer to find out more](#)



