

ADVISORY PANEL ON CLINICAL DIAGNOSTIC LABORATORY TESTS (CDLT)

Recommendations

July 18, 2016

New Code/ Reconsideration Request	Code Description	CDLT Panel Voting
803X1X	Drug test(s), presumptive, any number of drug classes, qualitative, any number of devices or procedures, (eg, immunoassay) capable of being read by direct optical observation only (eg, dipsticks, cups, cards, cartridges) includes sample validation when performed, per date of service.	11: Crosswalk to G0477 1: Crosswalk to unspecified code 1: Recuse 0: Gapfill
803X2X	Drug test(s), presumptive, any number of drug classes, qualitative, any number of devices or procedures, (eg, immunoassay) read by instrument assisted direct optical observation (eg, dipsticks, cups, cards, cartridges), includes sample validation when performed, per date of service.	11: Crosswalk to G0478 1: Crosswalk to unspecified code 1: Recuse 0: Gapfill
803X3X	Drug test(s), presumptive, any number of drug classes, qualitative, any number of devices or procedures by instrument chemistry analyzers (eg, utilizing immunoassay [eg, EIA, ELISA, EMIT, FPIA, IA, KIMS, RIA]), chromatography (eg, GC, HPLC), and mass spectrometry either with or without chromatography, (eg, DART, DESI, GC-MS, GC-MS/MS, LC-MS, LC-MS/MS, LDTD, MALDI, TOF) includes sample validation when performed, per date of service.	11: Crosswalk to G0479 1: Crosswalk to unspecified code 1: Recuse 0: Gapfill
813X7X	SEPT9 (Septin9) (eg, colorectal cancer) methylation analysis.	9: Crosswalk to 81288 3: Gapfill 1: Recuse
814X5X	Cardiac ion channelopathies (eg, Brugada syndrome, long QT syndrome, short QT syndrome, catecholaminergic polymorphic ventricular tachycardia); genomic sequence analysis panel, must include sequencing of at least 10 genes, including ANK2, CASQ2, CAV3, KCNE1, KCNE2, KCNH2, KCNJ2, KCNQ1, RYR2, and SCN5A	10: Crosswalk to 81435 1: Crosswalk to unspecified code 1: Recuse 1: Gapfill
814X6X	Cardiac ion channelopathies (eg, Brugada syndrome, long QT syndrome, short QT syndrome, catecholaminergic polymorphic ventricular tachycardia); duplication/deletion gene analysis panel, must include analysis of at least 2 genes, including KCNH2 and KCNQ1.	6: Crosswalk to 81436 5: Gapfill 1: Crosswalk to 0.5 X 81436 1: Recuse

ADVISORY PANEL ON CLINICAL DIAGNOSTIC LABORATORY TESTS (CDLT)

Recommendations

July 18, 2016

New Code/ Reconsideration Request	Code Description	CDLT Panel Voting
814X3X	Fetal chromosomal microdeletion(s) genomic sequence analysis (eg, DiGeorge syndrome, Cri-du-chat syndrome), circulating cell-free fetal DNA in maternal blood.	8: Gapfill 4: Crosswalk to 81435 1: Recuse
814X2X	Inherited cardiomyopathy (eg, hypertrophic cardiomyopathy, dilated cardiomyopathy, arrhythmogenic right ventricular cardiomyopathy) genomic sequence analysis panel, must include sequencing of at least 5 genes, including DSG2, MYBPC3, MYH7, PKP2, and TTN.	7: Crosswalk to 81435 5: Gapfill 1: Recuse
814X1X	Oncology (high-grade prostate cancer), biochemical assay of four proteins (Total PSA, Free PSA, Intact PSA and human kallikrein-2 [hK2]), utilizing plasma or serum, prognostic algorithm reported as a probability score.	5: Crosswalk to 0010M; if not feasible, then Gapfill 5: Crosswalk to 3 X 84153 + 84154 3: Gapfill
844XXX	Testosterone; bioavailable, direct measurement (eg, differential precipitation).	11: Crosswalk to 84402 + 84403 1: Crosswalk to 3 X 84403 1: Recuse
878XXX	Infectious agent detection by nucleic acid (DNA or RNA); central nervous system pathogen (eg, Neisseria meningitidis, Streptococcus pneumoniae, Listeria, Haemophilus influenzae, E. coli, Streptococcus agalactiae, enterovirus, human parechovirus, herpes simplex virus type 1 and 2, human herpes virus 6, cytomegalovirus, varicella zoster virus, Cryptococcus), includes multiplex reverse transcription, when performed, and multiplex amplified probe technique, multiple types or subtypes, 12-25 targets.	11: Crosswalk to 87633 1: Gapfill 1: Recuse
G0475	Hiv antigen/antibody, combination assay, screening.	11: Crosswalk to 87389 1: Crosswalk to 87806 1: Recuse
G0476	Infectious agent detection by nucleic acid (dna or rna); human papillomavirus (hpv), high-risk types (eg, 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 68) for cervical cancer screening, must be performed in addition to pap test.	10: Crosswalk to 87624 2: Gapfill 1: Recuse

ADVISORY PANEL ON CLINICAL DIAGNOSTIC LABORATORY TESTS (CDLT)

Recommendations

July 18, 2016

New Code/ Reconsideration Request	Code Description	CDLT Panel Voting
G0480 - RECONSIDERATION REQUEST	Drug test(s), definitive, utilizing drug identification methods able to identify individual drugs and distinguish between structural isomers (but not necessarily stereoisomers), including, but not limited to GC/MS (any type, single or tandem) and LC/MS (any type, single or tandem and excluding immunoassays (eg, IA, EIA, ELISA, EMIT, FPIA) and enzymatic methods (eg, alcohol dehydrogenase)); qualitative or quantitative, all sources, includes specimen validity testing, per day, 1-7 drug class(es), including metabolite(s) if performed.	8: Crosswalk to 6 X 82542 1: Crosswalk to 4.5 X 82542 1: Crosswalk to 82542 + [5 X 0.6 X 82542] 1: Gapfill 1: Current Price 1: Abstain
G0481 - RECONSIDERATION REQUEST	Drug test(s), definitive, utilizing drug identification methods able to identify individual drugs and distinguish between structural isomers (but not necessarily stereoisomers), including, but not limited to GC/MS (any type, single or tandem) and LC/MS (any type, single or tandem and excluding immunoassays (eg, IA, EIA, ELISA, EMIT, FPIA) and enzymatic methods (eg, alcohol dehydrogenase)); qualitative or quantitative, all sources, includes specimen validity testing, per day, 8-14 drug class(es), including metabolite(s) if performed.	10: Crosswalk to 8 X 82542 1: Crosswalk to 82542 + [12 X 0.6 X 82542] 1: Gapfill 1: Abstain
G0482 - RECONSIDERATION REQUEST	Drug test(s), definitive, utilizing drug identification methods able to identify individual drugs and distinguish between structural isomers (but not necessarily stereoisomers), including, but not limited to GC/MS (any type, single or tandem) and LC/MS (any type, single or tandem and excluding immunoassays (eg, IA, EIA, ELISA, EMIT, FPIA) and enzymatic methods (eg, alcohol dehydrogenase)); qualitative or quantitative, all sources, includes specimen validity testing, per day, 15-21 drug class(es), including metabolite(s) if performed.	10: Crosswalk to 10 X 82542 1: Crosswalk to 82542 + [19 X 0.6 X 82542] 1: Gapfill 1: Abstain

ADVISORY PANEL ON CLINICAL DIAGNOSTIC LABORATORY TESTS (CDLT)

Recommendations

July 18, 2016

New Code/ Reconsideration Request	Code Description	CDLT Panel Voting
G0483 - RECONSIDERATION REQUEST	Drug test(s), definitive, utilizing drug identification methods able to identify individual drugs and distinguish between structural isomers (but not necessarily stereoisomers), including, but not limited to GC/MS (any type, single or tandem) and LC/MS (any type, single or tandem and excluding immunoassays (eg, IA, EIA, ELISA, EMIT, FPIA) and enzymatic methods (eg, alcohol dehydrogenase)); qualitative or quantitative, all sources, includes specimen validity testing, per day, 22 or more drug class(es), including metabolite(s) if performed.	10: Crosswalk to 12 X 82542 1: Crosswalk to 82542 + [27 X 0.6 X 82542] 1: Gapfill 1: Abstain