

Medtronic

CRHF Economics & Reimbursement
8200 Coral Sea Street NE, MVS33
Mounds View, MN 55112
USA
www.medtronic.com

February 21, 2017

Ms. Tamara Syrek Jensen, J.D.
Director, Coverage & Analysis Group
Center for Clinical Standards and Quality
Centers for Medicare & Medicaid Services
7500 Security Boulevard, C1-14-15
Baltimore, Maryland 21244

Dear Ms. Syrek Jensen:

Medtronic is the world's leading medical technology company, specializing in implantable and interventional therapies that alleviate pain, restore health, and extend life. We are committed to the continual research and development necessary to produce high-quality products and to support innovative therapies that improve patients' health outcomes.

We appreciate the opportunity to comment on the Medicare Evidence Development & Coverage Advisory Committee (MEDCAC) meeting on heart failure. Medtronic appreciates CMS's interest in the relevant outcomes in studies for heart failure treatment technologies.

Medtronic has provided comments in response to the specific questions and discussion points that CMS posted in the MEDCAC Meeting notice. CMS specifically expressed "... medical technologies are receiving market authorization based on less long-term data with greater reliance upon intermediate and surrogate outcomes. As a result, assessments of medical technologies are being made with more frequent evidence gaps with respect to the clinically meaningful health outcomes for CMS beneficiaries."

Heart failure clinical trials with the traditional standard endpoints of mortality and morbidity are increasingly challenging to complete as event rates decline with the evolution of care and as we move toward studying healthier heart failure populations. These trials are exceedingly less feasible to conduct in a time and cost efficient manner. Mortality and morbidity as continued standards therefore, may delay or potentially reduce the availability of important heart failure innovations to patients who could benefit. We believe the aims of establishing safety and effectiveness of a therapy can confidently be assessed beyond traditional mortality and morbidity studies such that important innovations can be delivered in a timely manner to patients in need.

Medtronic presents consideration that there are important and reliable surrogate outcomes that should be considered as appropriate outcomes for heart failure treatment technologies. We highlight two viable surrogate outcomes to be considered.

Clinical Composite Score (CCS)

The Packer CCS was introduced in 2001 as a means to measure a HF intervention's intermediate effect on outcomes and symptoms. The CCS leverages clinically meaningful objective measures of death and HFH and subjective measures of NYHA functional class and global assessment. Importantly, CCS accounts for discontinuation of therapy due to clinical deterioration. The decision flow to identify whether the patient's status "improved," "worsened," or is "unchanged" is decided by a hierarchical flow, placing priority on death, HFH, crossovers, then subjective measures. Patient well-being is only relevant in the absence of outcomes or worsening NYHA class. Unlike other endpoints, every patient contributes to the analysis through the tested duration.

CCS has been used in notable trials as a primary endpoint^{i,ii,iii} and a key secondary endpoint^{iv v,vi,vii} and continues to be used^{viii,ix}. The commonality of the endpoint enables comparative effectiveness research against new products^x. A 2014 analysis of 1603 CRT patients from five trials shows six month CCS status to be predictive of long-term survival and healthcare utilization^{xi}.

Using CCS is likely to have a dramatic impact on the duration and cost efficiency of trial execution. In a review of nine pivotal CRT trials, the trials using CCS as a key endpoint were substantially shorter in duration (2.9 vs 4.7 yrs.) and 91% reduced in the patient-years sample compared to morbidity/mortality trials. Despite the smaller sample and shorter duration, to date we are unaware of an intervention showing a significant effect in CCS but not in mortality and morbidity outcomes (i.e. false positive). While REVERSE reported an insignificant CCS effect of CRT in select NYHA I-II patients at 12 months, the pre-specified endpoint only evaluated worsening vs non-worsening. When CCS was used conventionally, accounting also for unchanged and improved statuses, the effect was significant and at multiple time-points^{xii}. CRT for mild HF would further prove effective in the reduction of outcomes one and two years later when reported by the MADIT CRT^{xiii} and RAFT^{xiv} trials, respectively.

There are some limitations to CCS worth noting. CCS may not detect a favorable effect after early worsening. However, this limitation also resides in other outcome measures. CCS was built as an intermediate measure, and its sensitivity at measuring long-term effect is unknown. Further, components of the CCS can be influenced by bias thus patient and clinician blinding may be needed.

Myocardial Remodeling

Myocardial remodeling has an important relationship to HF as injury to the myocardium can result in adverse ventricular remodeling, impairing contractile function, and reducing stroke volume. Remodeling increases patient risk of HF symptoms, morbidity, and mortality. HF drug therapy^{xv} and CRT^{xvi,xvii ,xviii,xix,xx,xxi} have been demonstrated to limit or reverse LV remodeling and improve long-term survival. LVESVi was a prospectively powered secondary endpoint in the REVERSE trial.

Thus far, although limited to post-hoc analyses, LVESV has been shown to be the strongest echo measure predictive of survival^{xxii,xxiii,xxiv}. Change in myocardial function as a predictor of long-term outcomes in the context of a new HF intervention is biologically plausible and an objective measurement unlikely to be influenced by patient or clinician bias. This may be particularly appealing when blinded randomization coupled with long-term follow-up is unfeasible^{xxv}. We estimate that half of the qualifying patients for the MIRACLE EF study declined participation due to the chance of having a sham device implant for five years. Systematic measurement of cardiac volumes allows for specific detection of improvement and worsening of HF function.

Identifying echocardiographic measures to predict patient response to CRT has been challenging, as poor agreement to clinical measures has been reported^{xxvi}. Interpretation of results can be difficult for modest but statistically significant degrees of functional change. Additionally, linking remodeling to economic measures has yet to be established and further research is warranted. Remodeling measures, as a key intermediate endpoint^{xxvii} or as a component of long-term measures^{xxviii} has been useful and may continue to be, particularly when the intervention risks are well-known. For example, due to the established safety profile and robust evidence of CRT's effects on volumes in systolic dysfunction (i.e. $EF \leq 35\%$), volumes as a primary endpoint for patients with ejection fractions above 35% may be justified.

Thank you for your consideration of these recommendations. We look forward to the MEDCAC Meeting on Heart Failure outcome measures. Please do not hesitate to contact me about the information we have provided.

Sincerely,



Dan Schaber, PharmD
Vice President, Heart Failure Clinical Research
Medtronic, plc
Cardiac Rhythm and Heart Failure

References

- ⁱ Linde C, Abraham WT, Gold MR, St John Sutton M, Ghio S, Daubert C; REVERSE (REsynchronization reVERses Remodeling in Systolic left vEntricular dysfunction) Study Group. Randomized trial of cardiac resynchronization in mildly symptomatic heart failure patients and in asymptomatic patients with left ventricular dysfunction and previous heart failure symptoms. *J Am Coll Cardiol*. 2008;52(23):1834-43.
- ⁱⁱ Hindricks G, Tabor sky M, Glikson M, Heinrich U, Schumacher B, Katz A, Brachmann J, Lewalter T, Goette A, Block M, Kautzner J, Sack S, Husser D, Piorkowski C, Sogaard P; IN-TIME study group. *Lancet*. 2014;384(9943):583-90.
- ⁱⁱⁱ Martin DO, Day JD, Lai PY, Murphy AL, Nayak HM, Villareal RP, Weiner S, Kraus SM, Stolen KQ, Gold MR. Atrial support pacing in heart failure: results from the multicenter PEGASUS CRT trial. *J Cardiovasc Electrophysiol*. 2012;23(12):1317-25.
- ^{iv} Abraham WT, Fisher WG, Smith AL, Delurgio DB, Leon AR, Loh E, Kocovic DZ, Packer M, Clavell AL, Hayes DL, Ellestad M, Trupp RJ, Underwood J, Pickering F, Truex C, McAtee P, Messenger J; MIRACLE Study Group. Cardiac resynchronization in chronic heart failure. *N Engl J Med*. 2002;346(24):1845-53.
- ^v Young JB, Abraham WT, Smith AL, Leon AR, Lieberman R, Wilkoff B, Canby RC, Schroeder JS, Liem LB, Hall S, Wheelan K; Multicenter InSync ICD Randomized Clinical Evaluation (MIRACLE ICD) Trial Investigators. Combined cardiac resynchronization and implantable cardioversion defibrillation in advanced chronic heart failure: the MIRACLE ICD Trial. *JAMA*. 2003;289(20):2685-94.
- ^{vi} Curtis AB, Worley SJ, Adamson PB, Chung ES, Niazi I, Sherfese L, Shinn T, Sutton MS; Biventricular versus Right Ventricular Pacing in Heart Failure Patients with Atrioventricular Block (BLOCK HF) Trial Investigators. Biventricular pacing for atrioventricular block and systolic dysfunction. *N Engl J Med*. 2013;368(17):1585-93.
- ^{vii} Torre-Amione G, Anker SD, Bourge RC, Colucci WS, Greenberg BH, Hildebrandt P, Keren A, Motro M, Moyé LA, Otterstad JE, Pratt CM, Ponikowski P, Rouleau JL, Sestier F, Winkelmann BR, Young JB; Advanced Chronic Heart Failure CLinical Assessment of Immune Modulation Therapy Investigators. Results of a non-specific immunomodulation therapy in chronic heart failure (ACCLAIM trial): a placebo-controlled randomised trial. *Lancet*. 2008;371(9608):228-36.
- ^{viii} McMurray JJ, Packer M, Desai AS, Gong J, Lefkowitz MP, Rizkala AR, Rouleau J, et al. Dual angiotensin receptor and neprilysin inhibition as an alternative to angiotensin-converting enzyme inhibition in patients with chronic systolic heart failure: rationale for and design of the Prospective comparison of ARNI with ACEI to Determine Impact on Global Mortality and morbidity in Heart Failure trial (PARADIGM-HF). *Eur J Heart Fail* 2013; 15(9):1062-73.
- ^{ix} AdaptResponse clinical trial. [ClinicalTrials.gov ID NCT02205359](http://clinicaltrials.gov/ct2/show/study/NCT02205359).
<http://clinicaltrials.gov/ct2/show/study/NCT02205359>.
- ^x Singh JP, Abraham WT, Chung ES, Rogers T, Sambelashvili A, Coles JA Jr, Martin DO. Clinical response with adaptive CRT algorithm compared with CRT with echocardiography optimized atrioventricular delay: a retrospective analysis of multicentre trials. *Europace*. 2013;15(11):1622-8.
- ^{xi} Chung ES, Singh JP, Martin DO et al. Clinical and economic value of maximizing response to cardiac resynchronization therapy (CRT): Evidence from 5 randomized controlled trials. *Circulation* 2014, Issue 130, Supplement 2. Abstract No 16318. Available on-line at:
<https://aha.scientificposters.com/index.cfm?k=rtq1p07lq6>
- ^{xii} Linde C, Abraham WT, Gold MR, St John Sutton M, Ghio S, Daubert C; REVERSE (REsynchronization reVERses Remodeling in Systolic left vEntricular dysfunction) Study Group. Randomized trial of cardiac resynchronization in mildly symptomatic heart failure patients and in asymptomatic patients with left ventricular dysfunction and previous heart failure symptoms. *J Am Coll Cardiol*. 2008;52(23):1834-43.
- ^{xiii} Moss AJ, Hall WJ, Cannom DS, Klein H, Brown MW, Daubert JP, Estes NA 3rd, Foster E, Greenberg H, Higgins SL, Pfeffer MA, Solomon SD, Wilber D, Zareba W; MADIT-CRT Trial Investigators. Cardiac-resynchronization therapy for the prevention of heart-failure events. *N Engl J Med*. 2009;361(14):1329-38.
- ^{xiv} Tang AS, Wells GA, Talajic M, Arnold MO, Sheldon R, Connolly S, Hohnloser SH, Nichol G, Birnie DH, Sapp JL, Yee R, Healey JS, Rouleau JL; Resynchronization-Defibrillation for Ambulatory Heart Failure Trial Investigators. Cardiac-resynchronization therapy for mild-to-moderate heart failure. *N Engl J Med*. 2010;363(25):2385-95.

-
- ^{xv} Konstam MA, Rousseau MF, Kronenberg MW, Udelson JE, Melin J, Stewart D, Dolan N, et al. Effects of the angiotensin converting enzyme inhibitor enalapril on the long-term progression of left ventricular dysfunction in patients with heart failure. SOLVD Investigators. *Circulation*. 1992;86(2):431-8.
- ^{xvi} Abraham WT, Fisher WG, Smith AL, Delurgio DB, Leon AR, Loh E, Kocovic DZ, Packer M, Clavell AL, Hayes DL, Ellestad M, Trupp RJ, Underwood J, Pickering F, Truex C, McAtee P, Messenger J; MIRACLE Study Group. Cardiac resynchronization in chronic heart failure. *N Engl J Med*. 2002;346(24):1845-53.
- ^{xvii} Bristow MR, Saxon LA, Boehmer J, Krueger S, Kass DA, De Marco T, Carson P, et al. Cardiac-resynchronization therapy with or without an implantable defibrillator in advanced chronic heart failure. *N Engl J Med*. 2004;350(21):2140-50.
- ^{xviii} Cleland JG, Daubert JC, Erdmann E, Freemantle N, Gras D, Kappenberger L, Tavazzi L, et al. The effect of cardiac resynchronization on morbidity and mortality in heart failure. *N Engl J Med*. 2005;352(15):1539-49.
- ^{xix} Linde C, Abraham WT, Gold MR, St John Sutton M, Ghio S, Daubert C; REVERSE (REsynchronization reVERses Remodeling in Systolic left vEntricular dysfunction) Study Group. Randomized trial of cardiac resynchronization in mildly symptomatic heart failure patients and in asymptomatic patients with left ventricular dysfunction and previous heart failure symptoms. *J Am Coll Cardiol*. 2008;52(23):1834-43.
- ^{xx} Curtis AB, Worley SJ, Adamson PB, Chung ES, Niazi I, Sherfese L, Shinn T, Sutton MS; Biventricular versus Right Ventricular Pacing in Heart Failure Patients with Atrioventricular Block (BLOCK HF) Trial Investigators. Biventricular pacing for atrioventricular block and systolic dysfunction. *N Engl J Med*. 2013;368(17):1585-93.
- ^{xxi} Moss AJ, Hall WJ, Cannom DS, Klein H, Brown MW, Daubert JP, Estes NA 3rd, Foster E, Greenberg H, Higgins SL, Pfeffer MA, Solomon SD, Wilber D, Zareba W; MADIT-CRT Trial Investigators. Cardiac-resynchronization therapy for the prevention of heart-failure events. *N Engl J Med*. 2009;361(14):1329-38.
- ^{xxii} White HD, Norris RM, Brown MA, Brandt PW, Whitlock RM, Wild CJ. Left ventricular end-systolic volume as the major determinant of survival after recovery from myocardial infarction. *Circulation*. 1987;76:44-51.
- ^{xxiii} Yu CM, Bleeker GB, Fung JW, Schalij MJ, Zhang Q, van der Wall EE, Chan YS, et al. Left ventricular reverse remodeling but not clinical improvement predicts long-term survival after cardiac resynchronization therapy. *Circulation*. 2005;112(11):1580-6.
- ^{xxiv} Gold MR, Daubert C, Abraham WT, Ghio S, St. John Sutton, M, Hudnall, JH, Cerkenik J, et al. The Impact of Reverse Remodeling on Long Term Survival in Mildly Symptomatic Heart Failure Patients Receiving Cardiac Resynchronization Therapy: Results from the REVERSE study. *Heart Rhythm Journal*. 2015;12(3):524-530..
- ^{xxv} MIRACLE EF clinical trial. ClinicalTrials.gov ID NCT01735916. <http://clinicaltrials.gov/ct2/show/NCT01735916>.
- ^{xxvi} Fornwalt BK, Sprague WW, BeDell P, Suever JD, Gerritse B, Merlino JD, Fyfe DA, et al. Agreement is poor among current criteria used to define response to cardiac resynchronization therapy. *Circulation*. 2010;121(18):1985-91.
- ^{xxvii} Linde C, Abraham WT, Gold MR, St John Sutton M, Ghio S, Daubert C; REVERSE (REsynchronization reVERses Remodeling in Systolic left vEntricular dysfunction) Study Group. Randomized trial of cardiac resynchronization in mildly symptomatic heart failure patients and in asymptomatic patients with left ventricular dysfunction and previous heart failure symptoms. *J Am Coll Cardiol*. 2008;52(23):1834-43.
- ^{xxviii} Gold MR, Daubert C, Abraham WT, Ghio S, St. John Sutton, M, Hudnall, JH, Cerkenik J, et al. The Impact of Reverse Remodeling on Long Term Survival in Mildly Symptomatic Heart Failure Patients Receiving Cardiac Resynchronization Therapy: Results from the REVERSE study. *Heart Rhythm Journal*. 2015;12(3):524-530.