References

- 1. Aaronson et al., for the European Organization for Research and Treatment of Cancer QLQ-C30: A Quality-of-Life Instrument for Use in International Clinical Trials in Oncology. JNCI 1993; 85:365-376.
- 2. Anderson et al., Symptom burden in patients undergoing autologous stem-cell transplantation. Bone Marrow Transplantation. 2007; 39:759-766.
- 3. Berry et al., Enhancing patient-provider communication with the electronic self-report assessment for cancer: a randomized trial. JCO 2011; 29:1029-1035.
- 4. Berry et al., Electronic self-report assessment for cancer and self-care support: results of a multicenter randomized trial. JCO 2014; 32:199-205.
- 5. Cleeland et al., Assessing Symptom Distress in Cancer Patients: The MD Anderson Symptom Inventory. Cancer. 2000; 89:1634-1646.
- 6. Cook et al., PROMIS measures of pain, fatigue, negative affect, physical function, and social function demonstrate clinical validity across a range of chronic conditions. J Clin Epi. 2016; 73:89-102.
- 7. DeWalt et al., Evaluation of Item Candidates: The PROMIS Qualitative Item Review. Med Care. 2007; 45 (Suppl1):S12-S21.
- 8. Dueck et al., Validity and Reliability of the US NCI's PRO-CTCAE. JAMA Oncol. 2015; 1(8):1051-1059.
- 9. Kotronoulas et al., What is the value of the routine use of patient-reported outcome measures toward improvement of patient outcomes, processes of care, and health service outcomes in cancer care? A systematic review of controlled trials. JCO. 2014; 32:1480-1501.
- 10. Rogers et al., The addition of mood and anxiety domains to the university of Washington quality of life scale. Head & Neck. 2002; 24(6):521-529.
- 11. Rogers et al., The patients' account of outcome following primary surgery for oral and oropharyngeal cancer using a "quality of life" questionnaire. Eur J Cancer Care (Engl) 2008; 17(2):182-188.
- 12. Rogers et al., Screening for dysfunction to promote multidisciplinary intervention using the University of Washington Quality of Life Questionnaire. Arch Oto Head Neck Surg. 2009; 135(4):369-375.
- 13. Schipper et al., Measuring the quality of life of cancer patients: the functional living index-cancer: development and validation. JCO. 1984; 5:472-483.
- 14. Wildiers et al., End points and trial design in geriatric oncology research: a joint European organisation for research and treatment of cancer—Alliance for Clinical Trials in Oncology—International Society of Geriatric Oncology position article. JCO. 2013;31:3711-8.