

CENTER OF EXCELLENCE
FOR BEHAVIORAL HEALTH IN NURSING FACILITIES

Understanding Psychotic Disorders in Nursing Facility Residents

SUPPORTING MENTAL HEALTH WITH
COMPASSION AND KNOWLEDGE

What is a psychotic disorder?

Psychotic disorder is an umbrella term used to describe a group of mental health disorders characterized by delusions and/or hallucinations, often resulting in a loss of contact with reality. These symptoms can lead to significant emotional distress and interfere with daily functioning.



What are delusions and hallucinations?

Delusions are false, unshakeable beliefs that a person holds on to despite evidence that the belief is not real.

Common delusions that may be seen in nursing facility resident populations include:



- ❑ **Paranoid delusions**
(e.g., “people are trying to harm me, poison me”)



- ❑ **Delusions of reference**
(e.g., “people are talking about me negatively behind my back”)



- ❑ **Delusions of theft**
(e.g., “someone is stealing my belongings”)

Hallucinations are the false experiences of seeing, hearing, smelling, or feeling things that aren’t there. Hallucinations can involve any of the senses: smell, touch, hearing, vision, or taste.

Common hallucinations that may be seen in nursing facility resident populations include:



- ❑ **Visual hallucinations**
(e.g., seeing people or animals that are not there)



- ❑ **Auditory hallucinations**
(e.g., hearing voices)

Note: Delusions and hallucinations can occur briefly due to an adverse effect of medication or acute medical condition such as infection, delirium, or certain forms of dementia. These symptoms are not always part of a psychotic disorder and should be assessed to rule out underlying causes.

Common psychotic disorders seen in nursing facility residents

- ❑ Major depression with psychotic symptoms
- ❑ Bipolar disorder with psychotic symptoms
- ❑ Schizophrenia, schizoaffective disorders, and other related psychotic disorders
- ❑ Medication-induced psychotic disorder (e.g., steroid-induced psychosis)
- ❑ Substance-induced psychotic disorder (e.g., alcohol withdrawal, cannabis, methamphetamine intoxication)
- ❑ Parkinson’s-related psychosis
- ❑ Dementia-related psychosis



Common risks associated with untreated psychotic disorders

- ❑ Suicide
- ❑ Agitation
- ❑ Poor quality of life
- ❑ Worsening of symptoms
- ❑ Disruption of family relationships
- ❑ Physical aggression towards others
- ❑ Impairment in the ability to perform activities of daily living



How can staff help support residents?

- ❑ **Recognize early warning signs:** Watch for subtle changes in behavior or thinking that may indicate the onset of delusions or hallucinations.
- ❑ **Clarify and confirm symptoms:** Gently explore the resident's concerns and check the facts. For instance, if they report theft, verify whether items are actually missing.
- ❑ **Offer reassurance:** Let the resident know they are not alone, and support is available.
- ❑ **Identify and treat underlying causes:** Assess contributing factors such as substance use, medication side effects or other underlying medical conditions.
- ❑ **Apply person-centered interventions:** Use individualized non-pharmacological interventions to help reduce distress and promote safety and enhance quality of life. Include interventions shared by the family that have helped with symptom management in the past.
- ❑ **Communicate with the care team:** Report observations to the interdisciplinary team, including the physician or advanced practice provider, to ensure coordinated and responsive care.



Treatment of psychotic disorders

Psychotic disorders are highly treatable with a combination of evidence-based, comprehensive interventions.

- ❑ **Psychoeducation:** Ensure residents, family, and staff have basic knowledge about the disorder and treatment options.
- ❑ **Person-centered care plan:** Engage the resident in creating a care plan that reflects their strengths, abilities, preferences, and needs.
- ❑ **Non-pharmacological interventions:** Use individualized approaches to reduce distress and enhance daily functioning, including psychosocial skills training.
- ❑ **Psychotherapy:** Coordinate therapeutic support such as cognitive behavioral therapy or other relevant modalities to address symptoms and promote recovery.
- ❑ **Medications management:** Consider pharmacological treatment if clinically indicated, in consultation with the care team.



References

Understanding Psychosis. National Institute of Mental Health (NIMH).
 National Alliance on Mental Illness (NAMI)
 Untreated Psychosis.



This material was created by the Center of Excellence for Behavioral Health in Nursing Facilities. This work is made possible by grant number 1H79SM087155 from the Substance Abuse and Mental Health Services Administration (SAMHSA). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Substance Abuse and Mental Health Services Administration. 8/2025