

## 2016 Ophthalmology Preferred Specialty Measure Set

PQRS#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
12	0086	Claims, Registry, EHR	Effective Clinical Care	<b>Primary Open-Angle Glaucoma (POAG): Optic Nerve Evaluation:</b> Percentage of patients aged 18 years and older with a diagnosis of primary open-angle glaucoma (POAG) who have an optic nerve head evaluation during one or more office visits within 12 months
14	0087	Claims, Registry	Effective Clinical Care	<b>Age-Related Macular Degeneration (AMD): Dilated Macular Examination:</b> Percentage of patients aged 50 years and older with a diagnosis of age-related macular degeneration (AMD) who had a dilated macular examination performed which included documentation of the presence or absence of macular thickening or hemorrhage AND the level of macular degeneration severity during one or more office visits within 12 months
18	0088	EHR, Measures Groups	Effective Clinical Care	<b>Diabetic Retinopathy: Documentation of Presence or Absence of Macular Edema and Level of Severity of Retinopathy:</b> Percentage of patients aged 18 years and older with a diagnosis of diabetic retinopathy who had a dilated macular or fundus exam performed which included documentation of the level of severity of retinopathy and the presence or absence of macular edema during one or more office visits within 12 months
19	0089	Claims, Registry, EHR, Measures Groups	Communication and Care Coordination	<b>Diabetic Retinopathy: Communication with the Physician Managing Ongoing Diabetes Care:</b> Percentage of patients aged 18 years and older with a diagnosis of diabetic retinopathy who had a dilated macular or fundus exam performed with documented communication to the physician who manages the ongoing care of the patient with diabetes mellitus regarding the findings of the macular or fundus exam at least once within 12 months
117	0055	Registry, EHR, GPRO (Web Interface), Measures Groups	Effective Clinical Care	<b>Diabetes: Eye Exam:</b> Percentage of patients 18 - 75 years of age with diabetes who had a retinal or dilated eye exam by an eye care professional during the measurement period or a negative retinal or dilated eye exam (no evidence of retinopathy) in the 12 months prior to the measurement period

2016 Ophthalmology Preferred Specialty Measure Set

PQRS#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
140	0566	Claims, Registry	Effective Clinical Care	<b>Age-Related Macular Degeneration (AMD): Counseling on Antioxidant Supplement:</b> Percentage of patients aged 50 years and older with a diagnosis of age-related macular degeneration (AMD) or their caregiver(s) who were counseled within 12 months on the benefits and/or risks of the Age-Related Eye Disease Study (AREDS) formulation for preventing progression of AMD
141	0563	Claims, Registry	Communication and Care Coordination	<b>Primary Open-Angle Glaucoma (POAG): Reduction of Intraocular Pressure (IOP) by 15% OR Documentation of a Plan of Care:</b> Percentage of patients aged 18 years and older with a diagnosis of primary open-angle glaucoma (POAG) whose glaucoma treatment has not failed (the most recent IOP was reduced by at least 15% from the pre-intervention level) OR if the most recent IOP was not reduced by at least 15% from the pre-intervention level, a plan of care was documented within 12 months
191	0565	Registry, EHR, Measures Groups	Effective Clinical Care	<b>Cataracts: 20/40 or Better Visual Acuity within 90 Days Following Cataract Surgery:</b> Percentage of patients aged 18 years and older with a diagnosis of uncomplicated cataract who had cataract surgery and no significant ocular conditions impacting the visual outcome of surgery and had best-corrected visual acuity of 20/40 or better (distance or near) achieved within 90 days following the cataract surgery
192	0564	Registry, EHR, Measures Groups	Patient Safety	<b>Cataracts: Complications within 30 Days Following Cataract Surgery Requiring Additional Surgical Procedures:</b> Percentage of patients aged 18 years and older with a diagnosis of uncomplicated cataract who had cataract surgery and had any of a specified list of surgical procedures in the 30 days following cataract surgery which would indicate the occurrence of any of the following major complications: retained nuclear fragments, endophthalmitis, dislocated or wrong power IOL, retinal detachment, or wound dehiscence

2016 Ophthalmology Preferred Specialty Measure Set

PQRS#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
226	0028	Claims, Registry, EHR, GPRO (Web Interface), Measures Groups	Community/ Population Health	<b>Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention:</b> Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 24 months AND who received cessation counseling intervention if identified as a tobacco user
303	1536	Registry, Measures Groups	Patient and Caregiver-Centered Experience and Outcomes	<b>Cataracts: Improvement in Patient's Visual Function within 90 Days Following Cataract Surgery:</b> Percentage of patients aged 18 years and older in sample who had cataract surgery and had improvement in visual function achieved within 90 days following the cataract surgery, based on completing a pre-operative and post-operative visual function survey
304	N/A	Registry, Measures Groups	Patient and Caregiver-Centered Experience and Outcomes	<b>Cataracts: Patient Satisfaction within 90 Days Following Cataract Surgery:</b> Percentage of patients aged 18 years and older in sample who had cataract surgery and were satisfied with their care within 90 days following the cataract surgery, based on completion of the Consumer Assessment of Healthcare Providers and Systems Surgical Care Survey
384	N/A	Registry	Effective Clinical Care	<b>Adult Primary Rhegmatogenous Retinal Detachment Surgery: No Return to the Operating Room Within 90 Days of Surgery:</b> Patients aged 18 years and older who had surgery for primary rhegmatogenous retinal detachment who did not require a return to the operating room within 90 days of surgery
385	N/A	Registry	Effective Clinical Care	<b>Adult Primary Rhegmatogenous Retinal Detachment Surgery: Visual Acuity Improvement Within 90 Days of Surgery:</b> Patients aged 18 years and older who had surgery for primary rhegmatogenous retinal detachment and achieved an improvement in their visual acuity, from their preoperative level, within 90 days of surgery in the operative eye
388	N/A	Registry, Measures Groups	Patient Safety	<b>Cataract Surgery with Intra-Operative Complications (Unplanned Rupture of Posterior Capsule Requiring Unplanned Vitrectomy):</b> Percentage of patients aged 18 years and older who had cataract surgery performed and had an unplanned rupture of the posterior capsule requiring vitrectomy

## 2016 Ophthalmology Preferred Specialty Measure Set

PQRS#	NQF#	Reporting Method	National Quality Strategy Domain	Measure Title: Description
389	N/A	Registry, Measures Groups	Effective Clinical Care	<b>Cataract Surgery: Difference Between Planned and Final Refraction:</b> Percentage of patients aged 18 years and older who had cataract surgery performed and achieved a final refraction within +/-1,0 diopters of their planned (target) refraction

This table is a resource of potential measure that may be applicable to Ophthalmology. Please note that in addition to this table, a cross cutting measures list is also available for reporting at the following link: [2016 Cross-Cutting Measures List](#).