

Table 7. Part B therapy per patient expenditures, by setting and patient diagnosis, 1998-2000 (page 1 of 2)

Condition	Average therapy expenditures ¹									
	% of patients, all settings	All settings	Hospital outpatient	SNF inpatient	SNF outpatient	Rehabilitation agency	CORF	Independent practice	Physician practice	Other/ multiple
All										
1998	100.0%	\$709	\$286	\$1,299	\$1,350	\$962	\$1,642	\$361	\$230	\$1,383
1999	100.0	480	389	554	525	525	743	449	331	822
2000	100.0	642	432	838	769	728	1,199	607	335	1,238
Percent change, 1998-99		-32.3%	35.9%	-57.4%	-61.1%	-45.4%	-54.8%	24.4%	43.9%	-40.6%
Percent change, 1999-00		33.8%	11.1%	51.3%	46.5%	38.7%	61.4%	35.2%	1.2%	50.6%
Average annual change, 1998-00		-4.8%	22.9%	-19.7%	-24.5%	-13.0%	-14.5%	29.7%	20.7%	-5.4%
Musculoskeletal/soft tissue										
1998	32.1	590	280	1,371	1,343	800	1,369	364	239	1,023
1999	32.3	477	362	584	521	509	657	451	393	769
2000	32.0	640	418	872	791	696	1,081	614	385	1,223
Percent change, 1998-99		-19.2%	29.1%	-57.4%	-61.2%	-36.4%	-52.0%	23.9%	64.4%	-24.8%
Percent change, 1999-00		34.2%	15.6%	49.3%	51.8%	36.7%	64.5%	36.1%	-2.0%	59.0%
Average annual change, 1998-00		4.2%	22.1%	-20.2%	-23.3%	-6.7%	-11.1%	29.9%	26.9%	9.3%
Back disorder										
1998	18.5	465	249	1,189	997	616	1,132	349	351	887
1999	19.0	451	374	516	451	452	663	421	456	804
2000	19.3	605	377	772	583	603	1,046	561	484	1,270
Percent change, 1998-99		-3.0%	50.0%	-56.6%	-54.8%	-26.6%	-41.4%	20.6%	29.9%	-9.4%
Percent change, 1999-00		34.1%	0.7%	49.6%	29.3%	33.4%	57.8%	33.3%	6.1%	58.0%
Average annual change, 1998-00		14.1%	22.9%	-19.4%	-23.5%	-1.1%	-3.9%	26.8%	17.4%	19.7%
Neurologic										
1998	15.8	971	209	1,191	1,297	1,415	2,271	343	137	2,061
1999	15.4	461	341	506	502	599	967	439	172	929
2000	15.3	634	321	734	729	848	1,601	623	174	1,288
Percent change, 1998-99		-52.5%	63.0%	-57.5%	-61.3%	-57.7%	-57.4%	28.0%	25.5%	-54.9%
Percent change, 1999-00		37.5%	-5.9%	45.1%	45.2%	41.6%	65.6%	41.9%	1.2%	38.6%
Average annual change, 1998-00		-19.2%	23.8%	-21.5%	-25.0%	-22.6%	-16.0%	34.8%	12.7%	-20.9%
Unspecified rehabilitation										
1998	7.7	367	294	1,058	810	694	1,307	366	135	1,117
1999	8.9	434	409	539	559	383	488	517	132	930
2000	9.9	486	443	649	586	337	594	539	303	1,139
Percent change, 1998-99		18.3%	38.7%	-49.1%	-31.0%	-44.8%	-62.7%	41.3%	-2.2%	-16.7%
Percent change, 1999-00		12.0%	8.6%	20.4%	4.8%	-12.0%	21.7%	4.3%	129.5%	22.5%
Average annual change, 1998-00		15.1%	22.7%	-21.7%	-14.9%	-30.3%	-32.6%	21.4%	49.8%	1.0%
Ortho-surgical										
1998	4.8	656	352	1,556	1,379	928	1,605	399	320	818
1999	5.0	534	471	680	610	564	628	545	347	690
2000	4.8	711	528	1,023	876	836	845	715	363	1,063
Percent change, 1998-99		-18.6%	33.9%	-56.3%	-55.8%	-39.2%	-60.9%	36.6%	8.4%	-15.6%
Percent change, 1999-00		33.1%	12.1%	50.4%	43.6%	48.2%	34.6%	31.2%	4.6%	54.1%
Average annual change, 1998-00		4.1%	22.5%	-18.9%	-20.3%	-5.1%	-27.4%	33.9%	6.5%	14.0%

Table 7. Part B therapy per patient expenditures, by setting and patient diagnosis, 1998-2000 (page 2 of 2)

Condition	% of patients, all settings	Average therapy expenditures ¹								
		All settings	Hospital outpatient	SNF inpatient	SNF outpatient	Rehabilitation agency	CORF	Independent practice	Physician office	Other/ multiple
Stroke										
1998	4.8	1,489	564	1,529	1,697	2,053	2,804	417	342	2,809
1999	4.3	809	943	613	600	769	1,057	561	361	1,343
2000	4.2	1,140	974	1,002	925	1,315	1,849	873	587	1,963
Percent change, 1998-99		-45.7%	67.3%	-59.9%	-64.6%	-62.5%	-62.3%	34.5%	5.6%	-52.2%
Percent change, 1999-00		40.9%	3.3%	63.5%	54.2%	71.0%	74.9%	55.6%	62.6%	46.2%
Average annual change, 1998-00		-12.5%	31.5%	-19.0%	-26.2%	-20.0%	-18.8%	44.7%	31.0%	-16.4%
Cardiovascular										
1998	2.9	953	246	1,223	1,278	1,701	1,829	481	105	1,628
1999	2.7	455	272	563	538	684	972	318	140	757
2000	2.6	652	315	881	760	1,089	1,282	1,031	154	966
Percent change, 1998-99		-52.3%	10.9%	-54.0%	-57.9%	-59.8%	-46.9%	-33.9%	33.3%	-53.5%
Percent change, 1999-00		43.3%	15.7%	56.5%	41.3%	59.2%	31.9%	224.2%	10.0%	27.6%
Average annual change, 1998-00		-17.3%	13.3%	-15.1%	-22.9%	-20.0%	-16.3%	46.4%	21.1%	-23.0%
Hip fracture										
1998	1.6	1,352	364	1,354	1,415	1,818	2,271	419	339	2,205
1999	1.4	653	452	567	557	785	866	588	508	1,079
2000	1.3	930	609	910	837	1,214	1,598	727	418	1,393
Percent change, 1998-99		-51.7%	24.1%	-58.1%	-60.6%	-56.8%	-61.9%	40.3%	49.9%	-51.1%
Percent change, 1999-00		42.4%	34.9%	60.5%	50.3%	54.6%	84.5%	23.6%	-17.7%	29.1%
Average annual change, 1998-00		-17.1%	29.4%	-18.0%	-23.1%	-18.3%	-16.1%	31.7%	11.0%	-20.5%
Amputation/deformity										
1998	0.5	1,015	383	1,550	1,529	1,237	3,042	303	173	1,882
1999	0.4	532	420	732	605	490	716	384	198	886
2000	0.4	768	530	1,063	777	751	903	456	406	1,838
Percent change, 1998-99		-47.6%	9.7%	-52.8%	-60.4%	-60.4%	-76.5%	26.7%	14.5%	-52.9%
Percent change, 1999-00		44.4%	26.1%	45.2%	28.4%	53.3%	26.1%	18.8%	105.1%	107.4%
Average annual change, 1998-00		-13.0%	17.6%	-17.2%	-28.7%	-22.1%	-45.5%	22.7%	53.2%	-1.2%
Spinal cord										
1998	0.3	823	479	2,108	754	1,049	1,021	446	556	1,349
1999	0.3	664	567	609	277	794	1,018	639	911	930
2000	0.3	836	495	1,177	808	1,240	2,696	1,164	181	1,421
Percent change, 1998-99		-19.3%	18.3%	-71.1%	-63.3%	-24.3%	-0.3%	43.3%	63.8%	-31.1%
Percent change, 1999-00		25.9%	-12.7%	93.3%	191.7%	56.2%	164.8%	82.2%	-80.1%	52.8%
Average annual change, 1998-00		0.8%	1.6%	-25.3%	3.5%	8.7%	62.5%	61.6%	-42.9%	2.6%
All other diagnoses										
1998	10.9	841	288	1,288	1,380	1,470	2,615	332	103	1,560
1999	10.2	422	369	558	530	616	723	410	138	672
2000	9.8	596	369	865	775	813	1,127	546	145	1,018
Percent change, 1998-99		-49.8%	28.1%	-56.7%	-61.6%	-58.1%	-72.4%	23.5%	34.0%	-56.9%
Percent change, 1999-00		41.2%	0.2%	55.0%	46.2%	32.0%	55.9%	33.2%	5.1%	51.5%
Average annual change, 1998-00		-15.8%	13.3%	-18.0%	-25.1%	-25.6%	-34.4%	28.2%	18.6%	-19.2%

¹ Expenditures reflect Medicare program payments (this excludes patients' 20 percent coinsurance). Note: SNF is skilled nursing facility. CORF is comprehensive outpatient rehabilitation facility.

"Other/multiple" is hospital inpatient, home health agency outpatient, ambulatory surgical centers, and multiple (use of two or more settings).

Source: Urban Institute analysis of 5% samples of Medicare claims, 1998-2000.