

Treatment Resistant Depression

**Medicare Evidence Development and Coverage
Advisory Committee (MEDCAC)**

April 27, 2016



Purpose of Meeting

- **Obtain MEDCAC recommendations regarding**
 1. Definition of TRD
 2. Advise CMS on the use of the definition of TRD in the context of clinical studies, coverage with evidence development, and treatment outcomes

Voting Question # 1

- **How confident are you that there is a standard definition of TRD that can be applied to Medicare beneficiaries in clinical studies of therapies for this disease?**

Use the following scale identifying your level of confidence - with a score of 1 being low or no confidence and 5 representing high confidence.

1 — 2 — 3 — 4 — 5
Low Confidence Intermediate High Confidence

Voting Question # 2

- **If intermediate confidence (≥ 2.5) is noted for Question #1, please vote by “yes or no” as to whether the following are important defining characteristics of TRD that are to be considered in clinical research:**
 - a) The number, duration, dosage, and/or classes of antidepressants attempted
 - b) The use of augmentation/combination pharmacological therapies
 - c) Type of depressive episode (unipolar, bipolar, psychotic, atypical, other)
 - d) The use of nonpharmacological treatments such as electroconvulsive therapy
 - e) The use of psychotherapy
 - f) Score changes on standardized and validated depression rating instruments (e.g. Hamilton Depression Rating Scale)
 - g) Suicidal ideation and suicide attempts
 - h) Other

Question # 4 - Discussion

- **For each characteristic in Question #4 that receives intermediate confidence (≥ 2.5), please discuss the a priori parameters that define successful or failed treatment.**

- **Characteristics listed in Question #4**
 - a) Improvement or decline in function
 - b) Improvement or decline in quality of life
 - c) Decrease in suicide ideation
 - d) Decrease in suicidal attempts
 - e) Other

Voting Question # 5

- **How confident are you that the strategies below, when applied to Medicare beneficiaries, represent meaningful and realistic study designs in research investigations performed to evaluate interventions for TRD?**
 - a) Randomized sham-controlled double blinded trial
 - b) Randomized sham-controlled single blinded trial
 - c) Randomized controlled unblinded trial
 - d) Randomized crossover study
 - e) Nonrandomized crossover study
 - f) Pre/post study design
 - g) Other

Use the following scale identifying your level of confidence - with a score of 1 being low or no confidence and 5 representing high confidence.

1 — 2 — 3 — 4 — 5
Low — **Intermediate** — **High**
Confidence — **Confidence**