

Cognitive improvement following bariatric surgery

John Gunstad
Department of Psychological Sciences
Applied Psychology Center

Disclosures


- Research support (>\$10,000) from:
 - NIH, iHealth, IBM, Dairy Research Council
- No intellectual conflicts of interest

Neurological outcomes

- Mid-life obesity associated with stroke, Alzheimer's disease, vascular dementia, etc.


(Gustafson et al., 2012; Fitzpatrick et al., 2009)

- Obesity is associated with accelerated cognitive decline




Severe obesity and cognition

Age X
severity
interaction
for
executive
function




MCI in severe obesity



A majority of persons with severe obesity met criteria for MCI

LABS ancillary study


- Longitudinal Assessment of Bariatric Surgery (LABS)
- Cognitive testing before and after surgery
 - Baseline
 - 12 week f/u
 - 12-, 24-, and 36-month f/u



No cognitive side effects


<u>Test</u>	<u>>1.5 SD Decline</u>	<u>Differ from Controls?</u>
Learning	6.7%	No
Digit Span	9.5%	No
Switch Attn— Let/Num	0.9%	No
Animal Fluency	5.3%	No

Post-op improvement



Pilot data
suggests
benefits
persist to 7
years

Benefits in older adults



Cognitive benefits were similar in patients <55> years of age